

Instructor's Solution Manual  
Introduction to Electrodynamics  
Fourth Edition

David J. Griffiths

2014

# Contents

<b>1</b>	<b>Vector Analysis</b>	<b>4</b>
<b>2</b>	<b>Electrostatics</b>	<b>26</b>
<b>3</b>	<b>Potential</b>	<b>53</b>
<b>4</b>	<b>Electric Fields in Matter</b>	<b>92</b>
<b>5</b>	<b>Magnetostatics</b>	<b>110</b>
<b>6</b>	<b>Magnetic Fields in Matter</b>	<b>133</b>
<b>7</b>	<b>Electrodynamics</b>	<b>145</b>
<b>8</b>	<b>Conservation Laws</b>	<b>168</b>
<b>9</b>	<b>Electromagnetic Waves</b>	<b>185</b>
<b>10</b>	<b>Potentials and Fields</b>	<b>210</b>
<b>11</b>	<b>Radiation</b>	<b>231</b>
<b>12</b>	<b>Electrodynamics and Relativity</b>	<b>262</b>



# Preface

Although I wrote these solutions, much of the typesetting was done by Jonah Gollub, Christopher Lee, and James Terwilliger (any mistakes are, of course, entirely their fault). Chris also did many of the figures, and I would like to thank him particularly for all his help. If you find errors, please let me know ([griffith@reed.edu](mailto:griffith@reed.edu)).

David Griffiths

# Chapter 1

## Vector Analysis

### Problem 1.1

- (a) From the diagram,  $|\mathbf{B} + \mathbf{C}| \cos \theta_3 = |\mathbf{B}| \cos \theta_1 + |\mathbf{C}| \cos \theta_2$ . Multiply by  $|\mathbf{A}|$ .

$$|\mathbf{A}| |\mathbf{B} + \mathbf{C}| \cos \theta_3 = |\mathbf{A}| |\mathbf{B}| \cos \theta_1 + |\mathbf{A}| |\mathbf{C}| \cos \theta_2.$$


So:  $\mathbf{A} \cdot (\mathbf{B} + \mathbf{C}) = \mathbf{A} \cdot \mathbf{B} + \mathbf{A} \cdot \mathbf{C}$ . (Dot product is distributive)

Similarly:  $|\mathbf{B} + \mathbf{C}| \sin \theta_3 = |\mathbf{B}| \sin \theta_1 + |\mathbf{C}| \sin \theta_2$ . Multiply by  $|\mathbf{A}| \hat{\mathbf{n}}$ .

$$|\mathbf{A}| |\mathbf{B} + \mathbf{C}| \sin \theta_3 \hat{\mathbf{n}} = |\mathbf{A}| |\mathbf{B}| \sin \theta_1 \hat{\mathbf{n}} + |\mathbf{A}| |\mathbf{C}| \sin \theta_2 \hat{\mathbf{n}}.$$

If  $\hat{\mathbf{n}}$  is the unit vector pointing out of the page, it follows that


$$\mathbf{A} \times (\mathbf{B} + \mathbf{C}) = (\mathbf{A} \times \mathbf{B}) + (\mathbf{A} \times \mathbf{C}). \quad (\text{Cross product is distributive})$$



- (b) For the general case, see G. E. Hay's *Vector and Tensor Analysis*, Chapter 1, Section 7 (dot product) and Section 8 (cross product)

### Problem 1.2

The triple cross-product is *not* in general associative. For example, suppose  $\mathbf{A} = \mathbf{B}$  and  $\mathbf{C}$  is perpendicular to  $\mathbf{A}$ , as in the diagram. Then  $(\mathbf{B} \times \mathbf{C})$  points out-of-the-page, and  $\mathbf{A} \times (\mathbf{B} \times \mathbf{C})$  points *down*, and has magnitude  $ABC$ . But  $(\mathbf{A} \times \mathbf{B}) = \mathbf{0}$ , so  $(\mathbf{A} \times \mathbf{B}) \times \mathbf{C} = \mathbf{0} \neq \mathbf{A} \times (\mathbf{B} \times \mathbf{C})$ .




### Problem 1.3

$$\mathbf{A} = +1\hat{\mathbf{x}} + 1\hat{\mathbf{y}} - 1\hat{\mathbf{z}}; A = \sqrt{3}; \mathbf{B} = 1\hat{\mathbf{x}} + 1\hat{\mathbf{y}} + 1\hat{\mathbf{z}}; B = \sqrt{3}.$$

$$\mathbf{A} \cdot \mathbf{B} = +1 + 1 - 1 = 1 = AB \cos \theta = \sqrt{3}\sqrt{3} \cos \theta \Rightarrow \cos \theta = \frac{1}{3}.$$

$$\theta = \cos^{-1}\left(\frac{1}{3}\right) \approx 70.5288^\circ$$



### Problem 1.4

The cross-product of any two vectors in the plane will give a vector perpendicular to the plane. For example, we might pick the base ( $\mathbf{A}$ ) and the left side ( $\mathbf{B}$ ):

$$\mathbf{A} = -1\hat{\mathbf{x}} + 2\hat{\mathbf{y}} + 0\hat{\mathbf{z}}; \mathbf{B} = -1\hat{\mathbf{x}} + 0\hat{\mathbf{y}} + 3\hat{\mathbf{z}}.$$

$$\mathbf{A} \times \mathbf{B} = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ -1 & 2 & 0 \\ -1 & 0 & 3 \end{vmatrix} = 6\hat{\mathbf{x}} + 3\hat{\mathbf{y}} + 2\hat{\mathbf{z}}.$$

This has the right *direction*, but the wrong *magnitude*. To make a *unit* vector out of it, simply divide by its length:

$$|\mathbf{A} \times \mathbf{B}| = \sqrt{36 + 9 + 4} = 7. \quad \hat{\mathbf{n}} = \frac{\mathbf{A} \times \mathbf{B}}{|\mathbf{A} \times \mathbf{B}|} = \left[ \frac{6}{7}\hat{\mathbf{x}} + \frac{3}{7}\hat{\mathbf{y}} + \frac{2}{7}\hat{\mathbf{z}} \right].$$

**Problem 1.5**

$$\begin{aligned} \mathbf{A} \times (\mathbf{B} \times \mathbf{C}) &= \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ A_x & A_y & A_z \\ (B_y C_z - B_z C_y) & (B_z C_x - B_x C_z) & (B_x C_y - B_y C_x) \end{vmatrix} \\ &= \hat{\mathbf{x}}[A_y(B_x C_y - B_y C_x) - A_z(B_z C_x - B_x C_z)] + \hat{\mathbf{y}}(\dots) + \hat{\mathbf{z}}(\dots) \\ &\quad \text{(I'll just check the x-component; the others go the same way)} \\ &= \hat{\mathbf{x}}(A_y B_x C_y - A_y B_y C_x - A_z B_z C_x + A_z B_x C_z) + \hat{\mathbf{y}}(\dots) + \hat{\mathbf{z}}(\dots). \\ \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) &= [B_x(A_x C_x + A_y C_y + A_z C_z) - C_x(A_x B_x + A_y B_y + A_z B_z)]\hat{\mathbf{x}} + (\dots)\hat{\mathbf{y}} + (\dots)\hat{\mathbf{z}} \\ &= \hat{\mathbf{x}}(A_y B_x C_y + A_z B_x C_z - A_y B_y C_x - A_z B_z C_x) + \hat{\mathbf{y}}(\dots) + \hat{\mathbf{z}}(\dots). \text{ They agree.} \end{aligned}$$

**Problem 1.6**

$$\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) + \mathbf{B} \times (\mathbf{C} \times \mathbf{A}) + \mathbf{C} \times (\mathbf{A} \times \mathbf{B}) = \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) + \mathbf{C}(\mathbf{A} \cdot \mathbf{B}) - \mathbf{A}(\mathbf{C} \cdot \mathbf{B}) + \mathbf{A}(\mathbf{B} \cdot \mathbf{C}) - \mathbf{B}(\mathbf{C} \cdot \mathbf{A}) = \mathbf{0}.$$

So:  $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) - (\mathbf{A} \times \mathbf{B}) \times \mathbf{C} = -\mathbf{B} \times (\mathbf{C} \times \mathbf{A}) = \mathbf{A}(\mathbf{B} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B}).$

If this is zero, then either  $\mathbf{A}$  is parallel to  $\mathbf{C}$  (including the case in which they point in *opposite* directions, or one is zero), or else  $\mathbf{B} \cdot \mathbf{C} = \mathbf{B} \cdot \mathbf{A} = 0$ , in which case  $\mathbf{B}$  is perpendicular to  $\mathbf{A}$  and  $\mathbf{C}$  (including the case  $\mathbf{B} = \mathbf{0}$ .)

*Conclusion:*  $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = (\mathbf{A} \times \mathbf{B}) \times \mathbf{C} \iff$  either  $\mathbf{A}$  is parallel to  $\mathbf{C}$ , or  $\mathbf{B}$  is perpendicular to  $\mathbf{A}$  and  $\mathbf{C}$ .

**Problem 1.7**

$$\mathbf{r} = (4\hat{\mathbf{x}} + 6\hat{\mathbf{y}} + 8\hat{\mathbf{z}}) - (2\hat{\mathbf{x}} + 8\hat{\mathbf{y}} + 7\hat{\mathbf{z}}) = \boxed{2\hat{\mathbf{x}} - 2\hat{\mathbf{y}} + \hat{\mathbf{z}}}$$

$$r = \sqrt{4 + 4 + 1} = \boxed{3}$$

$$\hat{\mathbf{r}} = \frac{\mathbf{r}}{r} = \boxed{\frac{2}{3}\hat{\mathbf{x}} - \frac{2}{3}\hat{\mathbf{y}} + \frac{1}{3}\hat{\mathbf{z}}}$$


**Problem 1.8**

$$\begin{aligned} \text{(a)} \quad \bar{A}_y \bar{B}_y + \bar{A}_z \bar{B}_z &= (\cos \phi A_y + \sin \phi A_z)(\cos \phi B_y + \sin \phi B_z) + (-\sin \phi A_y + \cos \phi A_z)(-\sin \phi B_y + \cos \phi B_z) \\ &= \cos^2 \phi A_y B_y + \sin \phi \cos \phi (A_y B_z + A_z B_y) + \sin^2 \phi A_z B_z + \sin^2 \phi A_y B_y - \sin \phi \cos \phi (A_y B_z + A_z B_y) + \cos^2 \phi A_z B_z \\ &= (\cos^2 \phi + \sin^2 \phi) A_y B_y + (\sin^2 \phi + \cos^2 \phi) A_z B_z = A_y B_y + A_z B_z. \quad \checkmark \end{aligned}$$


$$\text{(b)} \quad (\bar{A}_x)^2 + (\bar{A}_y)^2 + (\bar{A}_z)^2 = \sum_{i=1}^3 \bar{A}_i \bar{A}_i = \sum_{i=1}^3 (\sum_{j=1}^3 R_{ij} A_j) (\sum_{k=1}^3 R_{ik} A_k) = \sum_{j,k} (\sum_i R_{ij} R_{ik}) A_j A_k.$$

$$\text{This equals } A_x^2 + A_y^2 + A_z^2 \text{ provided } \sum_{i=1}^3 R_{ij} R_{ik} = \begin{cases} 1 & \text{if } j = k \\ 0 & \text{if } j \neq k \end{cases}$$

Moreover, if  $R$  is to preserve lengths for *all* vectors  $\mathbf{A}$ , then this condition is not only *sufficient* but also *necessary*. For suppose  $\mathbf{A} = (1, 0, 0)$ . Then  $\sum_{j,k} (\sum_i R_{ij} R_{ik}) A_j A_k = \sum_i R_{i1} R_{i1}$ , and this must equal 1 (since we want  $\bar{A}_x^2 + \bar{A}_y^2 + \bar{A}_z^2 = 1$ ). Likewise,  $\sum_{i=1}^3 R_{i2} R_{i2} = \sum_{i=1}^3 R_{i3} R_{i3} = 1$ . To check the case  $j \neq k$ , choose  $\mathbf{A} = (1, 1, 0)$ . Then we want  $2 = \sum_{j,k} (\sum_i R_{ij} R_{ik}) A_j A_k = \sum_i R_{i1} R_{i1} + \sum_i R_{i2} R_{i2} + \sum_i R_{i1} R_{i2} + \sum_i R_{i2} R_{i1}$ . But we already know that the first two sums are both 1; the third and fourth are *equal*, so  $\sum_i R_{i1} R_{i2} = \sum_i R_{i2} R_{i1} = 0$ , and so on for other unequal combinations of  $j, k$ .  $\checkmark$  In matrix notation:  $\bar{R}R = 1$ , where  $\bar{R}$  is the *transpose* of  $R$ .

**Problem 1.9**

Looking down the axis:



A  $120^\circ$  rotation carries the  $z$  axis into the  $y$  ( $= \bar{z}$ ) axis,  $y$  into  $x$  ( $= \bar{y}$ ), and  $x$  into  $z$  ( $= \bar{x}$ ). So  $\bar{A}_x = A_z$ ,  $\bar{A}_y = A_x$ ,  $\bar{A}_z = A_y$ .

$$R = \begin{pmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}$$

**Problem 1.10**

(a) No change. ( $\bar{A}_x = A_x$ ,  $\bar{A}_y = A_y$ ,  $\bar{A}_z = A_z$ )

(b)  $\mathbf{A} \longrightarrow -\mathbf{A}$ , in the sense ( $\bar{A}_x = -A_x$ ,  $\bar{A}_y = -A_y$ ,  $\bar{A}_z = -A_z$ )

(c)  $(\mathbf{A} \times \mathbf{B}) \longrightarrow (-\mathbf{A}) \times (-\mathbf{B}) = (\mathbf{A} \times \mathbf{B})$ . That is, if  $\mathbf{C} = \mathbf{A} \times \mathbf{B}$ ,  $\boxed{\mathbf{C} \longrightarrow \mathbf{C}}$ . No minus sign, in contrast to behavior of an “ordinary” vector, as given by (b). If  $\mathbf{A}$  and  $\mathbf{B}$  are *pseudovectors*, then  $(\mathbf{A} \times \mathbf{B}) \longrightarrow (\mathbf{A}) \times (\mathbf{B}) = (\mathbf{A} \times \mathbf{B})$ . So the cross-product of two pseudovectors is again a *pseudovector*. In the cross-product of a vector and a pseudovector, one changes sign, the other doesn’t, and therefore the cross-product is itself a *vector*. *Angular momentum* ( $\mathbf{L} = \mathbf{r} \times \mathbf{p}$ ) and *torque* ( $\mathbf{N} = \mathbf{r} \times \mathbf{F}$ ) are pseudovectors.

(d)  $\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) \longrightarrow (-\mathbf{A}) \cdot ((-\mathbf{B}) \times (-\mathbf{C})) = -\mathbf{A} \cdot (\mathbf{B} \times \mathbf{C})$ . So, if  $a = \mathbf{A} \cdot (\mathbf{B} \times \mathbf{C})$ , then  $\boxed{a \longrightarrow -a}$ ; a pseudoscalar *changes sign* under inversion of coordinates.

**Problem 1.11**

$$(a) \nabla f = 2x \hat{\mathbf{x}} + 3y^2 \hat{\mathbf{y}} + 4z^3 \hat{\mathbf{z}}$$

$$(b) \nabla f = 2xy^3z^4 \hat{\mathbf{x}} + 3x^2y^2z^4 \hat{\mathbf{y}} + 4x^2y^3z^3 \hat{\mathbf{z}}$$

$$(c) \nabla f = e^x \sin y \ln z \hat{\mathbf{x}} + e^x \cos y \ln z \hat{\mathbf{y}} + e^x \sin y (1/z) \hat{\mathbf{z}}$$

**Problem 1.12**

(a)  $\nabla h = 10[(2y - 6x - 18) \hat{\mathbf{x}} + (2x - 8y + 28) \hat{\mathbf{y}}]$ .  $\nabla h = 0$  at summit, so

$$\left. \begin{aligned} 2y - 6x - 18 &= 0 \\ 2x - 8y + 28 &= 0 \end{aligned} \right\} \begin{aligned} 6x - 24y + 84 &= 0 \\ 2y - 18 - 24y + 84 &= 0. \end{aligned}$$

$$22y = 66 \implies y = 3 \implies 2x - 24 + 28 = 0 \implies x = -2.$$

Top is  $\boxed{3 \text{ miles north, 2 miles west, of South Hadley.}}$

(b) Putting in  $x = -2$ ,  $y = 3$ :

$$h = 10(-12 - 12 - 36 + 36 + 84 + 12) = \boxed{720 \text{ ft.}}$$

(c) Putting in  $x = 1$ ,  $y = 1$ :  $\nabla h = 10[(2 - 6 - 18) \hat{\mathbf{x}} + (2 - 8 + 28) \hat{\mathbf{y}}] = 10(-22 \hat{\mathbf{x}} + 22 \hat{\mathbf{y}}) = 220(-\hat{\mathbf{x}} + \hat{\mathbf{y}})$ .

$$|\nabla h| = 220\sqrt{2} \approx \boxed{311 \text{ ft/mile;}} \quad \text{direction: } \boxed{\text{northwest.}}$$

**Problem 1.13**

$$\mathbf{r} = (x - x')\hat{\mathbf{x}} + (y - y')\hat{\mathbf{y}} + (z - z')\hat{\mathbf{z}}; \quad r = \sqrt{(x - x')^2 + (y - y')^2 + (z - z')^2}.$$

$$(a) \nabla(r^2) = \frac{\partial}{\partial x}[(x - x')^2 + (y - y')^2 + (z - z')^2]\hat{\mathbf{x}} + \frac{\partial}{\partial y}[(x - x')^2 + (y - y')^2 + (z - z')^2]\hat{\mathbf{y}} + \frac{\partial}{\partial z}[(x - x')^2 + (y - y')^2 + (z - z')^2]\hat{\mathbf{z}} = 2(x - x')\hat{\mathbf{x}} + 2(y - y')\hat{\mathbf{y}} + 2(z - z')\hat{\mathbf{z}} = 2\mathbf{r}.$$

$$(b) \nabla\left(\frac{1}{r}\right) = \frac{\partial}{\partial x}[(x - x')^2 + (y - y')^2 + (z - z')^2]^{-\frac{1}{2}}\hat{\mathbf{x}} + \frac{\partial}{\partial y}[(x - x')^2 + (y - y')^2 + (z - z')^2]^{-\frac{1}{2}}\hat{\mathbf{y}} + \frac{\partial}{\partial z}[(x - x')^2 + (y - y')^2 + (z - z')^2]^{-\frac{1}{2}}\hat{\mathbf{z}} \\ = -\frac{1}{2}()^{-\frac{3}{2}}2(x - x')\hat{\mathbf{x}} - \frac{1}{2}()^{-\frac{3}{2}}2(y - y')\hat{\mathbf{y}} - \frac{1}{2}()^{-\frac{3}{2}}2(z - z')\hat{\mathbf{z}} \\ = -()^{-\frac{3}{2}}[(x - x')\hat{\mathbf{x}} + (y - y')\hat{\mathbf{y}} + (z - z')\hat{\mathbf{z}}] = -(1/r^3)\mathbf{r} = -(1/r^2)\hat{\mathbf{r}}.$$

$$(c) \frac{\partial}{\partial x}(r^n) = n r^{n-1} \frac{\partial r}{\partial x} = n r^{n-1} \left(\frac{1}{2} \frac{1}{r} 2 r_x\right) = n r^{n-1} \hat{\mathbf{r}}_x, \text{ so } \boxed{\nabla(r^n) = n r^{n-1} \hat{\mathbf{r}}}$$

**Problem 1.14**

$$\bar{y} = +y \cos \phi + z \sin \phi; \text{ multiply by } \sin \phi: \bar{y} \sin \phi = +y \sin \phi \cos \phi + z \sin^2 \phi.$$

$$\bar{z} = -y \sin \phi + z \cos \phi; \text{ multiply by } \cos \phi: \bar{z} \cos \phi = -y \sin \phi \cos \phi + z \cos^2 \phi.$$

$$\text{Add: } \bar{y} \sin \phi + \bar{z} \cos \phi = z(\sin^2 \phi + \cos^2 \phi) = z. \text{ Likewise, } \bar{y} \cos \phi - \bar{z} \sin \phi = y.$$

$$\text{So } \frac{\partial \bar{y}}{\partial y} = \cos \phi; \frac{\partial \bar{y}}{\partial \bar{z}} = -\sin \phi; \frac{\partial \bar{z}}{\partial y} = \sin \phi; \frac{\partial \bar{z}}{\partial \bar{z}} = \cos \phi. \text{ Therefore}$$


$$\left. \begin{aligned} \overline{(\nabla f)}_y &= \frac{\partial f}{\partial \bar{y}} = \frac{\partial f}{\partial y} \frac{\partial y}{\partial \bar{y}} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial \bar{y}} = +\cos \phi (\nabla f)_y + \sin \phi (\nabla f)_z \\ \overline{(\nabla f)}_z &= \frac{\partial f}{\partial \bar{z}} = \frac{\partial f}{\partial y} \frac{\partial y}{\partial \bar{z}} + \frac{\partial f}{\partial z} \frac{\partial z}{\partial \bar{z}} = -\sin \phi (\nabla f)_y + \cos \phi (\nabla f)_z \end{aligned} \right\} \text{ So } \nabla f \text{ transforms as a vector.} \quad \text{qed}$$

**Problem 1.15**

$$(a) \nabla \cdot \mathbf{v}_a = \frac{\partial}{\partial x}(x^2) + \frac{\partial}{\partial y}(3xz^2) + \frac{\partial}{\partial z}(-2xz) = 2x + 0 - 2x = 0.$$

$$(b) \nabla \cdot \mathbf{v}_b = \frac{\partial}{\partial x}(xy) + \frac{\partial}{\partial y}(2yz) + \frac{\partial}{\partial z}(3xz) = y + 2z + 3x.$$

$$(c) \nabla \cdot \mathbf{v}_c = \frac{\partial}{\partial x}(y^2) + \frac{\partial}{\partial y}(2xy + z^2) + \frac{\partial}{\partial z}(2yz) = 0 + (2x) + (2y) = 2(x + y)$$

**Problem 1.16**

$$\nabla \cdot \mathbf{v} = \frac{\partial}{\partial x}\left(\frac{x}{r^3}\right) + \frac{\partial}{\partial y}\left(\frac{y}{r^3}\right) + \frac{\partial}{\partial z}\left(\frac{z}{r^3}\right) = \frac{\partial}{\partial x} \left[ x(x^2 + y^2 + z^2)^{-\frac{3}{2}} \right] \\ + \frac{\partial}{\partial y} \left[ y(x^2 + y^2 + z^2)^{-\frac{3}{2}} \right] + \frac{\partial}{\partial z} \left[ z(x^2 + y^2 + z^2)^{-\frac{3}{2}} \right] \\ = ()^{-\frac{3}{2}} + x(-\frac{3}{2})()^{-\frac{5}{2}}2x + ()^{-\frac{3}{2}} + y(-\frac{3}{2})()^{-\frac{5}{2}}2y + ()^{-\frac{3}{2}} \\ + z(-\frac{3}{2})()^{-\frac{5}{2}}2z = 3r^{-3} - 3r^{-5}(x^2 + y^2 + z^2) = 3r^{-3} - 3r^{-3} = 0.$$

This conclusion is surprising, because, from the diagram, this vector field is obviously diverging away from the origin. How, then, can  $\nabla \cdot \mathbf{v} = 0$ ? The answer is that  $\nabla \cdot \mathbf{v} = 0$  everywhere *except* at the origin, but at the origin our calculation is no good, since  $r = 0$ , and the expression for  $\mathbf{v}$  blows up. In fact,  $\nabla \cdot \mathbf{v}$  is *infinite* at that one point, and zero elsewhere, as we shall see in Sect. 1.5.

**Problem 1.17**

$$\bar{v}_y = \cos \phi v_y + \sin \phi v_z; \quad \bar{v}_z = -\sin \phi v_y + \cos \phi v_z.$$

$$\frac{\partial \bar{v}_y}{\partial \bar{y}} = \frac{\partial v_y}{\partial y} \cos \phi + \frac{\partial v_z}{\partial y} \sin \phi = \left( \frac{\partial v_y}{\partial y} \frac{\partial y}{\partial \bar{y}} + \frac{\partial v_y}{\partial z} \frac{\partial z}{\partial \bar{y}} \right) \cos \phi + \left( \frac{\partial v_z}{\partial y} \frac{\partial y}{\partial \bar{y}} + \frac{\partial v_z}{\partial z} \frac{\partial z}{\partial \bar{y}} \right) \sin \phi. \text{ Use result in Prob. 1.14:}$$

$$= \left( \frac{\partial v_y}{\partial y} \cos \phi + \frac{\partial v_y}{\partial z} \sin \phi \right) \cos \phi + \left( \frac{\partial v_z}{\partial y} \cos \phi + \frac{\partial v_z}{\partial z} \sin \phi \right) \sin \phi.$$

$$\frac{\partial \bar{v}_z}{\partial \bar{z}} = -\frac{\partial v_y}{\partial y} \sin \phi + \frac{\partial v_z}{\partial y} \cos \phi = -\left( \frac{\partial v_y}{\partial y} \frac{\partial y}{\partial \bar{z}} + \frac{\partial v_y}{\partial z} \frac{\partial z}{\partial \bar{z}} \right) \sin \phi + \left( \frac{\partial v_z}{\partial y} \frac{\partial y}{\partial \bar{z}} + \frac{\partial v_z}{\partial z} \frac{\partial z}{\partial \bar{z}} \right) \cos \phi$$

$$= -\left( -\frac{\partial v_y}{\partial y} \sin \phi + \frac{\partial v_y}{\partial z} \cos \phi \right) \sin \phi + \left( -\frac{\partial v_z}{\partial y} \sin \phi + \frac{\partial v_z}{\partial z} \cos \phi \right) \cos \phi. \text{ So}$$


$$\begin{aligned}\frac{\partial \bar{v}_y}{\partial \bar{y}} + \frac{\partial \bar{v}_z}{\partial \bar{z}} &= \frac{\partial v_y}{\partial y} \cos^2 \phi + \frac{\partial v_y}{\partial z} \sin \phi \cos \phi + \frac{\partial v_z}{\partial y} \sin \phi \cos \phi + \frac{\partial v_z}{\partial z} \sin^2 \phi + \frac{\partial v_y}{\partial y} \sin^2 \phi - \frac{\partial v_y}{\partial z} \sin \phi \cos \phi \\ &\quad - \frac{\partial v_z}{\partial y} \sin \phi \cos \phi + \frac{\partial v_z}{\partial z} \cos^2 \phi \\ &= \frac{\partial v_y}{\partial y} (\cos^2 \phi + \sin^2 \phi) + \frac{\partial v_z}{\partial z} (\sin^2 \phi + \cos^2 \phi) = \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z}. \quad \checkmark\end{aligned}$$

**Problem 1.18**

$$(a) \nabla \times \mathbf{v}_a = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x^2 & 3xz^2 & -2xz \end{vmatrix} = \hat{\mathbf{x}}(0 - 6xz) + \hat{\mathbf{y}}(0 + 2z) + \hat{\mathbf{z}}(3z^2 - 0) = \boxed{-6xz \hat{\mathbf{x}} + 2z \hat{\mathbf{y}} + 3z^2 \hat{\mathbf{z}}}.$$

$$(b) \nabla \times \mathbf{v}_b = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ xy & 2yz & 3xz \end{vmatrix} = \hat{\mathbf{x}}(0 - 2y) + \hat{\mathbf{y}}(0 - 3z) + \hat{\mathbf{z}}(0 - x) = \boxed{-2y \hat{\mathbf{x}} - 3z \hat{\mathbf{y}} - x \hat{\mathbf{z}}}.$$

$$(c) \nabla \times \mathbf{v}_c = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y^2 & (2xy + z^2) & 2yz \end{vmatrix} = \hat{\mathbf{x}}(2z - 2z) + \hat{\mathbf{y}}(0 - 0) + \hat{\mathbf{z}}(2y - 2y) = \boxed{\mathbf{0}}.$$

**Problem 1.19**

As we go from point  $A$  to point  $B$  (9 o'clock to 10 o'clock),  $x$  increases,  $y$  increases,  $v_x$  increases, and  $v_y$  decreases, so  $\partial v_x / \partial y > 0$ , while  $\partial v_y / \partial y < 0$ . On the circle,  $v_z = 0$ , and there is no dependence on  $z$ , so Eq. 1.41 says

$$\nabla \times \mathbf{v} = \hat{\mathbf{z}} \left( \frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \right)$$

points in the negative  $z$  direction (into the page), as the right hand rule would suggest. (Pick any other nearby points on the circle and you will come to the same conclusion.) [I'm sorry, but I cannot remember who suggested this cute illustration.]

**Problem 1.20**

$$\begin{aligned}\mathbf{v} &= y \hat{\mathbf{x}} + x \hat{\mathbf{y}}; \text{ or } \mathbf{v} = yz \hat{\mathbf{x}} + xz \hat{\mathbf{y}} + xy \hat{\mathbf{z}}; \text{ or } \mathbf{v} = (3x^2z - z^3) \hat{\mathbf{x}} + 3\hat{\mathbf{y}} + (x^3 - 3xz^2) \hat{\mathbf{z}}; \\ \text{or } \mathbf{v} &= (\sin x)(\cosh y) \hat{\mathbf{x}} - (\cos x)(\sinh y) \hat{\mathbf{y}}; \text{ etc.}\end{aligned}$$

**Problem 1.21**

$$\begin{aligned}(i) \nabla(fg) &= \frac{\partial(fg)}{\partial x} \hat{\mathbf{x}} + \frac{\partial(fg)}{\partial y} \hat{\mathbf{y}} + \frac{\partial(fg)}{\partial z} \hat{\mathbf{z}} = \left( f \frac{\partial g}{\partial x} + g \frac{\partial f}{\partial x} \right) \hat{\mathbf{x}} + \left( f \frac{\partial g}{\partial y} + g \frac{\partial f}{\partial y} \right) \hat{\mathbf{y}} + \left( f \frac{\partial g}{\partial z} + g \frac{\partial f}{\partial z} \right) \hat{\mathbf{z}} \\ &= f \left( \frac{\partial g}{\partial x} \hat{\mathbf{x}} + \frac{\partial g}{\partial y} \hat{\mathbf{y}} + \frac{\partial g}{\partial z} \hat{\mathbf{z}} \right) + g \left( \frac{\partial f}{\partial x} \hat{\mathbf{x}} + \frac{\partial f}{\partial y} \hat{\mathbf{y}} + \frac{\partial f}{\partial z} \hat{\mathbf{z}} \right) = f(\nabla g) + g(\nabla f). \quad \text{qed}\end{aligned}$$

$$\begin{aligned}(iv) \nabla \cdot (\mathbf{A} \times \mathbf{B}) &= \frac{\partial}{\partial x} (A_y B_z - A_z B_y) + \frac{\partial}{\partial y} (A_z B_x - A_x B_z) + \frac{\partial}{\partial z} (A_x B_y - A_y B_x) \\ &= A_y \frac{\partial B_z}{\partial x} + B_z \frac{\partial A_y}{\partial x} - A_z \frac{\partial B_y}{\partial x} - B_y \frac{\partial A_z}{\partial x} + A_z \frac{\partial B_x}{\partial y} + B_x \frac{\partial A_z}{\partial y} - A_x \frac{\partial B_z}{\partial y} - B_z \frac{\partial A_x}{\partial y} \\ &\quad + A_x \frac{\partial B_y}{\partial z} + B_y \frac{\partial A_x}{\partial z} - A_y \frac{\partial B_x}{\partial z} - B_x \frac{\partial A_y}{\partial z} \\ &= B_x \left( \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right) + B_y \left( \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right) + B_z \left( \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right) - A_x \left( \frac{\partial B_z}{\partial y} - \frac{\partial B_y}{\partial z} \right) \\ &\quad - A_y \left( \frac{\partial B_x}{\partial z} - \frac{\partial B_z}{\partial x} \right) - A_z \left( \frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} \right) = \mathbf{B} \cdot (\nabla \times \mathbf{A}) - \mathbf{A} \cdot (\nabla \times \mathbf{B}). \quad \text{qed}\end{aligned}$$

$$(v) \nabla \times (f\mathbf{A}) = \left( \frac{\partial(fA_z)}{\partial y} - \frac{\partial(fA_y)}{\partial z} \right) \hat{\mathbf{x}} + \left( \frac{\partial(fA_x)}{\partial z} - \frac{\partial(fA_z)}{\partial x} \right) \hat{\mathbf{y}} + \left( \frac{\partial(fA_y)}{\partial x} - \frac{\partial(fA_x)}{\partial y} \right) \hat{\mathbf{z}}$$

$$\begin{aligned}
&= \left( f \frac{\partial A_z}{\partial y} + A_z \frac{\partial f}{\partial y} - f \frac{\partial A_y}{\partial z} - A_y \frac{\partial f}{\partial z} \right) \hat{\mathbf{x}} + \left( f \frac{\partial A_x}{\partial z} + A_x \frac{\partial f}{\partial z} - f \frac{\partial A_z}{\partial x} - A_z \frac{\partial f}{\partial x} \right) \hat{\mathbf{y}} \\
&\quad + \left( f \frac{\partial A_y}{\partial x} + A_y \frac{\partial f}{\partial x} - f \frac{\partial A_x}{\partial y} - A_x \frac{\partial f}{\partial y} \right) \hat{\mathbf{z}} \\
&= f \left[ \left( \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right) \hat{\mathbf{x}} + \left( \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right) \hat{\mathbf{y}} + \left( \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right) \hat{\mathbf{z}} \right] \\
&\quad - \left[ \left( A_y \frac{\partial f}{\partial z} - A_z \frac{\partial f}{\partial y} \right) \hat{\mathbf{x}} + \left( A_z \frac{\partial f}{\partial x} - A_x \frac{\partial f}{\partial z} \right) \hat{\mathbf{y}} + \left( A_x \frac{\partial f}{\partial y} - A_y \frac{\partial f}{\partial x} \right) \hat{\mathbf{z}} \right] \\
&= f (\nabla \times \mathbf{A}) - \mathbf{A} \times (\nabla f). \quad \text{qed}
\end{aligned}$$

**Problem 1.22**

$$\begin{aligned}
\text{(a)} \quad (\mathbf{A} \cdot \nabla) \mathbf{B} &= \left( A_x \frac{\partial B_x}{\partial x} + A_y \frac{\partial B_x}{\partial y} + A_z \frac{\partial B_x}{\partial z} \right) \hat{\mathbf{x}} + \left( A_x \frac{\partial B_y}{\partial x} + A_y \frac{\partial B_y}{\partial y} + A_z \frac{\partial B_y}{\partial z} \right) \hat{\mathbf{y}} \\
&\quad + \left( A_x \frac{\partial B_z}{\partial x} + A_y \frac{\partial B_z}{\partial y} + A_z \frac{\partial B_z}{\partial z} \right) \hat{\mathbf{z}}.
\end{aligned}$$

$$\text{(b)} \quad \hat{\mathbf{r}} = \frac{\mathbf{r}}{r} = \frac{x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}}}{\sqrt{x^2 + y^2 + z^2}}. \quad \text{Let's just do the } x \text{ component.}$$

$$\begin{aligned}
[(\hat{\mathbf{r}} \cdot \nabla) \hat{\mathbf{r}}]_x &= \frac{1}{\sqrt{x^2 + y^2 + z^2}} \left( x \frac{\partial}{\partial x} + y \frac{\partial}{\partial y} + z \frac{\partial}{\partial z} \right) \frac{x}{\sqrt{x^2 + y^2 + z^2}} \\
&= \frac{1}{r} \left\{ x \left[ \frac{1}{\sqrt{x^2 + y^2 + z^2}} + x \left( -\frac{1}{2} \right) \frac{1}{(\sqrt{x^2 + y^2 + z^2})^3} 2x \right] + yx \left[ -\frac{1}{2} \frac{1}{(\sqrt{x^2 + y^2 + z^2})^3} 2y \right] + zx \left[ -\frac{1}{2} \frac{1}{(\sqrt{x^2 + y^2 + z^2})^3} 2z \right] \right\} \\
&= \frac{1}{r} \left\{ \frac{x}{r} - \frac{1}{r^3} (x^3 + xy^2 + xz^2) \right\} = \frac{1}{r} \left\{ \frac{x}{r} - \frac{x}{r^3} (x^2 + y^2 + z^2) \right\} = \frac{1}{r} \left( \frac{x}{r} - \frac{x}{r} \right) = 0.
\end{aligned}$$

Same goes for the other components. Hence:  $(\hat{\mathbf{r}} \cdot \nabla) \hat{\mathbf{r}} = \mathbf{0}$ .

$$\begin{aligned}
\text{(c)} \quad (\mathbf{v}_a \cdot \nabla) \mathbf{v}_b &= \left( x^2 \frac{\partial}{\partial x} + 3xz^2 \frac{\partial}{\partial y} - 2xz \frac{\partial}{\partial z} \right) (xy\hat{\mathbf{x}} + 2yz\hat{\mathbf{y}} + 3xz\hat{\mathbf{z}}) \\
&= x^2 (y\hat{\mathbf{x}} + 0\hat{\mathbf{y}} + 3z\hat{\mathbf{z}}) + 3xz^2 (x\hat{\mathbf{x}} + 2z\hat{\mathbf{y}} + 0\hat{\mathbf{z}}) - 2xz (0\hat{\mathbf{x}} + 2y\hat{\mathbf{y}} + 3x\hat{\mathbf{z}}) \\
&= (x^2y + 3x^2z^2) \hat{\mathbf{x}} + (6xz^3 - 4xyz) \hat{\mathbf{y}} + (3x^2z - 6x^2z) \hat{\mathbf{z}} \\
&= \boxed{x^2 (y + 3z^2) \hat{\mathbf{x}} + 2xz (3z^2 - 2y) \hat{\mathbf{y}} - 3x^2z \hat{\mathbf{z}}}
\end{aligned}$$

**Problem 1.23**

$$\text{(ii)} \quad [\nabla(\mathbf{A} \cdot \mathbf{B})]_x = \frac{\partial}{\partial x} (A_x B_x + A_y B_y + A_z B_z) = \frac{\partial A_x}{\partial x} B_x + A_x \frac{\partial B_x}{\partial x} + \frac{\partial A_y}{\partial x} B_y + A_y \frac{\partial B_y}{\partial x} + \frac{\partial A_z}{\partial x} B_z + A_z \frac{\partial B_z}{\partial x}$$

$$[\mathbf{A} \times (\nabla \times \mathbf{B})]_x = A_y (\nabla \times \mathbf{B})_z - A_z (\nabla \times \mathbf{B})_y = A_y \left( \frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} \right) - A_z \left( \frac{\partial B_x}{\partial z} - \frac{\partial B_z}{\partial x} \right)$$

$$[\mathbf{B} \times (\nabla \times \mathbf{A})]_x = B_y \left( \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} \right) - B_z \left( \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} \right)$$

$$[(\mathbf{A} \cdot \nabla) \mathbf{B}]_x = \left( A_x \frac{\partial}{\partial x} + A_y \frac{\partial}{\partial y} + A_z \frac{\partial}{\partial z} \right) B_x = A_x \frac{\partial B_x}{\partial x} + A_y \frac{\partial B_x}{\partial y} + A_z \frac{\partial B_x}{\partial z}$$

$$[(\mathbf{B} \cdot \nabla) \mathbf{A}]_x = B_x \frac{\partial A_x}{\partial x} + B_y \frac{\partial A_x}{\partial y} + B_z \frac{\partial A_x}{\partial z}$$

$$\begin{aligned}
\text{So } [\mathbf{A} \times (\nabla \times \mathbf{B}) + \mathbf{B} \times (\nabla \times \mathbf{A}) + (\mathbf{A} \cdot \nabla) \mathbf{B} + (\mathbf{B} \cdot \nabla) \mathbf{A}]_x &= A_y \frac{\partial B_y}{\partial x} - A_y \frac{\partial B_x}{\partial y} - A_z \frac{\partial B_x}{\partial z} + A_z \frac{\partial B_z}{\partial x} + B_y \frac{\partial A_y}{\partial x} - B_y \frac{\partial A_x}{\partial y} - B_z \frac{\partial A_x}{\partial z} + B_z \frac{\partial A_z}{\partial x} \\
&\quad + A_x \frac{\partial B_x}{\partial x} + A_y \frac{\partial B_y}{\partial y} + A_z \frac{\partial B_z}{\partial z} + B_x \frac{\partial A_x}{\partial x} + B_y \frac{\partial A_y}{\partial y} + B_z \frac{\partial A_z}{\partial z} \\
&= B_x \frac{\partial A_x}{\partial x} + A_x \frac{\partial B_x}{\partial x} + B_y \left( \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} + \frac{\partial A_x}{\partial y} \right) + A_y \left( \frac{\partial B_y}{\partial x} - \frac{\partial B_x}{\partial y} + \frac{\partial B_x}{\partial y} \right) \\
&\quad + B_z \left( -\frac{\partial A_x}{\partial z} + \frac{\partial A_z}{\partial x} + \frac{\partial A_x}{\partial z} \right) + A_z \left( -\frac{\partial B_x}{\partial z} + \frac{\partial B_z}{\partial x} + \frac{\partial B_x}{\partial z} \right) \\
&= [\nabla(\mathbf{A} \cdot \mathbf{B})]_x \quad (\text{same for } y \text{ and } z)
\end{aligned}$$

$$\begin{aligned}
\text{(vi)} \quad [\nabla \times (\mathbf{A} \times \mathbf{B})]_x &= \frac{\partial}{\partial y} (\mathbf{A} \times \mathbf{B})_z - \frac{\partial}{\partial z} (\mathbf{A} \times \mathbf{B})_y = \frac{\partial}{\partial y} (A_x B_y - A_y B_x) - \frac{\partial}{\partial z} (A_z B_x - A_x B_z) \\
&= \frac{\partial A_x}{\partial y} B_y + A_x \frac{\partial B_y}{\partial y} - \frac{\partial A_y}{\partial y} B_x - A_y \frac{\partial B_x}{\partial y} - \frac{\partial A_z}{\partial z} B_x - A_z \frac{\partial B_x}{\partial z} + \frac{\partial A_x}{\partial z} B_z + A_x \frac{\partial B_z}{\partial z}
\end{aligned}$$

$$\begin{aligned}
[(\mathbf{B} \cdot \nabla) \mathbf{A} - (\mathbf{A} \cdot \nabla) \mathbf{B} + \mathbf{A}(\nabla \cdot \mathbf{B}) - \mathbf{B}(\nabla \cdot \mathbf{A})]_x &= B_x \frac{\partial A_x}{\partial x} + B_y \frac{\partial A_x}{\partial y} + B_z \frac{\partial A_x}{\partial z} - A_x \frac{\partial B_x}{\partial x} - A_y \frac{\partial B_x}{\partial y} - A_z \frac{\partial B_x}{\partial z} + A_x \left( \frac{\partial B_x}{\partial x} + \frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z} \right) - B_x \left( \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} \right)
\end{aligned}$$

$$\begin{aligned}
&= B_y \frac{\partial A_x}{\partial y} + A_x \left( -\frac{\partial B_y}{\partial x} + \frac{\partial B_x}{\partial y} + \frac{\partial B_z}{\partial z} \right) + B_x \left( \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} - \frac{\partial A_z}{\partial z} \right) \\
&\quad + A_y \left( -\frac{\partial B_x}{\partial y} \right) + A_z \left( -\frac{\partial B_z}{\partial z} \right) + B_z \left( \frac{\partial A_z}{\partial z} \right) \\
&= [\nabla \times (\mathbf{A} \times \mathbf{B})]_x \text{ (same for } y \text{ and } z)
\end{aligned}$$

**Problem 1.24**

$$\begin{aligned}
\nabla(f/g) &= \frac{\partial}{\partial x}(f/g) \hat{\mathbf{x}} + \frac{\partial}{\partial y}(f/g) \hat{\mathbf{y}} + \frac{\partial}{\partial z}(f/g) \hat{\mathbf{z}} \\
&= \frac{g \frac{\partial f}{\partial x} - f \frac{\partial g}{\partial x}}{g^2} \hat{\mathbf{x}} + \frac{g \frac{\partial f}{\partial y} - f \frac{\partial g}{\partial y}}{g^2} \hat{\mathbf{y}} + \frac{g \frac{\partial f}{\partial z} - f \frac{\partial g}{\partial z}}{g^2} \hat{\mathbf{z}} \\
&= \frac{1}{g^2} \left[ g \left( \frac{\partial f}{\partial x} \hat{\mathbf{x}} + \frac{\partial f}{\partial y} \hat{\mathbf{y}} + \frac{\partial f}{\partial z} \hat{\mathbf{z}} \right) - f \left( \frac{\partial g}{\partial x} \hat{\mathbf{x}} + \frac{\partial g}{\partial y} \hat{\mathbf{y}} + \frac{\partial g}{\partial z} \hat{\mathbf{z}} \right) \right] = \frac{g \nabla f - f \nabla g}{g^2}. \quad \text{qed}
\end{aligned}$$

$$\begin{aligned}
\nabla \cdot (\mathbf{A}/g) &= \frac{\partial}{\partial x}(A_x/g) + \frac{\partial}{\partial y}(A_y/g) + \frac{\partial}{\partial z}(A_z/g) \\
&= \frac{g \frac{\partial A_x}{\partial x} - A_x \frac{\partial g}{\partial x}}{g^2} + \frac{g \frac{\partial A_y}{\partial y} - A_y \frac{\partial g}{\partial y}}{g^2} + \frac{g \frac{\partial A_z}{\partial z} - A_z \frac{\partial g}{\partial z}}{g^2} \\
&= \frac{1}{g^2} \left[ g \left( \frac{\partial A_x}{\partial x} + \frac{\partial A_y}{\partial y} + \frac{\partial A_z}{\partial z} \right) - \left( A_x \frac{\partial g}{\partial x} + A_y \frac{\partial g}{\partial y} + A_z \frac{\partial g}{\partial z} \right) \right] = \frac{g \nabla \cdot \mathbf{A} - \mathbf{A} \cdot \nabla g}{g^2}. \quad \text{qed}
\end{aligned}$$

$$\begin{aligned}
[\nabla \times (\mathbf{A}/g)]_x &= \frac{\partial}{\partial y}(A_z/g) - \frac{\partial}{\partial z}(A_y/g) \\
&= \frac{g \frac{\partial A_z}{\partial y} - A_z \frac{\partial g}{\partial y}}{g^2} - \frac{g \frac{\partial A_y}{\partial z} - A_y \frac{\partial g}{\partial z}}{g^2} \\
&= \frac{1}{g^2} \left[ g \left( \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} \right) - \left( A_z \frac{\partial g}{\partial y} - A_y \frac{\partial g}{\partial z} \right) \right] \\
&= \frac{g(\nabla \times \mathbf{A})_x + (\mathbf{A} \times \nabla g)_x}{g^2} \text{ (same for } y \text{ and } z). \quad \text{qed}
\end{aligned}$$

**Problem 1.25**

$$(a) \mathbf{A} \times \mathbf{B} = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ x & 2y & 3z \\ 3y & -2x & 0 \end{vmatrix} = \hat{\mathbf{x}}(6xz) + \hat{\mathbf{y}}(9zy) + \hat{\mathbf{z}}(-2x^2 - 6y^2)$$

$$\nabla \cdot (\mathbf{A} \times \mathbf{B}) = \frac{\partial}{\partial x}(6xz) + \frac{\partial}{\partial y}(9zy) + \frac{\partial}{\partial z}(-2x^2 - 6y^2) = 6z + 9z + 0 = 15z$$

$$\nabla \times \mathbf{A} = \hat{\mathbf{x}} \left( \frac{\partial}{\partial y}(3z) - \frac{\partial}{\partial z}(2y) \right) + \hat{\mathbf{y}} \left( \frac{\partial}{\partial z}(x) - \frac{\partial}{\partial x}(3z) \right) + \hat{\mathbf{z}} \left( \frac{\partial}{\partial x}(2y) - \frac{\partial}{\partial y}(x) \right) = 0; \quad \mathbf{B} \cdot (\nabla \times \mathbf{A}) = 0$$

$$\nabla \times \mathbf{B} = \hat{\mathbf{x}} \left( \frac{\partial}{\partial y}(0) - \frac{\partial}{\partial z}(-2x) \right) + \hat{\mathbf{y}} \left( \frac{\partial}{\partial z}(3y) - \frac{\partial}{\partial x}(0) \right) + \hat{\mathbf{z}} \left( \frac{\partial}{\partial x}(-2x) - \frac{\partial}{\partial y}(3y) \right) = -5\hat{\mathbf{z}}; \quad \mathbf{A} \cdot (\nabla \times \mathbf{B}) = -15z$$

$$\nabla \cdot (\mathbf{A} \times \mathbf{B}) \stackrel{?}{=} \mathbf{B} \cdot (\nabla \times \mathbf{A}) - \mathbf{A} \cdot (\nabla \times \mathbf{B}) = 0 - (-15z) = 15z. \quad \checkmark$$

$$(b) \mathbf{A} \cdot \mathbf{B} = 3xy - 4xy = -xy; \quad \nabla(\mathbf{A} \cdot \mathbf{B}) = \nabla(-xy) = \hat{\mathbf{x}} \frac{\partial}{\partial x}(-xy) + \hat{\mathbf{y}} \frac{\partial}{\partial y}(-xy) = -y\hat{\mathbf{x}} - x\hat{\mathbf{y}}$$

$$\mathbf{A} \times (\nabla \times \mathbf{B}) = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ x & 2y & 3z \\ 0 & 0 & -5 \end{vmatrix} = \hat{\mathbf{x}}(-10y) + \hat{\mathbf{y}}(5x); \quad \mathbf{B} \times (\nabla \times \mathbf{A}) = 0$$

$$(\mathbf{A} \cdot \nabla) \mathbf{B} = \left( x \frac{\partial}{\partial x} + 2y \frac{\partial}{\partial y} + 3z \frac{\partial}{\partial z} \right) (3y\hat{\mathbf{x}} - 2x\hat{\mathbf{y}}) = \hat{\mathbf{x}}(6y) + \hat{\mathbf{y}}(-2x)$$

$$(\mathbf{B} \cdot \nabla) \mathbf{A} = \left( 3y \frac{\partial}{\partial x} - 2x \frac{\partial}{\partial y} \right) (x\hat{\mathbf{x}} + 2y\hat{\mathbf{y}} + 3z\hat{\mathbf{z}}) = \hat{\mathbf{x}}(3y) + \hat{\mathbf{y}}(-4x)$$

$$\begin{aligned} \mathbf{A} \times (\nabla \times \mathbf{B}) + \mathbf{B} \times (\nabla \times \mathbf{A}) + (\mathbf{A} \cdot \nabla) \mathbf{B} + (\mathbf{B} \cdot \nabla) \mathbf{A} \\ = -10y\hat{\mathbf{x}} + 5x\hat{\mathbf{y}} + 6y\hat{\mathbf{x}} - 2x\hat{\mathbf{y}} + 3y\hat{\mathbf{x}} - 4x\hat{\mathbf{y}} = -y\hat{\mathbf{x}} - x\hat{\mathbf{y}} = \nabla \cdot (\mathbf{A} \cdot \mathbf{B}). \quad \checkmark \end{aligned}$$

$$(c) \nabla \times (\mathbf{A} \times \mathbf{B}) = \hat{\mathbf{x}} \left( \frac{\partial}{\partial y}(-2x^2 - 6y^2) - \frac{\partial}{\partial z}(9zy) \right) + \hat{\mathbf{y}} \left( \frac{\partial}{\partial z}(6xz) - \frac{\partial}{\partial x}(-2x^2 - 6y^2) \right) + \hat{\mathbf{z}} \left( \frac{\partial}{\partial x}(9zy) - \frac{\partial}{\partial y}(6xz) \right) \\ = \hat{\mathbf{x}}(-12y - 9y) + \hat{\mathbf{y}}(6x + 4x) + \hat{\mathbf{z}}(0) = -21y\hat{\mathbf{x}} + 10x\hat{\mathbf{y}}$$

$$\nabla \cdot \mathbf{A} = \frac{\partial}{\partial x}(x) + \frac{\partial}{\partial y}(2y) + \frac{\partial}{\partial z}(3z) = 1 + 2 + 3 = 6; \quad \nabla \cdot \mathbf{B} = \frac{\partial}{\partial x}(3y) + \frac{\partial}{\partial y}(-2x) = 0$$



$$(\mathbf{B} \cdot \nabla) \mathbf{A} - (\mathbf{A} \cdot \nabla) \mathbf{B} + \mathbf{A}(\nabla \cdot \mathbf{B}) - \mathbf{B}(\nabla \cdot \mathbf{A}) = 3y \hat{\mathbf{x}} - 4x \hat{\mathbf{y}} - 6y \hat{\mathbf{x}} + 2x \hat{\mathbf{y}} - 18y \hat{\mathbf{x}} + 12x \hat{\mathbf{y}} = -21y \hat{\mathbf{x}} + 10x \hat{\mathbf{y}} \\ = \nabla \times (\mathbf{A} \times \mathbf{B}). \checkmark$$

**Problem 1.26**

- (a)  $\frac{\partial^2 T_a}{\partial x^2} = 2; \frac{\partial^2 T_a}{\partial y^2} = \frac{\partial^2 T_a}{\partial z^2} = 0 \Rightarrow \boxed{\nabla^2 T_a = 2.}$
- (b)  $\frac{\partial^2 T_b}{\partial x^2} = \frac{\partial^2 T_b}{\partial y^2} = \frac{\partial^2 T_b}{\partial z^2} = -T_b \Rightarrow \boxed{\nabla^2 T_b = -3T_b = -3 \sin x \sin y \sin z.}$
- (c)  $\frac{\partial^2 T_c}{\partial x^2} = 25T_c; \frac{\partial^2 T_c}{\partial y^2} = -16T_c; \frac{\partial^2 T_c}{\partial z^2} = -9T_c \Rightarrow \boxed{\nabla^2 T_c = 0.}$
- (d)  $\left. \begin{aligned} \frac{\partial^2 v_x}{\partial x^2} &= 2; \frac{\partial^2 v_x}{\partial y^2} = \frac{\partial^2 v_x}{\partial z^2} = 0 \Rightarrow \nabla^2 v_x = 2 \\ \frac{\partial^2 v_y}{\partial x^2} &= \frac{\partial^2 v_y}{\partial y^2} = 0; \frac{\partial^2 v_y}{\partial z^2} = 6x \Rightarrow \nabla^2 v_y = 6x \\ \frac{\partial^2 v_z}{\partial x^2} &= \frac{\partial^2 v_z}{\partial y^2} = \frac{\partial^2 v_z}{\partial z^2} = 0 \Rightarrow \nabla^2 v_z = 0 \end{aligned} \right\} \boxed{\nabla^2 \mathbf{v} = 2 \hat{\mathbf{x}} + 6x \hat{\mathbf{y}}.}$

**Problem 1.27**

$$\nabla \cdot (\nabla \times \mathbf{v}) = \frac{\partial}{\partial x} \left( \frac{\partial v_z}{\partial y} - \frac{\partial v_y}{\partial z} \right) + \frac{\partial}{\partial y} \left( \frac{\partial v_x}{\partial z} - \frac{\partial v_z}{\partial x} \right) + \frac{\partial}{\partial z} \left( \frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \right) \\ = \left( \frac{\partial^2 v_z}{\partial x \partial y} - \frac{\partial^2 v_z}{\partial y \partial x} \right) + \left( \frac{\partial^2 v_x}{\partial y \partial z} - \frac{\partial^2 v_x}{\partial z \partial y} \right) + \left( \frac{\partial^2 v_y}{\partial z \partial x} - \frac{\partial^2 v_y}{\partial x \partial z} \right) = 0, \text{ by equality of cross-derivatives.}$$

From Prob. 1.18:  $\nabla \times \mathbf{v}_a = -6xz \hat{\mathbf{x}} + 2z \hat{\mathbf{y}} + 3z^2 \hat{\mathbf{z}} \Rightarrow \nabla \cdot (\nabla \times \mathbf{v}_a) = \frac{\partial}{\partial x}(-6xz) + \frac{\partial}{\partial y}(2z) + \frac{\partial}{\partial z}(3z^2) = -6z + 6z = 0.$

**Problem 1.28**

$$\nabla \times (\nabla t) = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ \frac{\partial t}{\partial x} & \frac{\partial t}{\partial y} & \frac{\partial t}{\partial z} \end{vmatrix} = \hat{\mathbf{x}} \left( \frac{\partial^2 t}{\partial y \partial z} - \frac{\partial^2 t}{\partial z \partial y} \right) + \hat{\mathbf{y}} \left( \frac{\partial^2 t}{\partial z \partial x} - \frac{\partial^2 t}{\partial x \partial z} \right) + \hat{\mathbf{z}} \left( \frac{\partial^2 t}{\partial x \partial y} - \frac{\partial^2 t}{\partial y \partial x} \right) \\ = 0, \text{ by equality of cross-derivatives.}$$

In Prob. 1.11(b),  $\nabla f = 2xy^3z^4 \hat{\mathbf{x}} + 3x^2y^2z^4 \hat{\mathbf{y}} + 4x^2y^3z^3 \hat{\mathbf{z}}$ , so

$$\nabla \times (\nabla f) = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 2xy^3z^4 & 3x^2y^2z^4 & 4x^2y^3z^3 \end{vmatrix} \\ = \hat{\mathbf{x}}(3 \cdot 4x^2y^2z^3 - 4 \cdot 3x^2y^2z^3) + \hat{\mathbf{y}}(4 \cdot 2xy^3z^3 - 2 \cdot 4xy^3z^3) + \hat{\mathbf{z}}(2 \cdot 3xy^2z^4 - 3 \cdot 2xy^2z^4) = 0. \checkmark$$

**Problem 1.29**

- (a)  $(0, 0, 0) \rightarrow (1, 0, 0)$ .  $x: 0 \rightarrow 1, y = z = 0; d\mathbf{l} = dx \hat{\mathbf{x}}; \mathbf{v} \cdot d\mathbf{l} = x^2 dx; \int \mathbf{v} \cdot d\mathbf{l} = \int_0^1 x^2 dx = (x^3/3)|_0^1 = 1/3.$   
 $(1, 0, 0) \rightarrow (1, 1, 0)$ .  $x = 1, y: 0 \rightarrow 1, z = 0; d\mathbf{l} = dy \hat{\mathbf{y}}; \mathbf{v} \cdot d\mathbf{l} = 2yz dy = 0; \int \mathbf{v} \cdot d\mathbf{l} = 0.$   
 $(1, 1, 0) \rightarrow (1, 1, 1)$ .  $x = y = 1, z: 0 \rightarrow 1; d\mathbf{l} = dz \hat{\mathbf{z}}; \mathbf{v} \cdot d\mathbf{l} = y^2 dz = dz; \int \mathbf{v} \cdot d\mathbf{l} = \int_0^1 dz = z|_0^1 = 1.$   
**Total:**  $\int \mathbf{v} \cdot d\mathbf{l} = (1/3) + 0 + 1 = \boxed{4/3.}$
- (b)  $(0, 0, 0) \rightarrow (0, 0, 1)$ .  $x = y = 0, z: 0 \rightarrow 1; d\mathbf{l} = dz \hat{\mathbf{z}}; \mathbf{v} \cdot d\mathbf{l} = y^2 dz = 0; \int \mathbf{v} \cdot d\mathbf{l} = 0.$   
 $(0, 0, 1) \rightarrow (0, 1, 1)$ .  $x = 0, y: 0 \rightarrow 1, z = 1; d\mathbf{l} = dy \hat{\mathbf{y}}; \mathbf{v} \cdot d\mathbf{l} = 2yz dy = 2y dy; \int \mathbf{v} \cdot d\mathbf{l} = \int_0^1 2y dy = y^2|_0^1 = 1.$   
 $(0, 1, 1) \rightarrow (1, 1, 1)$ .  $x: 0 \rightarrow 1, y = z = 1; d\mathbf{l} = dx \hat{\mathbf{x}}; \mathbf{v} \cdot d\mathbf{l} = x^2 dx; \int \mathbf{v} \cdot d\mathbf{l} = \int_0^1 x^2 dx = (x^3/3)|_0^1 = 1/3.$   
**Total:**  $\int \mathbf{v} \cdot d\mathbf{l} = 0 + 1 + (1/3) = \boxed{4/3.}$
- (c)  $x = y = z: 0 \rightarrow 1; dx = dy = dz; \mathbf{v} \cdot d\mathbf{l} = x^2 dx + 2yz dy + y^2 dz = x^2 dx + 2x^2 dx + x^2 dx = 4x^2 dx;$   
 $\int \mathbf{v} \cdot d\mathbf{l} = \int_0^1 4x^2 dx = (4x^3/3)|_0^1 = \boxed{4/3.}$
- (d)  $\oint \mathbf{v} \cdot d\mathbf{l} = (4/3) - (4/3) = \boxed{0.}$

**Problem 1.30**

$x, y : 0 \rightarrow 1, z = 0; d\mathbf{a} = dx dy \hat{\mathbf{z}}; \mathbf{v} \cdot d\mathbf{a} = y(z^2 - 3) dx dy = -3y dx dy; \int \mathbf{v} \cdot d\mathbf{a} = -3 \int_0^2 dx \int_0^2 y dy = -3(x|_0^2)(\frac{y^2}{2}|_0^2) = -3(2)(2) = \boxed{-12.}$  In Ex. 1.7 we got 20, for the same boundary line (the square in the  $xy$ -plane), so the answer is no: the surface integral does *not* depend only on the boundary line. The *total* flux for the cube is  $20 + 12 = \boxed{32.}$

**Problem 1.31**

$\int T d\tau = \int z^2 dx dy dz$ . You can do the integrals in any order—here it is simplest to save  $z$  for last:

$$\int z^2 \left[ \int \left( \int dx \right) dy \right] dz.$$

The sloping surface is  $x + y + z = 1$ , so the  $x$  integral is  $\int_0^{(1-y-z)} dx = 1 - y - z$ . For a given  $z$ ,  $y$  ranges from 0 to  $1 - z$ , so the  $y$  integral is  $\int_0^{(1-z)} (1 - y - z) dy = [(1 - z)y - (y^2/2)]_0^{(1-z)} = (1 - z)^2 - [(1 - z)^2/2] = (1 - z)^2/2 = (1/2) - z + (z^2/2)$ . Finally, the  $z$  integral is  $\int_0^1 z^2(\frac{1}{2} - z + \frac{z^2}{2}) dz = \int_0^1 (\frac{z^2}{2} - z^3 + \frac{z^4}{2}) dz = (\frac{z^3}{6} - \frac{z^4}{4} + \frac{z^5}{10})|_0^1 = \frac{1}{6} - \frac{1}{4} + \frac{1}{10} = \boxed{1/60.}$

**Problem 1.32**

$$T(\mathbf{b}) = 1 + 4 + 2 = 7; T(\mathbf{a}) = 0. \Rightarrow \boxed{T(\mathbf{b}) - T(\mathbf{a}) = 7.}$$

$$\nabla T = (2x + 4y)\hat{\mathbf{x}} + (4x + 2z^3)\hat{\mathbf{y}} + (6yz^2)\hat{\mathbf{z}}; \nabla T \cdot d\mathbf{l} = (2x + 4y)dx + (4x + 2z^3)dy + (6yz^2)dz$$

- (a) Segment 1:  $x : 0 \rightarrow 1, y = z = dy = dz = 0. \int \nabla T \cdot d\mathbf{l} = \int_0^1 (2x) dx = x^2|_0^1 = 1.$   
 Segment 2:  $y : 0 \rightarrow 1, x = 1, z = 0, dx = dz = 0. \int \nabla T \cdot d\mathbf{l} = \int_0^1 (4) dy = 4y|_0^1 = 4.$   
 Segment 3:  $z : 0 \rightarrow 1, x = y = 1, dx = dy = 0. \int \nabla T \cdot d\mathbf{l} = \int_0^1 (6z^2) dz = 2z^3|_0^1 = 2.$   $\left. \begin{array}{l} \int_{\mathbf{a}}^{\mathbf{b}} \nabla T \cdot d\mathbf{l} = 7. \checkmark \end{array} \right\}$
- (b) Segment 1:  $z : 0 \rightarrow 1, x = y = dx = dy = 0. \int \nabla T \cdot d\mathbf{l} = \int_0^1 (0) dz = 0.$   
 Segment 2:  $y : 0 \rightarrow 1, x = 0, z = 1, dx = dz = 0. \int \nabla T \cdot d\mathbf{l} = \int_0^1 (2) dy = 2y|_0^1 = 2.$   
 Segment 3:  $x : 0 \rightarrow 1, y = z = 1, dy = dz = 0. \int \nabla T \cdot d\mathbf{l} = \int_0^1 (2x + 4) dx = (x^2 + 4x)|_0^1 = 1 + 4 = 5.$   $\left. \begin{array}{l} \int_{\mathbf{a}}^{\mathbf{b}} \nabla T \cdot d\mathbf{l} = 7. \checkmark \end{array} \right\}$
- (c)  $x : 0 \rightarrow 1, y = x, z = x^2, dy = dx, dz = 2x dx.$

$$\nabla T \cdot d\mathbf{l} = (2x + 4x)dx + (4x + 2x^6)dx + (6xx^4)2x dx = (10x + 14x^6)dx.$$

$$\int_{\mathbf{a}}^{\mathbf{b}} \nabla T \cdot d\mathbf{l} = \int_0^1 (10x + 14x^6)dx = (5x^2 + 2x^7)|_0^1 = 5 + 2 = 7. \checkmark$$

**Problem 1.33**

$$\nabla \cdot \mathbf{v} = y + 2z + 3x$$

$$\begin{aligned} \int (\nabla \cdot \mathbf{v}) d\tau &= \int (y + 2z + 3x) dx dy dz = \iint \left\{ \int_0^2 (y + 2z + 3x) dx \right\} dy dz \\ &\quad \hookrightarrow [(y + 2z)x + \frac{3}{2}x^2]_0^2 = 2(y + 2z) + 6 \\ &= \int \left\{ \int_0^2 (2y + 4z + 6) dy \right\} dz \\ &\quad \hookrightarrow [y^2 + (4z + 6)y]_0^2 = 4 + 2(4z + 6) = 8z + 16 \\ &= \int_0^2 (8z + 16) dz = (4z^2 + 16z)|_0^2 = 16 + 32 = \boxed{48.} \end{aligned}$$

Numbering the surfaces as in Fig. 1.29:

- (i)  $d\mathbf{a} = dy dz \hat{\mathbf{x}}, x = 2$ .  $\mathbf{v} \cdot d\mathbf{a} = 2y dy dz$ .  $\int \mathbf{v} \cdot d\mathbf{a} = \iint 2y dy dz = 2y^2 \Big|_0^2 = 8$ .  
(ii)  $d\mathbf{a} = -dy dz \hat{\mathbf{x}}, x = 0$ .  $\mathbf{v} \cdot d\mathbf{a} = 0$ .  $\int \mathbf{v} \cdot d\mathbf{a} = 0$ .  
(iii)  $d\mathbf{a} = dx dz \hat{\mathbf{y}}, y = 2$ .  $\mathbf{v} \cdot d\mathbf{a} = 4z dx dz$ .  $\int \mathbf{v} \cdot d\mathbf{a} = \iint 4z dx dz = 16$ .  
(iv)  $d\mathbf{a} = -dx dz \hat{\mathbf{y}}, y = 0$ .  $\mathbf{v} \cdot d\mathbf{a} = 0$ .  $\int \mathbf{v} \cdot d\mathbf{a} = 0$ .  
(v)  $d\mathbf{a} = dx dy \hat{\mathbf{z}}, z = 2$ .  $\mathbf{v} \cdot d\mathbf{a} = 6x dx dy$ .  $\int \mathbf{v} \cdot d\mathbf{a} = 24$ .  
(vi)  $d\mathbf{a} = -dx dy \hat{\mathbf{z}}, z = 0$ .  $\mathbf{v} \cdot d\mathbf{a} = 0$ .  $\int \mathbf{v} \cdot d\mathbf{a} = 0$ .  
 $\Rightarrow \int \mathbf{v} \cdot d\mathbf{a} = 8 + 16 + 24 = 48 \checkmark$


**Problem 1.34**

$$\nabla \times \mathbf{v} = \hat{\mathbf{x}}(0 - 2y) + \hat{\mathbf{y}}(0 - 3z) + \hat{\mathbf{z}}(0 - x) = -2y \hat{\mathbf{x}} - 3z \hat{\mathbf{y}} - x \hat{\mathbf{z}}.$$


$d\mathbf{a} = dy dz \hat{\mathbf{x}}$ , if we agree that the path integral shall run counterclockwise. So

$$(\nabla \times \mathbf{v}) \cdot d\mathbf{a} = -2y dy dz.$$

$$\begin{aligned} \int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} &= \int \left\{ \int_0^{2-z} (-2y) dy \right\} dz \\ &\quad \hookrightarrow y^2 \Big|_0^{2-z} = -(2-z)^2 \\ &= - \int_0^2 (4 - 4z + z^2) dz = - \left( 4z - 2z^2 + \frac{z^3}{3} \right) \Big|_0^2 \\ &= - \left( 8 - 8 + \frac{8}{3} \right) = \boxed{-\frac{8}{3}} \end{aligned}$$



Meanwhile,  $\mathbf{v} \cdot d\mathbf{l} = (xy)dx + (2yz)dy + (3zx)dz$ . There are three segments.



- (1)  $x = z = 0$ ;  $dx = dz = 0$ .  $y : 0 \rightarrow 2$ .  $\int \mathbf{v} \cdot d\mathbf{l} = 0$ .  
(2)  $x = 0$ ;  $z = 2 - y$ ;  $dx = 0$ ,  $dz = -dy$ ,  $y : 2 \rightarrow 0$ .  $\mathbf{v} \cdot d\mathbf{l} = 2yz dy$ .  
 $\int \mathbf{v} \cdot d\mathbf{l} = \int_2^0 2y(2-y)dy = - \int_0^2 (4y - 2y^2)dy = - \left( 2y^2 - \frac{2}{3}y^3 \right) \Big|_0^2 = - \left( 8 - \frac{2}{3} \cdot 8 \right) = -\frac{8}{3}$ .  
(3)  $x = y = 0$ ;  $dx = dy = 0$ ;  $z : 2 \rightarrow 0$ .  $\mathbf{v} \cdot d\mathbf{l} = 0$ .  $\int \mathbf{v} \cdot d\mathbf{l} = 0$ . So  $\oint \mathbf{v} \cdot d\mathbf{l} = -\frac{8}{3}$ .  $\checkmark$

**Problem 1.35**

By Corollary 1,  $\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a}$  should equal  $\frac{4}{3}$ .  $\nabla \times \mathbf{v} = (4z^2 - 2x)\hat{\mathbf{x}} + 2z\hat{\mathbf{z}}$ .

- (i)  $d\mathbf{a} = dy dz \hat{\mathbf{x}}$ ,  $x = 1$ ;  $y, z : 0 \rightarrow 1$ .  $(\nabla \times \mathbf{v}) \cdot d\mathbf{a} = (4z^2 - 2)dy dz$ ;  $\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = \int_0^1 (4z^2 - 2)dz$   
 $= \left( \frac{4}{3}z^3 - 2z \right) \Big|_0^1 = \frac{4}{3} - 2 = -\frac{2}{3}$ .  
(ii)  $d\mathbf{a} = -dx dy \hat{\mathbf{z}}$ ,  $z = 0$ ;  $x, y : 0 \rightarrow 1$ .  $(\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 0$ ;  $\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 0$ .  
(iii)  $d\mathbf{a} = dx dz \hat{\mathbf{y}}$ ,  $y = 1$ ;  $x, z : 0 \rightarrow 1$ .  $(\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 0$ ;  $\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 0$ .  
(iv)  $d\mathbf{a} = -dx dz \hat{\mathbf{y}}$ ,  $y = 0$ ;  $x, z : 0 \rightarrow 1$ .  $(\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 0$ ;  $\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 0$ .  
(v)  $d\mathbf{a} = dx dy \hat{\mathbf{z}}$ ,  $z = 1$ ;  $x, y : 0 \rightarrow 1$ .  $(\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 2 dx dy$ ;  $\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 2$ .  
 $\Rightarrow \int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = -\frac{2}{3} + 2 = \frac{4}{3}$ .  $\checkmark$

**Problem 1.36**

- (a) Use the product rule
- $\nabla \times (f\mathbf{A}) = f(\nabla \times \mathbf{A}) - \mathbf{A} \times (\nabla f)$
- :

$$\int_S f(\nabla \times \mathbf{A}) \cdot d\mathbf{a} = \int_S \nabla \times (f\mathbf{A}) \cdot d\mathbf{a} + \int_S [\mathbf{A} \times (\nabla f)] \cdot d\mathbf{a} = \oint_{\mathcal{P}} f\mathbf{A} \cdot d\mathbf{l} + \int_S [\mathbf{A} \times (\nabla f)] \cdot d\mathbf{a}. \quad \text{qed}$$

(I used Stokes' theorem in the last step.)

- (b) Use the product rule
- $\nabla \cdot (\mathbf{A} \times \mathbf{B}) = \mathbf{B} \cdot (\nabla \times \mathbf{A}) - \mathbf{A} \cdot (\nabla \times \mathbf{B})$
- :

$$\int_V \mathbf{B} \cdot (\nabla \times \mathbf{A}) d\tau = \int_V \nabla \cdot (\mathbf{A} \times \mathbf{B}) d\tau + \int_V \mathbf{A} \cdot (\nabla \times \mathbf{B}) d\tau = \oint_S (\mathbf{A} \times \mathbf{B}) \cdot d\mathbf{a} + \int_V \mathbf{A} \cdot (\nabla \times \mathbf{B}) d\tau. \quad \text{qed}$$

(I used the divergence theorem in the last step.)

**Problem 1.37**

$$r = \sqrt{x^2 + y^2 + z^2}; \quad \theta = \cos^{-1} \left( \frac{z}{\sqrt{x^2 + y^2 + z^2}} \right); \quad \phi = \tan^{-1} \left( \frac{y}{x} \right).$$

**Problem 1.38**

There are many ways to do this one—probably the most illuminating way is to work it out by trigonometry from Fig. 1.36. The most systematic approach is to study the expression:

$$\mathbf{r} = x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}} = r \sin \theta \cos \phi \hat{\mathbf{x}} + r \sin \theta \sin \phi \hat{\mathbf{y}} + r \cos \theta \hat{\mathbf{z}}.$$

If I only vary  $r$  slightly, then  $d\mathbf{r} = \frac{\partial \mathbf{r}}{\partial r}(\mathbf{r})dr$  is a short vector pointing in the direction of increase in  $r$ . To make it a unit vector, I must divide by its length. Thus:

$$\hat{\mathbf{r}} = \frac{\frac{\partial \mathbf{r}}{\partial r}}{\left| \frac{\partial \mathbf{r}}{\partial r} \right|}; \quad \hat{\boldsymbol{\theta}} = \frac{\frac{\partial \mathbf{r}}{\partial \theta}}{\left| \frac{\partial \mathbf{r}}{\partial \theta} \right|}; \quad \hat{\boldsymbol{\phi}} = \frac{\frac{\partial \mathbf{r}}{\partial \phi}}{\left| \frac{\partial \mathbf{r}}{\partial \phi} \right|}.$$

$$\begin{aligned} \frac{\partial \mathbf{r}}{\partial r} &= \sin \theta \cos \phi \hat{\mathbf{x}} + \sin \theta \sin \phi \hat{\mathbf{y}} + \cos \theta \hat{\mathbf{z}}; \quad \left| \frac{\partial \mathbf{r}}{\partial r} \right|^2 = \sin^2 \theta \cos^2 \phi + \sin^2 \theta \sin^2 \phi + \cos^2 \theta = 1. \\ \frac{\partial \mathbf{r}}{\partial \theta} &= r \cos \theta \cos \phi \hat{\mathbf{x}} + r \cos \theta \sin \phi \hat{\mathbf{y}} - r \sin \theta \hat{\mathbf{z}}; \quad \left| \frac{\partial \mathbf{r}}{\partial \theta} \right|^2 = r^2 \cos^2 \theta \cos^2 \phi + r^2 \cos^2 \theta \sin^2 \phi + r^2 \sin^2 \theta = r^2. \\ \frac{\partial \mathbf{r}}{\partial \phi} &= -r \sin \theta \sin \phi \hat{\mathbf{x}} + r \sin \theta \cos \phi \hat{\mathbf{y}}; \quad \left| \frac{\partial \mathbf{r}}{\partial \phi} \right|^2 = r^2 \sin^2 \theta \sin^2 \phi + r^2 \sin^2 \theta \cos^2 \phi = r^2 \sin^2 \theta. \end{aligned}$$

$$\Rightarrow \begin{cases} \hat{\mathbf{r}} = \sin \theta \cos \phi \hat{\mathbf{x}} + \sin \theta \sin \phi \hat{\mathbf{y}} + \cos \theta \hat{\mathbf{z}}. \\ \hat{\boldsymbol{\theta}} = \cos \theta \cos \phi \hat{\mathbf{x}} + \cos \theta \sin \phi \hat{\mathbf{y}} - \sin \theta \hat{\mathbf{z}}. \\ \hat{\boldsymbol{\phi}} = -\sin \phi \hat{\mathbf{x}} + \cos \phi \hat{\mathbf{y}}. \end{cases}$$

$$\text{Check: } \hat{\mathbf{r}} \cdot \hat{\mathbf{r}} = \sin^2 \theta (\cos^2 \phi + \sin^2 \phi) + \cos^2 \theta = \sin^2 \theta + \cos^2 \theta = 1, \quad \checkmark \\ \hat{\boldsymbol{\theta}} \cdot \hat{\boldsymbol{\phi}} = -\cos \theta \sin \phi \cos \phi + \cos \theta \sin \phi \cos \phi = 0, \quad \checkmark \quad \text{etc.}$$

$$\begin{aligned} \sin \theta \hat{\mathbf{r}} &= \sin^2 \theta \cos \phi \hat{\mathbf{x}} + \sin^2 \theta \sin \phi \hat{\mathbf{y}} + \sin \theta \cos \theta \hat{\mathbf{z}}. \\ \cos \theta \hat{\boldsymbol{\theta}} &= \cos^2 \theta \cos \phi \hat{\mathbf{x}} + \cos^2 \theta \sin \phi \hat{\mathbf{y}} - \sin \theta \cos \theta \hat{\mathbf{z}}. \end{aligned}$$

Add these:

$$\begin{aligned} (1) \quad \sin \theta \hat{\mathbf{r}} + \cos \theta \hat{\boldsymbol{\theta}} &= +\cos \phi \hat{\mathbf{x}} + \sin \phi \hat{\mathbf{y}}; \\ (2) \quad \hat{\boldsymbol{\phi}} &= -\sin \phi \hat{\mathbf{x}} + \cos \phi \hat{\mathbf{y}}. \end{aligned}$$

Multiply (1) by  $\cos \phi$ , (2) by  $\sin \phi$ , and subtract:

$$\hat{\mathbf{x}} = \sin \theta \cos \phi \hat{\mathbf{r}} + \cos \theta \cos \phi \hat{\boldsymbol{\theta}} - \sin \phi \hat{\boldsymbol{\phi}}.$$

Multiply (1) by  $\sin \phi$ , (2) by  $\cos \phi$ , and add:

$$\hat{\mathbf{y}} = \sin \theta \sin \phi \hat{\mathbf{r}} + \cos \theta \sin \phi \hat{\boldsymbol{\theta}} + \cos \phi \hat{\boldsymbol{\phi}}.$$

$$\cos \theta \hat{\mathbf{r}} = \sin \theta \cos \theta \cos \phi \hat{\mathbf{x}} + \sin \theta \cos \theta \sin \phi \hat{\mathbf{y}} + \cos^2 \theta \hat{\mathbf{z}}.$$

$$\sin \theta \hat{\boldsymbol{\theta}} = \sin \theta \cos \theta \cos \phi \hat{\mathbf{x}} + \sin \theta \cos \theta \sin \phi \hat{\mathbf{y}} - \sin^2 \theta \hat{\mathbf{z}}.$$

Subtract these:

$$\hat{\mathbf{z}} = \cos \theta \hat{\mathbf{r}} - \sin \theta \hat{\boldsymbol{\theta}}.$$

### Problem 1.39

$$(a) \nabla \cdot \mathbf{v}_1 = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 r^2) = \frac{1}{r^2} 4r^3 = 4r$$

$$\int (\nabla \cdot \mathbf{v}_1) d\tau = \int (4r) (r^2 \sin \theta dr d\theta d\phi) = (4) \int_0^R r^3 dr \int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\phi = (4) \left( \frac{R^4}{4} \right) (2)(2\pi) = 4\pi R^4$$

$$\int \mathbf{v}_1 \cdot d\mathbf{a} = \int (r^2 \hat{\mathbf{r}}) \cdot (r^2 \sin \theta d\theta d\phi \hat{\mathbf{r}}) = r^4 \int_0^\pi \sin \theta d\theta \int_0^{2\pi} d\phi = 4\pi R^4 \quad \checkmark \quad (\text{Note: at surface of sphere } r = R.)$$

$$(b) \nabla \cdot \mathbf{v}_2 = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \frac{1}{r^2}) = 0 \Rightarrow \int (\nabla \cdot \mathbf{v}_2) d\tau = 0$$

$$\int \mathbf{v}_2 \cdot d\mathbf{a} = \int \left( \frac{1}{r^2} \hat{\mathbf{r}} \right) (r^2 \sin \theta d\theta d\phi \hat{\mathbf{r}}) = \int \sin \theta d\theta d\phi = 4\pi.$$

They *don't* agree! The point is that this divergence is zero *except at the origin*, where it blows up, so our calculation of  $\int (\nabla \cdot \mathbf{v}_2)$  is *incorrect*. The right answer is  $4\pi$ .

### Problem 1.40

$$\begin{aligned} \nabla \cdot \mathbf{v} &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 r \cos \theta) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta r \sin \theta) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} (r \sin \theta \cos \phi) \\ &= \frac{1}{r^2} 3r^2 \cos \theta + \frac{1}{r \sin \theta} r 2 \sin \theta \cos \theta + \frac{1}{r \sin \theta} r \sin \theta (-\sin \phi) \\ &= 3 \cos \theta + 2 \cos \theta - \sin \phi = 5 \cos \theta - \sin \phi \end{aligned}$$

$$\int (\nabla \cdot \mathbf{v}) d\tau = \int (5 \cos \theta - \sin \phi) r^2 \sin \theta dr d\theta d\phi = \int_0^R r^2 dr \int_0^{\frac{\pi}{2}} \left[ \int_0^{2\pi} (5 \cos \theta - \sin \phi) d\phi \right] d\theta \sin \theta$$

$$\hookrightarrow 2\pi(5 \cos \theta)$$

$$= \left( \frac{R^3}{3} \right) (10\pi) \int_0^{\frac{\pi}{2}} \sin \theta \cos \theta d\theta$$

$$\hookrightarrow \frac{\sin^2 \theta}{2} \Big|_0^{\frac{\pi}{2}} = \frac{1}{2}$$

$$= \frac{5\pi}{3} R^3.$$

Two surfaces—one the hemisphere:  $d\mathbf{a} = R^2 \sin \theta d\theta d\phi \hat{\mathbf{r}}$ ;  $r = R$ ;  $\phi : 0 \rightarrow 2\pi$ ,  $\theta : 0 \rightarrow \frac{\pi}{2}$ .

$$\int \mathbf{v} \cdot d\mathbf{a} = \int (r \cos \theta) R^2 \sin \theta d\theta d\phi = R^3 \int_0^{\frac{\pi}{2}} \sin \theta \cos \theta d\theta \int_0^{2\pi} d\phi = R^3 \left( \frac{1}{2} \right) (2\pi) = \pi R^3.$$

other the flat bottom:  $d\mathbf{a} = (dr)(r \sin \theta d\phi)(+\hat{\boldsymbol{\theta}}) = r dr d\phi \hat{\boldsymbol{\theta}}$  (here  $\theta = \frac{\pi}{2}$ ).  $r : 0 \rightarrow R$ ,  $\phi : 0 \rightarrow 2\pi$ .

$$\int \mathbf{v} \cdot d\mathbf{a} = \int (r \sin \theta)(r dr d\phi) = \int_0^R r^2 dr \int_0^{2\pi} d\phi = 2\pi \frac{R^3}{3}.$$

$$\text{Total: } \int \mathbf{v} \cdot d\mathbf{a} = \pi R^3 + \frac{2}{3} \pi R^3 = \frac{5}{3} \pi R^3. \quad \checkmark$$

$$\text{Problem 1.41} \quad \nabla t = (\cos \theta + \sin \theta \cos \phi) \hat{\mathbf{r}} + (-\sin \theta + \cos \theta \cos \phi) \hat{\boldsymbol{\theta}} + \frac{1}{\sin \theta} (-\sin \theta \sin \phi) \hat{\boldsymbol{\phi}}$$

$$\begin{aligned} \nabla^2 t &= \nabla \cdot (\nabla t) \\ &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 (\cos \theta + \sin \theta \cos \phi)) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta (-\sin \theta + \cos \theta \cos \phi)) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} (-\sin \phi) \\ &= \frac{1}{r^2} 2r (\cos \theta + \sin \theta \cos \phi) + \frac{1}{r \sin \theta} (-2 \sin \theta \cos \theta + \cos^2 \theta \cos \phi - \sin^2 \theta \cos \phi) - \frac{1}{r \sin \theta} \cos \phi \\ &= \frac{1}{r \sin \theta} [2 \sin \theta \cos \theta + 2 \sin^2 \theta \cos \phi - 2 \sin \theta \cos \theta + \cos^2 \theta \cos \phi - \sin^2 \theta \cos \phi - \cos \phi] \\ &= \frac{1}{r \sin \theta} [(\sin^2 \theta + \cos^2 \theta) \cos \phi - \cos \phi] = 0. \end{aligned}$$

$$\Rightarrow \boxed{\nabla^2 t = 0}$$

Check:  $r \cos \theta = z$ ,  $r \sin \theta \cos \phi = x \Rightarrow$  in Cartesian coordinates  $t = x + z$ . Obviously Laplacian is zero.

Gradient Theorem:  $\int_{\mathbf{a}}^{\mathbf{b}} \nabla t \cdot d\mathbf{l} = t(\mathbf{b}) - t(\mathbf{a})$

Segment 1:  $\theta = \frac{\pi}{2}$ ,  $\phi = 0$ ,  $r : 0 \rightarrow 2$ .  $d\mathbf{l} = dr \hat{\mathbf{r}}$ ;  $\nabla t \cdot d\mathbf{l} = (\cos \theta + \sin \theta \cos \phi)dr = (0 + 1)dr = dr$ .

$$\int \nabla t \cdot d\mathbf{l} = \int_0^2 dr = 2.$$

Segment 2:  $\theta = \frac{\pi}{2}$ ,  $r = 2$ ,  $\phi : 0 \rightarrow \frac{\pi}{2}$ .  $d\mathbf{l} = r \sin \theta d\phi \hat{\phi} = 2 d\phi \hat{\phi}$ .

$$\nabla t \cdot d\mathbf{l} = (-\sin \phi)(2 d\phi) = -2 \sin \phi d\phi. \int \nabla t \cdot d\mathbf{l} = -\int_0^{\frac{\pi}{2}} 2 \sin \phi d\phi = 2 \cos \phi \Big|_0^{\frac{\pi}{2}} = -2.$$

Segment 3:  $r = 2$ ,  $\phi = \frac{\pi}{2}$ ;  $\theta : \frac{\pi}{2} \rightarrow 0$ .

$$d\mathbf{l} = r d\theta \hat{\theta} = 2 d\theta \hat{\theta}; \nabla t \cdot d\mathbf{l} = (-\sin \theta + \cos \theta \cos \phi)(2 d\theta) = -2 \sin \theta d\theta.$$

$$\int \nabla t \cdot d\mathbf{l} = -\int_{\frac{\pi}{2}}^0 2 \sin \theta d\theta = 2 \cos \theta \Big|_{\frac{\pi}{2}}^0 = 2.$$

Total:  $\int_{\mathbf{a}}^{\mathbf{b}} \nabla t \cdot d\mathbf{l} = 2 - 2 + 2 = \boxed{2}$ . Meanwhile,  $t(\mathbf{b}) - t(\mathbf{a}) = [2(1 + 0)] - [0(\quad)] = 2$ . ✓

**Problem 1.42** From Fig. 1.42,  $\hat{\mathbf{s}} = \cos \phi \hat{\mathbf{x}} + \sin \phi \hat{\mathbf{y}}$ ;  $\hat{\phi} = -\sin \phi \hat{\mathbf{x}} + \cos \phi \hat{\mathbf{y}}$ ;  $\hat{\mathbf{z}} = \hat{\mathbf{z}}$

Multiply first by  $\cos \phi$ , second by  $\sin \phi$ , and subtract:

$$\hat{\mathbf{s}} \cos \phi - \hat{\phi} \sin \phi = \cos^2 \phi \hat{\mathbf{x}} + \cos \phi \sin \phi \hat{\mathbf{y}} + \sin^2 \phi \hat{\mathbf{x}} - \sin \phi \cos \phi \hat{\mathbf{y}} = \hat{\mathbf{x}}(\sin^2 \phi + \cos^2 \phi) = \hat{\mathbf{x}}.$$

$$\text{So } \boxed{\hat{\mathbf{x}} = \cos \phi \hat{\mathbf{s}} - \sin \phi \hat{\phi}}.$$

Multiply first by  $\sin \phi$ , second by  $\cos \phi$ , and add:

$$\hat{\mathbf{s}} \sin \phi + \hat{\phi} \cos \phi = \sin \phi \cos \phi \hat{\mathbf{x}} + \sin^2 \phi \hat{\mathbf{y}} - \sin \phi \cos \phi \hat{\mathbf{x}} + \cos^2 \phi \hat{\mathbf{y}} = \hat{\mathbf{y}}(\sin^2 \phi + \cos^2 \phi) = \hat{\mathbf{y}}.$$

$$\text{So } \boxed{\hat{\mathbf{y}} = \sin \phi \hat{\mathbf{s}} + \cos \phi \hat{\phi}}. \quad \boxed{\hat{\mathbf{z}} = \hat{\mathbf{z}}}.$$

**Problem 1.43**

$$\begin{aligned} \text{(a) } \nabla \cdot \mathbf{v} &= \frac{1}{s} \frac{\partial}{\partial s} (s s (2 + \sin^2 \phi)) + \frac{1}{s} \frac{\partial}{\partial \phi} (s \sin \phi \cos \phi) + \frac{\partial}{\partial z} (3z) \\ &= \frac{1}{s} 2s(2 + \sin^2 \phi) + \frac{1}{s} s(\cos^2 \phi - \sin^2 \phi) + 3 \\ &= 4 + 2 \sin^2 \phi + \cos^2 \phi - \sin^2 \phi + 3 \\ &= 4 + \sin^2 \phi + \cos^2 \phi + 3 = \boxed{8}. \end{aligned}$$

$$\text{(b) } \int (\nabla \cdot \mathbf{v}) d\tau = \int (8) s ds d\phi dz = 8 \int_0^2 s ds \int_0^{\frac{\pi}{2}} d\phi \int_0^5 dz = 8(2) \left(\frac{\pi}{2}\right) (5) = \boxed{40\pi}.$$

Meanwhile, the surface integral has five parts:

$$\text{top: } z = 5, d\mathbf{a} = s ds d\phi \hat{\mathbf{z}}; \mathbf{v} \cdot d\mathbf{a} = 3z s ds d\phi = 15s ds d\phi. \int \mathbf{v} \cdot d\mathbf{a} = 15 \int_0^2 s ds \int_0^{\frac{\pi}{2}} d\phi = 15\pi.$$

$$\text{bottom: } z = 0, d\mathbf{a} = -s ds d\phi \hat{\mathbf{z}}; \mathbf{v} \cdot d\mathbf{a} = -3z s ds d\phi = 0. \int \mathbf{v} \cdot d\mathbf{a} = 0.$$

$$\text{back: } \phi = \frac{\pi}{2}, d\mathbf{a} = ds dz \hat{\phi}; \mathbf{v} \cdot d\mathbf{a} = s \sin \phi \cos \phi ds dz = 0. \int \mathbf{v} \cdot d\mathbf{a} = 0.$$

$$\text{left: } \phi = 0, d\mathbf{a} = -ds dz \hat{\phi}; \mathbf{v} \cdot d\mathbf{a} = -s \sin \phi \cos \phi ds dz = 0. \int \mathbf{v} \cdot d\mathbf{a} = 0.$$

$$\text{front: } s = 2, d\mathbf{a} = s d\phi dz \hat{\mathbf{s}}; \mathbf{v} \cdot d\mathbf{a} = s(2 + \sin^2 \phi) s d\phi dz = 4(2 + \sin^2 \phi) d\phi dz.$$

$$\int \mathbf{v} \cdot d\mathbf{a} = 4 \int_0^{\frac{\pi}{2}} (2 + \sin^2 \phi) d\phi \int_0^5 dz = (4)(\pi + \frac{\pi}{4})(5) = 25\pi.$$

$$\text{So } \oint \mathbf{v} \cdot d\mathbf{a} = 15\pi + 25\pi = 40\pi. \checkmark$$

$$\begin{aligned} \text{(c) } \nabla \times \mathbf{v} &= \left( \frac{1}{s} \frac{\partial}{\partial \phi} (3z) - \frac{\partial}{\partial z} (s \sin \phi \cos \phi) \right) \hat{\mathbf{s}} + \left( \frac{\partial}{\partial z} (s(2 + \sin^2 \phi)) - \frac{\partial}{\partial s} (3z) \right) \hat{\phi} \\ &\quad + \frac{1}{s} \left( \frac{\partial}{\partial s} (s^2 \sin \phi \cos \phi) - \frac{\partial}{\partial \phi} (s(2 + \sin^2 \phi)) \right) \hat{\mathbf{z}} \\ &= \frac{1}{s} (2s \sin \phi \cos \phi - s 2 \sin \phi \cos \phi) \hat{\mathbf{z}} = \boxed{\mathbf{0}}. \end{aligned}$$

**Problem 1.44**

(a)  $3(3^2) - 2(3) - 1 = 27 - 6 - 1 = \boxed{20}.$

(b)  $\cos \pi = \boxed{-1}.$

(c)  $\boxed{\text{zero}}.$

(d)  $\ln(-2 + 3) = \ln 1 = \boxed{\text{zero}}.$

**Problem 1.45**

(a)  $\int_{-2}^2 (2x + 3)^{\frac{1}{3}} \delta(x) dx = \frac{1}{3}(0 + 3) = \boxed{1}.$

(b) By Eq. 1.94,  $\delta(1 - x) = \delta(x - 1)$ , so  $1 + 3 + 2 = \boxed{6}.$

(c)  $\int_{-1}^1 9x^2 \frac{1}{3} \delta(x + \frac{1}{3}) dx = 9 \left(-\frac{1}{3}\right)^2 \frac{1}{3} = \boxed{\frac{1}{3}}.$

(d)  $\boxed{1 \text{ (if } a > b), 0 \text{ (if } a < b)}.$

**Problem 1.46**

(a)  $\int_{-\infty}^{\infty} f(x) \left[ x \frac{d}{dx} \delta(x) \right] dx = x f(x) \delta(x) \Big|_{-\infty}^{\infty} - \int_{-\infty}^{\infty} \frac{d}{dx} (x f(x)) \delta(x) dx.$

The first term is zero, since  $\delta(x) = 0$  at  $\pm\infty$ ;  $\frac{d}{dx} (x f(x)) = x \frac{df}{dx} + \frac{dx}{dx} f = x \frac{df}{dx} + f.$

So the integral is  $-\int_{-\infty}^{\infty} \left( x \frac{df}{dx} + f \right) \delta(x) dx = 0 - f(0) = -f(0) = -\int_{-\infty}^{\infty} f(x) \delta(x) dx.$

So,  $x \frac{d}{dx} \delta(x) = -\delta(x).$     qed

(b)  $\int_{-\infty}^{\infty} f(x) \frac{d\theta}{dx} dx = f(x) \theta(x) \Big|_{-\infty}^{\infty} - \int_{-\infty}^{\infty} \frac{df}{dx} \theta(x) dx = f(\infty) - \int_0^{\infty} \frac{df}{dx} dx = f(\infty) - (f(\infty) - f(0))$   
 $= f(0) = \int_{-\infty}^{\infty} f(x) \delta(x) dx.$  So  $\frac{d\theta}{dx} = \delta(x).$     qed

**Problem 1.47**

(a)  $\boxed{\rho(\mathbf{r}) = q\delta^3(\mathbf{r} - \mathbf{r}')}.$  Check:  $\int \rho(\mathbf{r}) d\tau = q \int \delta^3(\mathbf{r} - \mathbf{r}') d\tau = q.$     ✓

(b)  $\boxed{\rho(\mathbf{r}) = q\delta^3(\mathbf{r} - \mathbf{a}) - q\delta^3(\mathbf{r}).}$

(c) Evidently  $\rho(r) = A\delta(r - R).$  To determine the constant  $A$ , we require

$$Q = \int \rho d\tau = \int A\delta(r - R) 4\pi r^2 dr = A 4\pi R^2. \quad \text{So } A = \frac{Q}{4\pi R^2}. \quad \boxed{\rho(r) = \frac{Q}{4\pi R^2} \delta(r - R).}$$

**Problem 1.48**

(a)  $a^2 + \mathbf{a} \cdot \mathbf{a} + a^2 = \boxed{3a^2}.$

(b)  $\int (\mathbf{r} - \mathbf{b})^2 \frac{1}{5^3} \delta^3(\mathbf{r}) d\tau = \frac{1}{125} b^2 = \frac{1}{125} (4^2 + 3^2) = \boxed{\frac{1}{5}}.$

(c)  $c^2 = 25 + 9 + 4 = 38 > 36 = 6^2$ , so  $\mathbf{c}$  is outside  $\mathcal{V}$ , so the integral is  $\boxed{\text{zero}}.$

(d)  $(\mathbf{e} - (2\hat{\mathbf{x}} + 2\hat{\mathbf{y}} + 2\hat{\mathbf{z}}))^2 = (1\hat{\mathbf{x}} + 0\hat{\mathbf{y}} + (-1)\hat{\mathbf{z}})^2 = 1 + 1 = 2 < (1.5)^2 = 2.25$ , so  $\mathbf{e}$  is inside  $\mathcal{V}$ ,  
and hence the integral is  $\mathbf{e} \cdot (\mathbf{d} - \mathbf{e}) = (3, 2, 1) \cdot (-2, 0, 2) = -6 + 0 + 2 = \boxed{-4}.$

**Problem 1.49**

First method: use Eq. 1.99 to write  $J = \int e^{-r} (4\pi\delta^3(\mathbf{r})) d\tau = 4\pi e^{-0} = \boxed{4\pi}.$

Second method: integrating by parts (use Eq. 1.59).

$$\begin{aligned}
J &= - \int_V \frac{\hat{\mathbf{r}}}{r^2} \cdot \nabla(e^{-r}) d\tau + \oint_S e^{-r} \frac{\hat{\mathbf{r}}}{r^2} \cdot d\mathbf{a}. \quad \text{But } \nabla(e^{-r}) = \left(\frac{\partial}{\partial r} e^{-r}\right) \hat{\mathbf{r}} = -e^{-r} \hat{\mathbf{r}}. \\
&= \int \frac{1}{r^2} e^{-r} 4\pi r^2 dr + \int e^{-r} \frac{\hat{\mathbf{r}}}{r^2} \cdot r^2 \sin\theta d\theta d\phi \hat{\mathbf{r}} = 4\pi \int_0^R e^{-r} dr + e^{-R} \int \sin\theta d\theta d\phi \\
&= 4\pi (-e^{-r}) \Big|_0^R + 4\pi e^{-R} = 4\pi (-e^{-R} + e^{-0}) + 4\pi e^{-R} = 4\pi \checkmark \quad (\text{Here } R = \infty, \text{ so } e^{-R} = 0.)
\end{aligned}$$

**Problem 1.50** (a)  $\nabla \cdot \mathbf{F}_1 = \frac{\partial}{\partial x}(0) + \frac{\partial}{\partial y}(0) + \frac{\partial}{\partial z}(x^2) = \boxed{0}$ ;  $\nabla \cdot \mathbf{F}_2 = \frac{\partial x}{\partial x} + \frac{\partial y}{\partial y} + \frac{\partial z}{\partial z} = 1 + 1 + 1 = \boxed{3}$

$$\nabla \times \mathbf{F}_1 = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 0 & 0 & x^2 \end{vmatrix} = -\hat{\mathbf{y}} \frac{\partial}{\partial x}(x^2) = \boxed{-2x\hat{\mathbf{y}}}; \quad \nabla \times \mathbf{F}_2 = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ x & y & z \end{vmatrix} = \boxed{\mathbf{0}}$$

$\mathbf{F}_2$  is a gradient;  $\mathbf{F}_1$  is a curl  $\left[ U_2 = \frac{1}{2}(x^3 + y^2 + z^2) \right]$  would do ( $\mathbf{F}_2 = \nabla U_2$ ).

For  $\mathbf{A}_1$ , we want  $\left(\frac{\partial A_y}{\partial z} - \frac{\partial A_z}{\partial y}\right) = \left(\frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x}\right) = 0$ ;  $\frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} = x^2$ .  $A_y = \frac{x^3}{3}$ ,  $A_x = A_z = 0$  would do it.

$\mathbf{A}_1 = \frac{1}{3}x^2 \hat{\mathbf{y}}$  ( $\mathbf{F}_1 = \nabla \times \mathbf{A}_1$ ). (But these are not unique.)

$$\text{(b) } \nabla \cdot \mathbf{F}_3 = \frac{\partial}{\partial x}(yz) + \frac{\partial}{\partial y}(xz) + \frac{\partial}{\partial z}(xy) = 0; \quad \nabla \times \mathbf{F}_3 = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ yz & xz & xy \end{vmatrix} = \hat{\mathbf{x}}(x-x) + \hat{\mathbf{y}}(y-y) + \hat{\mathbf{z}}(z-z) = \mathbf{0}.$$

So  $\mathbf{F}_3$  can be written as the gradient of a scalar ( $\mathbf{F}_3 = \nabla U_3$ ) and as the curl of a vector ( $\mathbf{F}_3 = \nabla \times \mathbf{A}_3$ ). In fact,  $\left[ U_3 = xyz \right]$  does the job. For the vector potential, we have

$$\left\{ \begin{array}{l} \frac{\partial A_z}{\partial y} - \frac{\partial A_y}{\partial z} = yz, \text{ which suggests } A_z = \frac{1}{4}y^2z + f(x, z); A_y = -\frac{1}{4}yz^2 + g(x, y) \\ \frac{\partial A_x}{\partial z} - \frac{\partial A_z}{\partial x} = xz, \text{ suggesting } A_x = \frac{1}{4}z^2x + h(x, y); A_z = -\frac{1}{4}zx^2 + j(y, z) \\ \frac{\partial A_y}{\partial x} - \frac{\partial A_x}{\partial y} = xy, \text{ so } A_y = \frac{1}{4}x^2y + k(y, z); A_x = -\frac{1}{4}xy^2 + l(x, z) \end{array} \right\}$$

Putting this all together:  $\mathbf{A}_3 = \frac{1}{4} \{ x(z^2 - y^2) \hat{\mathbf{x}} + y(x^2 - z^2) \hat{\mathbf{y}} + z(y^2 - x^2) \hat{\mathbf{z}} \}$  (again, not unique).

### Problem 1.51

(d)  $\Rightarrow$  (a):  $\nabla \times \mathbf{F} = \nabla \times (-\nabla U) = \mathbf{0}$  (Eq. 1.44 – curl of gradient is always zero).

(a)  $\Rightarrow$  (c):  $\oint \mathbf{F} \cdot d\mathbf{l} = \int (\nabla \times \mathbf{F}) \cdot d\mathbf{a} = 0$  (Eq. 1.57–Stokes' theorem).

(c)  $\Rightarrow$  (b):  $\int_a^b \mathbf{F} \cdot d\mathbf{l} - \int_a^b \mathbf{F} \cdot d\mathbf{l} = \int_a^b \mathbf{F} \cdot d\mathbf{l} + \int_b^a \mathbf{F} \cdot d\mathbf{l} = \oint \mathbf{F} \cdot d\mathbf{l} = 0$ , so

$$\int_a^b \mathbf{F} \cdot d\mathbf{l} = \int_a^b \mathbf{F} \cdot d\mathbf{l}.$$

(b)  $\Rightarrow$  (c): same as (c)  $\Rightarrow$  (b), only in reverse; (c)  $\Rightarrow$  (a): same as (a)  $\Rightarrow$  (c).

### Problem 1.52

(d)  $\Rightarrow$  (a):  $\nabla \cdot \mathbf{F} = \nabla \cdot (\nabla \times \mathbf{W}) = 0$  (Eq 1.46—divergence of curl is always zero).

(a)  $\Rightarrow$  (c):  $\oint \mathbf{F} \cdot d\mathbf{a} = \int (\nabla \cdot \mathbf{F}) d\tau = 0$  (Eq. 1.56—divergence theorem).



(c)  $\Rightarrow$  (b):  $\int_I \mathbf{F} \cdot d\mathbf{a} - \int_{II} \mathbf{F} \cdot d\mathbf{a} = \oint \mathbf{F} \cdot d\mathbf{a} = 0$ , so

$$\int_I \mathbf{F} \cdot d\mathbf{a} = \int_{II} \mathbf{F} \cdot d\mathbf{a}.$$

(Note: sign change because for  $\oint \mathbf{F} \cdot d\mathbf{a}$ ,  $d\mathbf{a}$  is *outward*, whereas for surface II it is *inward*.)

(b)  $\Rightarrow$  (c): same as (c)  $\Rightarrow$  (b), in reverse; (c)  $\Rightarrow$  (a): same as (a)  $\Rightarrow$  (c).

### Problem 1.53

In Prob. 1.15 we found that  $\nabla \cdot \mathbf{v}_a = 0$ ; in Prob. 1.18 we found that  $\nabla \times \mathbf{v}_c = \mathbf{0}$ . So

$\mathbf{v}_c$  can be written as the gradient of a scalar;  $\mathbf{v}_a$  can be written as the curl of a vector.

(a) To find  $t$ :

$$(1) \quad \frac{\partial t}{\partial x} = y^2 \Rightarrow t = y^2 x + f(y, z)$$

$$(2) \quad \frac{\partial t}{\partial y} = (2xy + z^2)$$

$$(3) \quad \frac{\partial t}{\partial z} = 2yz$$

From (1) & (3) we get  $\frac{\partial f}{\partial z} = 2yz \Rightarrow f = yz^2 + g(y) \Rightarrow t = y^2 x + yz^2 + g(y)$ , so  $\frac{\partial t}{\partial y} = 2xy + z^2 + \frac{\partial g}{\partial y} = 2xy + z^2$  (from (2))  $\Rightarrow \frac{\partial g}{\partial y} = 0$ . We may as well pick  $g = 0$ ; then  $t = xy^2 + yz^2$ .

(b) To find  $\mathbf{W}$ :  $\frac{\partial W_z}{\partial y} - \frac{\partial W_y}{\partial z} = x^2$ ;  $\frac{\partial W_x}{\partial z} - \frac{\partial W_z}{\partial x} = 3z^2 x$ ;  $\frac{\partial W_y}{\partial x} - \frac{\partial W_x}{\partial y} = -2xz$ .

Pick  $W_x = 0$ ; then

$$\frac{\partial W_z}{\partial x} = -3xz^2 \Rightarrow W_z = -\frac{3}{2}x^2 z^2 + f(y, z)$$

$$\frac{\partial W_y}{\partial x} = -2xz \Rightarrow W_y = -x^2 z + g(y, z).$$

$\frac{\partial W_z}{\partial y} - \frac{\partial W_y}{\partial z} = \frac{\partial f}{\partial y} + x^2 - \frac{\partial g}{\partial z} = x^2 \Rightarrow \frac{\partial f}{\partial y} - \frac{\partial g}{\partial z} = 0$ . May as well pick  $f = g = 0$ .

$$\mathbf{W} = -x^2 z \hat{\mathbf{y}} - \frac{3}{2} x^2 z^2 \hat{\mathbf{z}}.$$

$$\text{Check: } \nabla \times \mathbf{W} = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 0 & -x^2 z & -\frac{3}{2} x^2 z^2 \end{vmatrix} = \hat{\mathbf{x}} (x^2) + \hat{\mathbf{y}} (3xz^2) + \hat{\mathbf{z}} (-2xz). \checkmark$$

You can add any gradient ( $\nabla t$ ) to  $\mathbf{W}$  without changing its curl, so this answer is far from unique. Some other solutions:

$$\mathbf{W} = xz^3 \hat{\mathbf{x}} - x^2 z \hat{\mathbf{y}};$$

$$\mathbf{W} = (2xyz + xz^3) \hat{\mathbf{x}} + x^2 y \hat{\mathbf{z}};$$

$$\mathbf{W} = xyz \hat{\mathbf{x}} - \frac{1}{2} x^2 z \hat{\mathbf{y}} + \frac{1}{2} x^2 (y - 3z^2) \hat{\mathbf{z}}.$$

**Problem 1.54**

$$\begin{aligned}
\nabla \cdot \mathbf{v} &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 r^2 \cos \theta) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta r^2 \cos \phi) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} (-r^2 \cos \theta \sin \phi) \\
&= \frac{1}{r^2} 4r^3 \cos \theta + \frac{1}{r \sin \theta} \cos \theta r^2 \cos \phi + \frac{1}{r \sin \theta} (-r^2 \cos \theta \cos \phi) \\
&= \frac{r \cos \theta}{\sin \theta} [4 \sin \theta + \cos \phi - \cos \phi] = 4r \cos \theta.
\end{aligned}$$

$$\begin{aligned}
\int (\nabla \cdot \mathbf{v}) d\tau &= \int (4r \cos \theta) r^2 \sin \theta dr d\theta d\phi = 4 \int_0^R r^3 dr \int_0^{\pi/2} \cos \theta \sin \theta d\theta \int_0^{\pi/2} d\phi \\
&= (R^4) \left( \frac{1}{2} \right) \left( \frac{\pi}{2} \right) = \boxed{\frac{\pi R^4}{4}}.
\end{aligned}$$

Surface consists of four parts:

(1) *Curved*:  $d\mathbf{a} = R^2 \sin \theta d\theta d\phi \hat{\mathbf{r}}$ ;  $r = R$ .  $\mathbf{v} \cdot d\mathbf{a} = (R^2 \cos \theta) (R^2 \sin \theta d\theta d\phi)$ .

$$\int \mathbf{v} \cdot d\mathbf{a} = R^4 \int_0^{\pi/2} \cos \theta \sin \theta d\theta \int_0^{\pi/2} d\phi = R^4 \left( \frac{1}{2} \right) \left( \frac{\pi}{2} \right) = \frac{\pi R^4}{4}.$$

(2) *Left*:  $d\mathbf{a} = -r dr d\theta \hat{\phi}$ ;  $\phi = 0$ .  $\mathbf{v} \cdot d\mathbf{a} = (r^2 \cos \theta \sin \phi) (r dr d\theta) = 0$ .  $\int \mathbf{v} \cdot d\mathbf{a} = 0$ .

(3) *Back*:  $d\mathbf{a} = r dr d\theta \hat{\phi}$ ;  $\phi = \pi/2$ .  $\mathbf{v} \cdot d\mathbf{a} = (-r^2 \cos \theta \sin \phi) (r dr d\theta) = -r^3 \cos \theta dr d\theta$ .

$$\int \mathbf{v} \cdot d\mathbf{a} = \int_0^R r^3 dr \int_0^{\pi/2} \cos \theta d\theta = -\left( \frac{1}{4} R^4 \right) (+1) = -\frac{1}{4} R^4.$$

(4) *Bottom*:  $d\mathbf{a} = r \sin \theta dr d\phi \hat{\theta}$ ;  $\theta = \pi/2$ .  $\mathbf{v} \cdot d\mathbf{a} = (r^2 \cos \phi) (r dr d\phi)$ .

$$\int \mathbf{v} \cdot d\mathbf{a} = \int_0^R r^3 dr \int_0^{\pi/2} \cos \phi d\phi = \frac{1}{4} R^4.$$

$$\text{Total: } \oint \mathbf{v} \cdot d\mathbf{a} = \pi R^4/4 + 0 - \frac{1}{4} R^4 + \frac{1}{4} R^4 = \frac{\pi R^4}{4}. \quad \checkmark$$

**Problem 1.55**

$$\nabla \times \mathbf{v} = \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ ay & bx & 0 \end{vmatrix} = \hat{\mathbf{z}}(b-a). \quad \text{So } \int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = (b-a)\pi R^2.$$

$\mathbf{v} \cdot d\mathbf{l} = (ay \hat{\mathbf{x}} + bx \hat{\mathbf{y}}) \cdot (dx \hat{\mathbf{x}} + dy \hat{\mathbf{y}} + dz \hat{\mathbf{z}}) = ay dx + bx dy$ ;  $x^2 + y^2 = R^2 \Rightarrow 2x dx + 2y dy = 0$ ,  
so  $dy = -(x/y) dx$ . So  $\mathbf{v} \cdot d\mathbf{l} = ay dx + bx(-x/y) dx = \frac{1}{y} (ay^2 - bx^2) dx$ .

For the “upper” semicircle,  $y = \sqrt{R^2 - x^2}$ , so  $\mathbf{v} \cdot d\mathbf{l} = \frac{a(R^2 - x^2) - bx^2}{\sqrt{R^2 - x^2}} dx$ .

$$\begin{aligned} \int \mathbf{v} \cdot d\mathbf{l} &= \int_R^{-R} \frac{aR^2 - (a+b)x^2}{\sqrt{R^2 - x^2}} dx = \left\{ aR^2 \sin^{-1}\left(\frac{x}{R}\right) - (a+b) \left[ -\frac{x}{2} \sqrt{R^2 - x^2} + \frac{R^2}{2} \sin^{-1}\left(\frac{x}{R}\right) \right] \right\} \Big|_{+R}^{-R} \\ &= \frac{1}{2} R^2 (a-b) \sin^{-1}(x/R) \Big|_{+R}^{-R} = \frac{1}{2} R^2 (a-b) (\sin^{-1}(-1) - \sin^{-1}(+1)) = \frac{1}{2} R^2 (a-b) \left( -\frac{\pi}{2} - \frac{\pi}{2} \right) \\ &= \frac{1}{2} \pi R^2 (b-a). \end{aligned}$$

And the same for the lower semicircle ( $y$  changes sign, but the limits on the integral are reversed) so

$$\oint \mathbf{v} \cdot d\mathbf{l} = \pi R^2 (b-a). \quad \checkmark$$

### Problem 1.56

$$(1) \quad x = z = 0; \quad dx = dz = 0; \quad y : 0 \rightarrow 1. \quad \mathbf{v} \cdot d\mathbf{l} = (yz^2) dy = 0; \quad \int \mathbf{v} \cdot d\mathbf{l} = 0.$$

$$(2) \quad x = 0; \quad z = 2 - 2y; \quad dz = -2 dy; \quad y : 1 \rightarrow 0. \quad \mathbf{v} \cdot d\mathbf{l} = (yz^2) dy + (3y + z) dz = y(2 - 2y)^2 dy - (3y + 2 - 2y) 2 dy;$$

$$\int \mathbf{v} \cdot d\mathbf{l} = 2 \int_1^0 (2y^3 - 4y^2 + y - 2) dy = 2 \left( \frac{y^4}{2} - \frac{4y^3}{3} + \frac{y^2}{2} - 2y \right) \Big|_1^0 = \frac{14}{3}.$$

$$(3) \quad x = y = 0; \quad dx = dy = 0; \quad z : 2 \rightarrow 0. \quad \mathbf{v} \cdot d\mathbf{l} = (3y + z) dz = z dz;$$

$$\int \mathbf{v} \cdot d\mathbf{l} = \int_2^0 z dz = \frac{z^2}{2} \Big|_2^0 = -2.$$

$$\text{Total: } \oint \mathbf{v} \cdot d\mathbf{l} = 0 + \frac{14}{3} - 2 = \boxed{\frac{8}{3}}.$$

Meanwhile, Stokes' theorem says  $\oint \mathbf{v} \cdot d\mathbf{l} = \int (\nabla \times \mathbf{v}) \cdot d\mathbf{a}$ . Here  $d\mathbf{a} = dy dz \hat{\mathbf{x}}$ , so all we need is  $(\nabla \times \mathbf{v})_x = \frac{\partial}{\partial y}(3y + z) - \frac{\partial}{\partial z}(yz^2) = 3 - 2yz$ . Therefore

$$\begin{aligned} \int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} &= \int \int (3 - 2yz) dy dz = \int_0^1 \left[ \int_0^{2-2y} (3 - 2yz) dz \right] dy \\ &= \int_0^1 [3(2 - 2y) - 2y \frac{1}{2} (2 - 2y)^2] dy = \int_0^1 (-4y^3 + 8y^2 - 10y + 6) dy \\ &= \left( -y^4 + \frac{8}{3}y^3 - 5y^2 + 6y \right) \Big|_0^1 = -1 + \frac{8}{3} - 5 + 6 = \frac{8}{3}. \quad \checkmark \end{aligned}$$

### Problem 1.57

Start at the origin.

$$(1) \quad \theta = \frac{\pi}{2}, \quad \phi = 0; \quad r : 0 \rightarrow 1. \quad \mathbf{v} \cdot d\mathbf{l} = (r \cos^2 \theta) (dr) = 0. \quad \int \mathbf{v} \cdot d\mathbf{l} = 0.$$

$$(2) \quad r = 1, \quad \theta = \frac{\pi}{2}; \quad \phi : 0 \rightarrow \pi/2. \quad \mathbf{v} \cdot d\mathbf{l} = (3r)(r \sin \theta d\phi) = 3 d\phi. \quad \int \mathbf{v} \cdot d\mathbf{l} = 3 \int_0^{\pi/2} d\phi = \frac{3\pi}{2}.$$

(3)  $\phi = \frac{\pi}{2}$ ;  $r \sin \theta = y = 1$ , so  $r = \frac{1}{\sin \theta}$ ,  $dr = \frac{-1}{\sin^2 \theta} \cos \theta d\theta$ ,  $\theta : \frac{\pi}{2} \rightarrow \theta_0 \equiv \tan^{-1}(1/2)$ .

$$\begin{aligned} \mathbf{v} \cdot d\mathbf{l} &= (r \cos^2 \theta) (dr) - (r \cos \theta \sin \theta)(r d\theta) = \frac{\cos^2 \theta}{\sin \theta} \left( -\frac{\cos \theta}{\sin^2 \theta} \right) d\theta - \frac{\cos \theta \sin \theta}{\sin^2 \theta} d\theta \\ &= -\left( \frac{\cos^3 \theta}{\sin^3 \theta} + \frac{\cos \theta}{\sin \theta} \right) d\theta = -\frac{\cos \theta}{\sin \theta} \left( \frac{\cos^2 \theta + \sin^2 \theta}{\sin^2 \theta} \right) d\theta = -\frac{\cos \theta}{\sin^3 \theta} d\theta. \end{aligned}$$

Therefore

$$\int \mathbf{v} \cdot d\mathbf{l} = - \int_{\pi/2}^{\theta_0} \frac{\cos \theta}{\sin^3 \theta} d\theta = \frac{1}{2 \sin^2 \theta} \Big|_{\pi/2}^{\theta_0} = \frac{1}{2 \cdot (1/5)} - \frac{1}{2 \cdot (1)} = \frac{5}{2} - \frac{1}{2} = 2.$$

(4)  $\theta = \theta_0$ ,  $\phi = \frac{\pi}{2}$ ;  $r : \sqrt{5} \rightarrow 0$ .  $\mathbf{v} \cdot d\mathbf{l} = (r \cos^2 \theta) (dr) = \frac{4}{5} r dr$ .

$$\int \mathbf{v} \cdot d\mathbf{l} = \frac{4}{5} \int_{\sqrt{5}}^0 r dr = \frac{4}{5} \frac{r^2}{2} \Big|_{\sqrt{5}}^0 = -\frac{4}{5} \cdot \frac{5}{2} = -2.$$

Total:

$$\oint \mathbf{v} \cdot d\mathbf{l} = 0 + \frac{3\pi}{2} + 2 - 2 = \boxed{\frac{3\pi}{2}}.$$

Stokes' theorem says this should equal  $\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a}$

$$\begin{aligned} \nabla \times \mathbf{v} &= \frac{1}{r \sin \theta} \left[ \frac{\partial}{\partial \theta} (\sin \theta 3r) - \frac{\partial}{\partial \phi} (-r \sin \theta \cos \theta) \right] \hat{\mathbf{r}} + \frac{1}{r} \left[ \frac{1}{\sin \theta} \frac{\partial}{\partial \phi} (r \cos^2 \theta) - \frac{\partial}{\partial r} (r 3r) \right] \hat{\boldsymbol{\theta}} \\ &\quad + \frac{1}{r} \left[ \frac{\partial}{\partial r} (-rr \cos \theta \sin \theta) - \frac{\partial}{\partial \theta} (r \cos^2 \theta) \right] \hat{\boldsymbol{\phi}} \\ &= \frac{1}{r \sin \theta} [3r \cos \theta] \hat{\mathbf{r}} + \frac{1}{r} [-6r] \hat{\boldsymbol{\theta}} + \frac{1}{r} [-2r \cos \theta \sin \theta + 2r \cos \theta \sin \theta] \hat{\boldsymbol{\phi}} \\ &= 3 \cot \theta \hat{\mathbf{r}} - 6 \hat{\boldsymbol{\theta}}. \end{aligned}$$

(1) Back face:  $d\mathbf{a} = -r dr d\theta \hat{\boldsymbol{\phi}}$ ;  $(\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 0$ .  $\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 0$ .

(2) Bottom:  $d\mathbf{a} = -r \sin \theta dr d\phi \hat{\boldsymbol{\theta}}$ ;  $(\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 6r \sin \theta dr d\phi$ .  $\theta = \frac{\pi}{2}$ , so  $(\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 6r dr d\phi$

$$\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = \int_0^1 6r dr \int_0^{\pi/2} d\phi = 6 \cdot \frac{1}{2} \cdot \frac{\pi}{2} = \frac{3\pi}{2}. \quad \checkmark$$

### Problem 1.58

$\mathbf{v} \cdot d\mathbf{l} = y dz$ .

(1) Left side:  $z = a - x$ ;  $dz = -dx$ ;  $y = 0$ . Therefore  $\int \mathbf{v} \cdot d\mathbf{l} = 0$ .

(2) Bottom:  $dz = 0$ . Therefore  $\int \mathbf{v} \cdot d\mathbf{l} = 0$ .

$$(3) \text{ Back: } z = a - \frac{1}{2}y; \quad dz = -\frac{1}{2}dy; \quad y: 2a \rightarrow 0. \quad \int \mathbf{v} \cdot d\mathbf{l} = \int_{2a}^0 y \left(-\frac{1}{2}dy\right) = -\frac{1}{2}\frac{y^2}{2}\bigg|_{2a}^0 = \frac{4a^2}{4} = \boxed{a^2}.$$

Meanwhile,  $\nabla \times \mathbf{v} = \hat{\mathbf{x}}$ , so  $\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a}$  is the projection of this surface on the  $xy$  plane  $= \frac{1}{2} \cdot a \cdot 2a = a^2$ .  $\checkmark$

### Problem 1.59

$$\begin{aligned} \nabla \cdot \mathbf{v} &= \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 r^2 \sin \theta) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} (\sin \theta 4r^2 \cos \theta) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} (r^2 \tan \theta) \\ &= \frac{1}{r^2} 4r^3 \sin \theta + \frac{1}{r \sin \theta} 4r^2 (\cos^2 \theta - \sin^2 \theta) = \frac{4r}{\sin \theta} (\sin^2 \theta + \cos^2 \theta - \sin^2 \theta) \\ &= 4r \frac{\cos^2 \theta}{\sin \theta}. \end{aligned}$$

$$\begin{aligned} \int (\nabla \cdot \mathbf{v}) d\tau &= \int \left( 4r \frac{\cos^2 \theta}{\sin \theta} \right) (r^2 \sin \theta dr d\theta d\phi) = \int_0^R 4r^3 dr \int_0^{\pi/6} \cos^2 \theta d\theta \int_0^{2\pi} d\phi = (R^4) (2\pi) \left[ \frac{\theta}{2} + \frac{\sin 2\theta}{4} \right]_0^{\pi/6} \\ &= 2\pi R^4 \left( \frac{\pi}{12} + \frac{\sin 60^\circ}{4} \right) = \frac{\pi R^4}{6} \left( \pi + 3 \frac{\sqrt{3}}{2} \right) = \boxed{\frac{\pi R^4}{12} (2\pi + 3\sqrt{3})}. \end{aligned}$$

Surface consists of two parts:

- (1) *The ice cream:*  $r = R$ ;  $\phi: 0 \rightarrow 2\pi$ ;  $\theta: 0 \rightarrow \pi/6$ ;  $d\mathbf{a} = R^2 \sin \theta d\theta d\phi \hat{\mathbf{r}}$ ;  $\mathbf{v} \cdot d\mathbf{a} = (R^2 \sin \theta) (R^2 \sin \theta d\theta d\phi) = R^4 \sin^2 \theta d\theta d\phi$ .

$$\int \mathbf{v} \cdot d\mathbf{a} = R^4 \int_0^{\pi/6} \sin^2 \theta d\theta \int_0^{2\pi} d\phi = (R^4) (2\pi) \left[ \frac{1}{2}\theta - \frac{1}{4}\sin 2\theta \right]_0^{\pi/6} = 2\pi R^4 \left( \frac{\pi}{12} - \frac{1}{4}\sin 60^\circ \right) = \frac{\pi R^4}{6} \left( \pi - 3 \frac{\sqrt{3}}{2} \right)$$

- (2) *The cone:*  $\theta = \frac{\pi}{6}$ ;  $\phi: 0 \rightarrow 2\pi$ ;  $r: 0 \rightarrow R$ ;  $d\mathbf{a} = r \sin \theta d\phi dr \hat{\boldsymbol{\theta}} = \frac{\sqrt{3}}{2} r dr d\phi \hat{\boldsymbol{\theta}}$ ;  $\mathbf{v} \cdot d\mathbf{a} = \sqrt{3} r^3 dr d\phi$

$$\int \mathbf{v} \cdot d\mathbf{a} = \sqrt{3} \int_0^R r^3 dr \int_0^{2\pi} d\phi = \sqrt{3} \cdot \frac{R^4}{4} \cdot 2\pi = \frac{\sqrt{3}}{2} \pi R^4.$$

$$\text{Therefore } \int \mathbf{v} \cdot d\mathbf{a} = \frac{\pi R^4}{2} \left( \frac{\pi}{3} - \frac{\sqrt{3}}{2} + \sqrt{3} \right) = \frac{\pi R^4}{12} (2\pi + 3\sqrt{3}). \quad \checkmark.$$

### Problem 1.60

(a) Corollary 2 says  $\oint (\nabla T) \cdot d\mathbf{l} = 0$ . Stokes' theorem says  $\oint (\nabla T) \cdot d\mathbf{l} = \int [\nabla \times (\nabla T)] \cdot d\mathbf{a}$ . So  $\int [\nabla \times (\nabla T)] \cdot d\mathbf{a} = 0$ , and since this is true for *any* surface, the integrand must vanish:  $\nabla \times (\nabla T) = \mathbf{0}$ , confirming Eq. 1.44.

(b) Corollary 2 says  $\oint (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = 0$ . Divergence theorem says  $\oint (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = \int \nabla \cdot (\nabla \times \mathbf{v}) d\tau$ . So  $\int \nabla \cdot (\nabla \times \mathbf{v}) d\tau = 0$ , and since this is true for *any* volume, the integrand must vanish:  $\nabla \cdot (\nabla \times \mathbf{v}) = 0$ , confirming Eq. 1.46.

### Problem 1.61

(a) Divergence theorem:  $\oint \mathbf{v} \cdot d\mathbf{a} = \int (\nabla \cdot \mathbf{v}) d\tau$ . Let  $\mathbf{v} = \mathbf{c}T$ , where  $\mathbf{c}$  is a constant vector. Using product rule #5 in front cover:  $\nabla \cdot \mathbf{v} = \nabla \cdot (\mathbf{c}T) = T(\nabla \cdot \mathbf{c}) + \mathbf{c} \cdot (\nabla T)$ . But  $\mathbf{c}$  is constant so  $\nabla \cdot \mathbf{c} = 0$ . Therefore we have:  $\int \mathbf{c} \cdot (\nabla T) d\tau = \int T \mathbf{c} \cdot d\mathbf{a}$ . Since  $\mathbf{c}$  is constant, take it outside the integrals:  $\mathbf{c} \cdot \int \nabla T d\tau = \mathbf{c} \cdot \int T d\mathbf{a}$ . But  $\mathbf{c}$

is *any* constant vector—in particular, it could be  $\hat{\mathbf{x}}$ , or  $\hat{\mathbf{y}}$ , or  $\hat{\mathbf{z}}$ —so each *component* of the integral on left equals corresponding component on the right, and hence

$$\int \nabla T d\tau = \int T d\mathbf{a}. \quad \text{qed}$$

(b) Let  $\mathbf{v} \rightarrow (\mathbf{v} \times \mathbf{c})$  in divergence theorem. Then  $\int \nabla \cdot (\mathbf{v} \times \mathbf{c}) d\tau = \int (\mathbf{v} \times \mathbf{c}) \cdot d\mathbf{a}$ . Product rule #6  $\Rightarrow \nabla \cdot (\mathbf{v} \times \mathbf{c}) = \mathbf{c} \cdot (\nabla \times \mathbf{v}) - \mathbf{v} \cdot (\nabla \times \mathbf{c}) = \mathbf{c} \cdot (\nabla \times \mathbf{v})$ . (Note:  $\nabla \times \mathbf{c} = \mathbf{0}$ , since  $\mathbf{c}$  is constant.) Meanwhile vector identity (1) says  $d\mathbf{a} \cdot (\mathbf{v} \times \mathbf{c}) = \mathbf{c} \cdot (d\mathbf{a} \times \mathbf{v}) = -\mathbf{c} \cdot (\mathbf{v} \times d\mathbf{a})$ . Thus  $\int \mathbf{c} \cdot (\nabla \times \mathbf{v}) d\tau = -\int \mathbf{c} \cdot (\mathbf{v} \times d\mathbf{a})$ . Take  $\mathbf{c}$  outside, and again let  $\mathbf{c}$  be  $\hat{\mathbf{x}}$ ,  $\hat{\mathbf{y}}$ ,  $\hat{\mathbf{z}}$  then:

$$\int (\nabla \times \mathbf{v}) d\tau = -\int \mathbf{v} \times d\mathbf{a}. \quad \text{qed}$$

(c) Let  $\mathbf{v} = T\nabla U$  in divergence theorem:  $\int \nabla \cdot (T\nabla U) d\tau = \int T\nabla U \cdot d\mathbf{a}$ . Product rule #5  $\Rightarrow \nabla \cdot (T\nabla U) = T\nabla \cdot (\nabla U) + (\nabla U) \cdot (\nabla T) = T\nabla^2 U + (\nabla U) \cdot (\nabla T)$ . Therefore

$$\int (T\nabla^2 U + (\nabla U) \cdot (\nabla T)) d\tau = \int (T\nabla U) \cdot d\mathbf{a}. \quad \text{qed}$$

(d) Rewrite (c) with  $T \leftrightarrow U$ :  $\int (U\nabla^2 T + (\nabla T) \cdot (\nabla U)) d\tau = \int (U\nabla T) \cdot d\mathbf{a}$ . Subtract this from (c), noting that the  $(\nabla U) \cdot (\nabla T)$  terms cancel:

$$\int (T\nabla^2 U - U\nabla^2 T) d\tau = \int (T\nabla U - U\nabla T) \cdot d\mathbf{a}. \quad \text{qed}$$

(e) Stokes' theorem:  $\int (\nabla \times \mathbf{v}) \cdot d\mathbf{a} = \oint \mathbf{v} \cdot d\mathbf{l}$ . Let  $\mathbf{v} = \mathbf{c}T$ . By Product Rule #7:  $\nabla \times (\mathbf{c}T) = T(\nabla \times \mathbf{c}) - \mathbf{c} \times (\nabla T) = -\mathbf{c} \times (\nabla T)$  (since  $\mathbf{c}$  is constant). Therefore,  $-\int (\mathbf{c} \times (\nabla T)) \cdot d\mathbf{a} = \oint \mathbf{c}T \cdot d\mathbf{l}$ . Use vector identity #1 to rewrite the first term  $(\mathbf{c} \times (\nabla T)) \cdot d\mathbf{a} = \mathbf{c} \cdot (\nabla T \times d\mathbf{a})$ . So  $-\int \mathbf{c} \cdot (\nabla T \times d\mathbf{a}) = \oint \mathbf{c} \cdot T d\mathbf{l}$ . Pull  $\mathbf{c}$  outside, and let  $\mathbf{c} \rightarrow \hat{\mathbf{x}}$ ,  $\hat{\mathbf{y}}$ , and  $\hat{\mathbf{z}}$  to prove:

$$\int \nabla T \times d\mathbf{a} = -\oint T d\mathbf{l}. \quad \text{qed}$$

### Problem 1.62

(a)  $d\mathbf{a} = R^2 \sin \theta d\theta d\phi \hat{\mathbf{r}}$ . Let the surface be the northern hemisphere. The  $\hat{\mathbf{x}}$  and  $\hat{\mathbf{y}}$  components clearly integrate to zero, and the  $\hat{\mathbf{z}}$  component of  $\hat{\mathbf{r}}$  is  $\cos \theta$ , so

$$\mathbf{a} = \int R^2 \sin \theta \cos \theta d\theta d\phi \hat{\mathbf{z}} = 2\pi R^2 \hat{\mathbf{z}} \int_0^{\pi/2} \sin \theta \cos \theta d\theta = 2\pi R^2 \hat{\mathbf{z}} \frac{\sin^2 \theta}{2} \Big|_0^{\pi/2} = \boxed{\pi R^2 \hat{\mathbf{z}}}.$$

(b) Let  $T = 1$  in Prob. 1.61(a). Then  $\nabla T = 0$ , so  $\oint d\mathbf{a} = 0$ . qed

(c) This follows from (b). For suppose  $\mathbf{a}_1 \neq \mathbf{a}_2$ ; then if you put them together to make a closed surface,  $\oint d\mathbf{a} = \mathbf{a}_1 - \mathbf{a}_2 \neq 0$ .

(d) For one such triangle,  $d\mathbf{a} = \frac{1}{2}(\mathbf{r} \times d\mathbf{l})$  (since  $\mathbf{r} \times d\mathbf{l}$  is the area of the parallelogram, and the direction is perpendicular to the surface), so for the entire conical surface,  $\mathbf{a} = \frac{1}{2} \oint \mathbf{r} \times d\mathbf{l}$ .

(e) Let  $T = \mathbf{c} \cdot \mathbf{r}$ , and use product rule #4:  $\nabla T = \nabla(\mathbf{c} \cdot \mathbf{r}) = \mathbf{c} \times (\nabla \times \mathbf{r}) + (\mathbf{c} \cdot \nabla)\mathbf{r}$ . But  $\nabla \times \mathbf{r} = 0$ , and  $(\mathbf{c} \cdot \nabla)\mathbf{r} = (c_x \frac{\partial}{\partial x} + c_y \frac{\partial}{\partial y} + c_z \frac{\partial}{\partial z})(x\hat{\mathbf{x}} + y\hat{\mathbf{y}} + z\hat{\mathbf{z}}) = c_x \hat{\mathbf{x}} + c_y \hat{\mathbf{y}} + c_z \hat{\mathbf{z}} = \mathbf{c}$ . So Prob. 1.61(e) says

$$\oint T d\mathbf{l} = \oint (\mathbf{c} \cdot \mathbf{r}) d\mathbf{l} = -\int (\nabla T) \times d\mathbf{a} = -\int \mathbf{c} \times d\mathbf{a} = -\mathbf{c} \times \int d\mathbf{a} = -\mathbf{c} \times \mathbf{a} = \mathbf{a} \times \mathbf{c}. \quad \text{qed}$$

**Problem 1.63**

(1)

$$\nabla \cdot \mathbf{v} = \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \cdot \frac{1}{r} \right) = \frac{1}{r^2} \frac{\partial}{\partial r} (r) = \boxed{\frac{1}{r^2}}.$$

For a sphere of radius  $R$ :

$$\left. \begin{aligned} \int \mathbf{v} \cdot d\mathbf{a} &= \int \left( \frac{1}{R} \hat{\mathbf{r}} \right) \cdot (R^2 \sin \theta d\theta d\phi \hat{\mathbf{r}}) = R \int \sin \theta d\theta d\phi = 4\pi R. \\ \int (\nabla \cdot \mathbf{v}) d\tau &= \int \left( \frac{1}{r^2} \right) (r^2 \sin \theta dr d\theta d\phi) = \left( \int_0^R dr \right) \left( \int \sin \theta d\theta d\phi \right) = 4\pi R. \end{aligned} \right\} \text{So divergence theorem checks.}$$

Evidently there is *no* delta function at the origin.

$$\nabla \times (r^n \hat{\mathbf{r}}) = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 r^n) = \frac{1}{r^2} \frac{\partial}{\partial r} (r^{n+2}) = \frac{1}{r^2} (n+2) r^{n+1} = \boxed{(n+2) r^{n-1}}$$

(except for  $n = -2$ , for which we already know (Eq. 1.99) that the divergence is  $4\pi\delta^3(\mathbf{r})$ ).

- (2) *Geometrically*, it should be zero. Likewise, the curl in the spherical coordinates obviously gives zero. To be certain there is no lurking delta function here, we integrate over a sphere of radius  $R$ , using Prob. 1.61(b): If  $\nabla \times (r^n \hat{\mathbf{r}}) = \mathbf{0}$ , then  $\int (\nabla \times \mathbf{v}) d\tau = \mathbf{0} \stackrel{?}{=} -\oint \mathbf{v} \times d\mathbf{a}$ . But  $\mathbf{v} = r^n \hat{\mathbf{r}}$  and  $d\mathbf{a} = R^2 \sin \theta d\theta d\phi \hat{\mathbf{r}}$  are both in the  $\hat{\mathbf{r}}$  directions, so  $\mathbf{v} \times d\mathbf{a} = \mathbf{0}$ . ✓

**Problem 1.64**

(a) Since the argument is not a function of angle, Eq. 1.73 says

$$\begin{aligned} D &= -\frac{1}{4\pi} \frac{1}{r^2} \frac{d}{dr} \left[ r^2 \left( -\frac{1}{2} \right) \frac{2r}{(r^2 + \epsilon^2)^{3/2}} \right] = \frac{1}{4\pi r^2} \frac{d}{dr} \left[ \frac{r^3}{(r^2 + \epsilon^2)^{3/2}} \right] \\ &= \frac{1}{4\pi r^2} \left[ \frac{3r^2}{(r^2 + \epsilon^2)^{3/2}} - \frac{3}{2} \frac{r^3 \cdot 2r}{(r^2 + \epsilon^2)^{5/2}} \right] = \frac{1}{4\pi r^2} \frac{3r^2}{(r^2 + \epsilon^2)^{5/2}} (r^2 + \epsilon^2 - r^2) = \frac{3\epsilon^2}{4\pi (r^2 + \epsilon^2)^{5/2}}. \quad \checkmark \end{aligned}$$

(b) Setting  $r \rightarrow 0$ :

$$D(0, \epsilon) = \frac{3\epsilon^2}{4\pi\epsilon^5} = \frac{3}{4\pi\epsilon^3},$$

which goes to infinity as  $\epsilon \rightarrow 0$ . ✓(c) From (a) it is clear that  $D(r, 0) = 0$  for  $r \neq 0$ . ✓

(d)

$$\int D(r, \epsilon) 4\pi r^2 dr = 3\epsilon^2 \int_0^\infty \frac{r^2}{(r^2 + \epsilon^2)^{5/2}} dr = 3\epsilon^2 \left( \frac{1}{3\epsilon^2} \right) = 1. \quad \checkmark$$

(I looked up the integral.) Note that (b), (c), and (d) are the defining conditions for  $\delta^3(\mathbf{r})$ .

## Chapter 2

# Electrostatics

### Problem 2.1

(a) Zero.

(b)  $F = \frac{1}{4\pi\epsilon_0} \frac{qQ}{r^2}$ , where  $r$  is the distance from center to each numeral.  $\mathbf{F}$  points *toward* the missing  $q$ .

*Explanation:* by superposition, this is equivalent to (a), with an extra  $-q$  at 6 o'clock—since the force of all twelve is zero, the net force is that of  $-q$  only.

(c) Zero.


(d)  $\frac{1}{4\pi\epsilon_0} \frac{qQ}{r^2}$ , pointing toward the missing  $q$ . Same reason as (b). Note, however, that if you explained (b) as a cancellation in pairs of opposite charges (1 o'clock against 7 o'clock; 2 against 8, etc.), with one unpaired  $q$  doing the job, then you'll need a *different* explanation for (d).

### Problem 2.2

This time the “vertical” components cancel, leaving


$$\mathbf{E} = \frac{1}{4\pi\epsilon_0} 2 \frac{q}{z^{\frac{3}{2}}} \sin \theta \hat{\mathbf{x}}, \text{ or}$$

$$\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{qd}{\left(z^2 + \left(\frac{d}{2}\right)^2\right)^{3/2}} \hat{\mathbf{x}}.$$



From far away, ( $z \gg d$ ), the field goes like  $\mathbf{E} \approx \frac{1}{4\pi\epsilon_0} \frac{qd}{z^3} \hat{\mathbf{z}}$ , which, as we shall see, is the field of a *dipole*. (If we set  $d \rightarrow 0$ , we get  $\mathbf{E} = \mathbf{0}$ , as is appropriate; to the extent that this configuration looks like a single point charge from far away, the net charge is zero, so  $\mathbf{E} \rightarrow \mathbf{0}$ .)



**Problem 2.3**

$$\begin{aligned}
 E_z &= \frac{1}{4\pi\epsilon_0} \int_0^L \frac{\lambda dx}{r^2} \cos \theta; \quad (r^2 = z^2 + x^2; \cos \theta = \frac{z}{r}) \\
 &= \frac{1}{4\pi\epsilon_0} \lambda z \int_0^L \frac{1}{(z^2 + x^2)^{3/2}} dx \\
 &= \frac{1}{4\pi\epsilon_0} \lambda z \left[ \frac{1}{z^2} \frac{x}{\sqrt{z^2 + x^2}} \right]_0^L = \frac{1}{4\pi\epsilon_0} \frac{\lambda}{z} \frac{L}{\sqrt{z^2 + L^2}}. \\
 E_x &= -\frac{1}{4\pi\epsilon_0} \int_0^L \frac{\lambda dx}{r^2} \sin \theta = -\frac{1}{4\pi\epsilon_0} \lambda \int \frac{x dx}{(x^2 + z^2)^{3/2}} \\
 &= -\frac{1}{4\pi\epsilon_0} \lambda \left[ -\frac{1}{\sqrt{x^2 + z^2}} \right]_0^L = -\frac{1}{4\pi\epsilon_0} \lambda \left[ \frac{1}{z} - \frac{1}{\sqrt{z^2 + L^2}} \right].
 \end{aligned}$$

$$\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{\lambda}{z} \left[ \left( -1 + \frac{z}{\sqrt{z^2 + L^2}} \right) \hat{\mathbf{x}} + \left( \frac{L}{\sqrt{z^2 + L^2}} \right) \hat{\mathbf{z}} \right].$$

For  $z \gg L$  you expect it to look like a point charge  $q = \lambda L$ :  $\mathbf{E} \rightarrow \frac{1}{4\pi\epsilon_0} \frac{\lambda L}{z^2} \hat{\mathbf{z}}$ . It checks, for with  $z \gg L$  the  $\hat{\mathbf{x}}$  term  $\rightarrow 0$ , and the  $\hat{\mathbf{z}}$  term  $\rightarrow \frac{1}{4\pi\epsilon_0} \frac{\lambda L}{z^2} \hat{\mathbf{z}}$ .


**Problem 2.4**

From Ex. 2.2, with  $L \rightarrow \frac{a}{2}$  and  $z \rightarrow \sqrt{z^2 + \left(\frac{a}{2}\right)^2}$  (distance from center of edge to  $P$ ), field of *one* edge is:

$$E_1 = \frac{1}{4\pi\epsilon_0} \frac{\lambda a}{\sqrt{z^2 + \frac{a^2}{4}} \sqrt{z^2 + \frac{a^2}{4} + \frac{a^2}{4}}}.$$

There are 4 sides, and we want vertical components only, so multiply by  $4 \cos \theta = 4 \frac{z}{\sqrt{z^2 + \frac{a^2}{4}}}$ :

$$\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{4\lambda a z}{\left(z^2 + \frac{a^2}{4}\right) \sqrt{z^2 + \frac{a^2}{2}}} \hat{\mathbf{z}}.$$

**Problem 2.5**

“Horizontal” components cancel, leaving:  $\mathbf{E} = \frac{1}{4\pi\epsilon_0} \left\{ \int \frac{\lambda dl}{r^2} \cos \theta \right\} \hat{\mathbf{z}}$ . Here,  $r^2 = r^2 + z^2$ ,  $\cos \theta = \frac{z}{r}$  (both constants), while  $\int dl = 2\pi r$ . So

$$\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{\lambda(2\pi r)z}{(r^2 + z^2)^{3/2}} \hat{\mathbf{z}}.$$

**Problem 2.6**

Break it into rings of radius  $r$ , and thickness  $dr$ , and use Prob. 2.5 to express the field of each ring. Total charge of a ring is  $\sigma \cdot 2\pi r \cdot dr = \lambda \cdot 2\pi r$ , so  $\lambda = \sigma dr$  is the “line charge” of each ring.

$$E_{\text{ring}} = \frac{1}{4\pi\epsilon_0} \frac{(\sigma dr)2\pi r z}{(r^2 + z^2)^{3/2}}; \quad E_{\text{disk}} = \frac{1}{4\pi\epsilon_0} 2\pi\sigma z \int_0^R \frac{r}{(r^2 + z^2)^{3/2}} dr.$$

$$\mathbf{E}_{\text{disk}} = \frac{1}{4\pi\epsilon_0} 2\pi\sigma z \left[ \frac{1}{z} - \frac{1}{\sqrt{R^2 + z^2}} \right] \hat{\mathbf{z}}.$$

For  $R \gg z$  the second term  $\rightarrow 0$ , so  $\mathbf{E}_{\text{plane}} = \frac{1}{4\pi\epsilon_0} 2\pi\sigma\hat{\mathbf{z}} = \boxed{\frac{\sigma}{2\epsilon_0}\hat{\mathbf{z}}}$ .

For  $z \gg R$ ,  $\frac{1}{\sqrt{R^2+z^2}} = \frac{1}{z} \left(1 + \frac{R^2}{z^2}\right)^{-1/2} \approx \frac{1}{z} \left(1 - \frac{1}{2} \frac{R^2}{z^2}\right)$ , so  $[\ ] \approx \frac{1}{z} - \frac{1}{z} + \frac{1}{2} \frac{R^2}{z^3} = \frac{R^2}{2z^3}$ ,

and  $E = \frac{1}{4\pi\epsilon_0} \frac{2\pi R^2\sigma}{2z^2} = \frac{1}{4\pi\epsilon_0} \frac{Q}{z^2}$ , where  $Q = \pi R^2\sigma$ .  $\checkmark$

### Problem 2.7


$\mathbf{E}$  is clearly in the  $z$  direction. From the diagram,

$$dq = \sigma da = \sigma R^2 \sin \theta d\theta d\phi,$$

$$r^2 = R^2 + z^2 - 2Rz \cos \theta,$$

$$\cos \psi = \frac{z - R \cos \theta}{r}.$$

So



$$\begin{aligned} E_z &= \frac{1}{4\pi\epsilon_0} \int \frac{\sigma R^2 \sin \theta d\theta d\phi (z - R \cos \theta)}{(R^2 + z^2 - 2Rz \cos \theta)^{3/2}}. \quad \int d\phi = 2\pi. \\ &= \frac{1}{4\pi\epsilon_0} (2\pi R^2 \sigma) \int_0^\pi \frac{(z - R \cos \theta) \sin \theta}{(R^2 + z^2 - 2Rz \cos \theta)^{3/2}} d\theta. \quad \text{Let } u = \cos \theta; du = -\sin \theta d\theta; \left\{ \begin{array}{l} \theta = 0 \Rightarrow u = +1 \\ \theta = \pi \Rightarrow u = -1 \end{array} \right\} \\ &= \frac{1}{4\pi\epsilon_0} (2\pi R^2 \sigma) \int_{-1}^1 \frac{z - Ru}{(R^2 + z^2 - 2Rzu)^{3/2}} du. \quad \text{Integral can be done by partial fractions—or look it up.} \\ &= \frac{1}{4\pi\epsilon_0} (2\pi R^2 \sigma) \left[ \frac{1}{z^2} \frac{zu - R}{\sqrt{R^2 + z^2 - 2Rzu}} \right]_{-1}^1 = \frac{1}{4\pi\epsilon_0} \frac{2\pi R^2 \sigma}{z^2} \left\{ \frac{(z - R)}{|z - R|} - \frac{(-z - R)}{|z + R|} \right\}. \end{aligned}$$

For  $z > R$  (outside the sphere),  $E_z = \frac{1}{4\pi\epsilon_0} \frac{4\pi R^2 \sigma}{z^2} = \frac{1}{4\pi\epsilon_0} \frac{q}{z^2}$ , so  $\mathbf{E} = \boxed{\frac{1}{4\pi\epsilon_0} \frac{q}{z^2} \hat{\mathbf{z}}}$ .

For  $z < R$  (inside),  $E_z = 0$ , so  $\mathbf{E} = \mathbf{0}$ .

### Problem 2.8

According to Prob. 2.7, all shells *interior* to the point (i.e. at smaller  $r$ ) contribute as though their charge were concentrated at the center, while all exterior shells contribute nothing. Therefore:

$$\mathbf{E}(r) = \frac{1}{4\pi\epsilon_0} \frac{Q_{\text{int}}}{r^2} \hat{\mathbf{r}},$$

where  $Q_{\text{int}}$  is the total charge interior to the point. *Outside* the sphere, *all* the charge is interior, so

$$\mathbf{E} = \boxed{\frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \hat{\mathbf{r}}}.$$

*Inside* the sphere, only that fraction of the total which is interior to the point counts:

$$Q_{\text{int}} = \frac{\frac{4}{3}\pi r^3}{\frac{4}{3}\pi R^3} Q = \frac{r^3}{R^3} Q, \quad \text{so} \quad \mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{r^3}{R^3} Q \frac{1}{r^2} \hat{\mathbf{r}} = \boxed{\frac{1}{4\pi\epsilon_0} \frac{Q}{R^3} \mathbf{r}}.$$

### Problem 2.9

(a)  $\rho = \epsilon_0 \nabla \cdot \mathbf{E} = \epsilon_0 \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 \cdot kr^3) = \epsilon_0 \frac{1}{r^2} k(5r^4) = \boxed{5\epsilon_0 k r^2}.$


(b) By Gauss's law:  $Q_{\text{enc}} = \epsilon_0 \oint \mathbf{E} \cdot d\mathbf{a} = \epsilon_0 (kR^3)(4\pi R^2) = \boxed{4\pi\epsilon_0 kR^5}$ .

By direct integration:  $Q_{\text{enc}} = \int \rho d\tau = \int_0^R (5\epsilon_0 k r^2)(4\pi r^2 dr) = 20\pi\epsilon_0 k \int_0^R r^4 dr = 4\pi\epsilon_0 k R^5$ . ✓


**Problem 2.10**

Think of this cube as one of 8 surrounding the charge. Each of the 24 squares which make up the surface of this larger cube gets the same flux as every other one, so:

$$\int_{\text{one face}} \mathbf{E} \cdot d\mathbf{a} = \frac{1}{24} \int_{\text{whole large cube}} \mathbf{E} \cdot d\mathbf{a}.$$




The latter is  $\frac{1}{\epsilon_0}q$ , by Gauss's law. Therefore  $\int_{\text{one face}} \mathbf{E} \cdot d\mathbf{a} = \frac{q}{24\epsilon_0}$ .

**Problem 2.11**

Gaussian surface: *Inside*:  $\oint \mathbf{E} \cdot d\mathbf{a} = E(4\pi r^2) = \frac{1}{\epsilon_0} Q_{\text{enc}} = 0 \Rightarrow \boxed{\mathbf{E} = 0}$ .

Gaussian surface: *Outside*:  $E(4\pi r^2) = \frac{1}{\epsilon_0}(\sigma 4\pi R^2) \Rightarrow \boxed{\mathbf{E} = \frac{\sigma R^2}{\epsilon_0 r^2} \hat{\mathbf{r}}}$ .

} (As in Prob. 2.7.)


**Problem 2.12**

Gaussian surface

$$\oint \mathbf{E} \cdot d\mathbf{a} = E \cdot 4\pi r^2 = \frac{1}{\epsilon_0} Q_{\text{enc}} = \frac{1}{\epsilon_0} \frac{4}{3} \pi r^3 \rho. \quad \text{So}$$

$$\boxed{\mathbf{E} = \frac{1}{3\epsilon_0} \rho r \hat{\mathbf{r}}}$$


Since  $Q_{\text{tot}} = \frac{4}{3} \pi R^3 \rho$ ,  $\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{Q}{R^3} \hat{\mathbf{r}}$  (as in Prob. 2.8).

**Problem 2.13**

Gaussian surface

$$\oint \mathbf{E} \cdot d\mathbf{a} = E \cdot 2\pi s \cdot l = \frac{1}{\epsilon_0} Q_{\text{enc}} = \frac{1}{\epsilon_0} \lambda l. \quad \text{So}$$

$$\boxed{\mathbf{E} = \frac{\lambda}{2\pi\epsilon_0 s} \hat{\mathbf{s}}} \quad (\text{same as Eq. 2.9}).$$

**Problem 2.14**

Gaussian surface

$$\begin{aligned} \oint \mathbf{E} \cdot d\mathbf{a} &= E \cdot 4\pi r^2 = \frac{1}{\epsilon_0} Q_{\text{enc}} = \frac{1}{\epsilon_0} \int \rho d\tau = \frac{1}{\epsilon_0} \int (k\bar{r})(\bar{r}^2 \sin\theta d\bar{r} d\theta d\phi) \\ &= \frac{1}{\epsilon_0} k 4\pi \int_0^r \bar{r}^3 d\bar{r} = \frac{4\pi k}{\epsilon_0} \frac{r^4}{4} = \frac{\pi k}{\epsilon_0} r^4. \end{aligned}$$

$$\therefore \boxed{\mathbf{E} = \frac{1}{4\pi\epsilon_0} \pi k r^2 \hat{\mathbf{r}}}$$


**Problem 2.15**(i)  $Q_{\text{enc}} = 0$ , so  $\mathbf{E} = \mathbf{0}$ .

$$(ii) \oint \mathbf{E} \cdot d\mathbf{a} = E(4\pi r^2) = \frac{1}{\epsilon_0} Q_{\text{enc}} = \frac{1}{\epsilon_0} \int \rho d\tau = \frac{1}{\epsilon_0} \int \frac{k}{\tilde{r}^2} \tilde{r}^2 \sin \theta d\tilde{r} d\theta d\phi$$

$$= \frac{4\pi k}{\epsilon_0} \int_a^r d\tilde{r} = \frac{4\pi k}{\epsilon_0} (r - a) \therefore \mathbf{E} = \frac{k}{\epsilon_0} \left( \frac{r - a}{r^2} \right) \hat{\mathbf{r}}.$$


(iii)  $E(4\pi r^2) = \frac{4\pi k}{\epsilon_0} \int_a^b d\tilde{r} = \frac{4\pi k}{\epsilon_0} (b - a)$ , so

$$\mathbf{E} = \frac{k}{\epsilon_0} \left( \frac{b - a}{r^2} \right) \hat{\mathbf{r}}.$$

**Problem 2.16**


$$\oint \mathbf{E} \cdot d\mathbf{a} = E \cdot 2\pi s \cdot l = \frac{1}{\epsilon_0} Q_{\text{enc}} = \frac{1}{\epsilon_0} \rho \pi s^2 l;$$

$$\mathbf{E} = \frac{\rho s}{2\epsilon_0} \hat{\mathbf{s}}.$$




$$\oint \mathbf{E} \cdot d\mathbf{a} = E \cdot 2\pi s \cdot l = \frac{1}{\epsilon_0} Q_{\text{enc}} = \frac{1}{\epsilon_0} \rho \pi a^2 l;$$

$$\mathbf{E} = \frac{\rho a^2}{2\epsilon_0 s} \hat{\mathbf{s}}.$$




$$\oint \mathbf{E} \cdot d\mathbf{a} = E \cdot 2\pi s \cdot l = \frac{1}{\epsilon_0} Q_{\text{enc}} = 0;$$

$$\mathbf{E} = \mathbf{0}.$$




**Problem 2.17** On the  $xz$  plane  $E = 0$  by symmetry. Set up a Gaussian “pillbox” with one face in this plane and the other at  $y$ .



$$\int \mathbf{E} \cdot d\mathbf{a} = E \cdot A = \frac{1}{\epsilon_0} Q_{\text{enc}} = \frac{1}{\epsilon_0} A y \rho;$$

$$\mathbf{E} = \frac{\rho}{\epsilon_0} y \hat{\mathbf{y}} \quad (\text{for } |y| < d).$$


$$Q_{\text{enc}} = \frac{1}{\epsilon_0} A d \rho \Rightarrow \boxed{\mathbf{E} = \frac{\rho}{\epsilon_0} d \hat{\mathbf{y}}} \text{ (for } y > d \text{)}.$$

**Problem 2.18**

From Prob. 2.12, the field inside the positive sphere is  $\mathbf{E}_+ = \frac{\rho}{3\epsilon_0} \mathbf{r}_+$ , where  $\mathbf{r}_+$  is the vector from the positive center to the point in question. Likewise, the field of the negative sphere is  $-\frac{\rho}{3\epsilon_0} \mathbf{r}_-$ . So the *total* field is

$$\mathbf{E} = \frac{\rho}{3\epsilon_0} (\mathbf{r}_+ - \mathbf{r}_-)$$

But (see diagram)  $\mathbf{r}_+ - \mathbf{r}_- = \mathbf{d}$ . So  $\boxed{\mathbf{E} = \frac{\rho}{3\epsilon_0} \mathbf{d}}$ .

**Problem 2.19**

$$\begin{aligned} \nabla \times \mathbf{E} &= \frac{1}{4\pi\epsilon_0} \nabla \times \int \frac{\hat{\mathbf{z}}}{r^2} \rho d\tau = \frac{1}{4\pi\epsilon_0} \int \left[ \nabla \times \left( \frac{\hat{\mathbf{z}}}{r^2} \right) \right] \rho d\tau \quad (\text{since } \rho \text{ depends on } \mathbf{r}', \text{ not } \mathbf{r}) \\ &= \mathbf{0} \quad (\text{since } \nabla \times \left( \frac{\hat{\mathbf{z}}}{r^2} \right) = \mathbf{0}, \text{ from Prob. 1.63}). \end{aligned}$$

**Problem 2.20**

$$(1) \nabla \times \mathbf{E}_1 = k \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ xy & 2yz & 3zx \end{vmatrix} = k [\hat{\mathbf{x}}(0 - 2y) + \hat{\mathbf{y}}(0 - 3z) + \hat{\mathbf{z}}(0 - x)] \neq \mathbf{0},$$

so  $\mathbf{E}_1$  is an *impossible* electrostatic field.

$$(2) \nabla \times \mathbf{E}_2 = k \begin{vmatrix} \hat{\mathbf{x}} & \hat{\mathbf{y}} & \hat{\mathbf{z}} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ y^2 & 2xy + z^2 & 2yz \end{vmatrix} = k [\hat{\mathbf{x}}(2z - 2z) + \hat{\mathbf{y}}(0 - 0) + \hat{\mathbf{z}}(2y - 2y)] = \mathbf{0},$$

so  $\mathbf{E}_2$  is a *possible* electrostatic field.

Let's go by the indicated path:

$$\mathbf{E} \cdot d\mathbf{l} = (y^2 dx + (2xy + z^2)dy + 2yz dz)k$$


Step I:  $y = z = 0$ ;  $dy = dz = 0$ .  $\mathbf{E} \cdot d\mathbf{l} = ky^2 dx = 0$ .

Step II:  $x = x_0$ ,  $y : 0 \rightarrow y_0$ ,  $z = 0$ .  $dx = dz = 0$ .

$$\begin{aligned} \mathbf{E} \cdot d\mathbf{l} &= k(2x_0 y + z^2)dy = 2kx_0 y dy. \\ \int_{II} \mathbf{E} \cdot d\mathbf{l} &= 2kx_0 \int_0^{y_0} y dy = kx_0 y_0^2. \end{aligned}$$

Step III:  $x = x_0$ ,  $y = y_0$ ,  $z : 0 \rightarrow z_0$ ;  $dx = dy = 0$ .

$$\mathbf{E} \cdot d\mathbf{l} = 2ky_0 z dz = 2ky_0 dz.$$



$$\int_{III} \mathbf{E} \cdot d\mathbf{l} = 2y_0 k \int_0^{z_0} z \, dz = ky_0 z_0^2.$$

$$V(x_0, y_0, z_0) = - \int_0^{(x_0, y_0, z_0)} \mathbf{E} \cdot d\mathbf{l} = -k(x_0 y_0^2 + y_0 z_0^2), \text{ or } \boxed{V(x, y, z) = -k(xy^2 + yz^2)}.$$

$$\text{Check: } -\nabla V = k \left[ \frac{\partial}{\partial x}(xy^2 + yz^2) \hat{\mathbf{x}} + \frac{\partial}{\partial y}(xy^2 + yz^2) \hat{\mathbf{y}} + \frac{\partial}{\partial z}(xy^2 + yz^2) \hat{\mathbf{z}} \right] = k[y^2 \hat{\mathbf{x}} + (2xy + z^2) \hat{\mathbf{y}} + 2yz \hat{\mathbf{z}}] = \mathbf{E}. \checkmark$$

**Problem 2.21**


$$V(r) = - \int_{\infty}^r \mathbf{E} \cdot d\mathbf{l} \quad \begin{cases} \text{Outside the sphere } (r > R) : \mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{\mathbf{r}}. \\ \text{Inside the sphere } (r < R) : \mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{R^3} r \hat{\mathbf{r}}. \end{cases}$$

$$\text{So for } r > R: V(r) = - \int_{\infty}^r \left( \frac{1}{4\pi\epsilon_0} \frac{q}{\bar{r}^2} \right) d\bar{r} = \frac{1}{4\pi\epsilon_0} q \left( \frac{1}{\bar{r}} \right) \Big|_{\infty}^r = \boxed{\frac{q}{4\pi\epsilon_0} \frac{1}{r}},$$

$$\begin{aligned} \text{and for } r < R: V(r) &= - \int_{\infty}^R \left( \frac{1}{4\pi\epsilon_0} \frac{q}{\bar{r}^2} \right) d\bar{r} - \int_R^r \left( \frac{1}{4\pi\epsilon_0} \frac{q}{R^3} \bar{r} \right) d\bar{r} = \frac{q}{4\pi\epsilon_0} \left[ \frac{1}{R} - \frac{1}{R^3} \left( \frac{r^2 - R^2}{2} \right) \right] \\ &= \boxed{\frac{q}{4\pi\epsilon_0} \frac{1}{2R} \left( 3 - \frac{r^2}{R^2} \right)}. \end{aligned}$$

$$\text{When } r > R, \nabla V = \frac{q}{4\pi\epsilon_0} \frac{\partial}{\partial r} \left( \frac{1}{r} \right) \hat{\mathbf{r}} = -\frac{q}{4\pi\epsilon_0} \frac{1}{r^2} \hat{\mathbf{r}}, \text{ so } \mathbf{E} = -\nabla V = \frac{q}{4\pi\epsilon_0} \frac{1}{r^2} \hat{\mathbf{r}}. \checkmark$$

$$\text{When } r < R, \nabla V = \frac{q}{4\pi\epsilon_0} \frac{1}{2R} \frac{\partial}{\partial r} \left( 3 - \frac{r^2}{R^2} \right) \hat{\mathbf{r}} = \frac{q}{4\pi\epsilon_0} \frac{1}{2R} \left( -\frac{2r}{R^2} \right) \hat{\mathbf{r}} = -\frac{q}{4\pi\epsilon_0} \frac{r}{R^3} \hat{\mathbf{r}}; \text{ so } \mathbf{E} = -\nabla V = \frac{1}{4\pi\epsilon_0} \frac{q}{R^3} r \hat{\mathbf{r}}. \checkmark$$



(In the figure,  $r$  is in units of  $R$ , and  $V(r)$  is in units of  $q/4\pi\epsilon_0 R$ .)

**Problem 2.22**

$\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{2\lambda}{s} \hat{\mathbf{s}}$  (Prob. 2.13). In this case we cannot set the reference point at  $\infty$ , since the charge itself extends to  $\infty$ . Let's set it at  $s = a$ . Then

$$V(s) = - \int_a^s \left( \frac{1}{4\pi\epsilon_0} \frac{2\lambda}{\bar{s}} \right) d\bar{s} = \boxed{-\frac{1}{4\pi\epsilon_0} 2\lambda \ln \left( \frac{s}{a} \right)}.$$

(In this form it is clear why  $a = \infty$  would be no good—likewise the other “natural” point,  $a = 0$ .)

$$\nabla V = -\frac{1}{4\pi\epsilon_0} 2\lambda \frac{\partial}{\partial s} \left( \ln \left( \frac{s}{a} \right) \right) \hat{\mathbf{s}} = -\frac{1}{4\pi\epsilon_0} 2\lambda \frac{1}{s} \hat{\mathbf{s}} = -\mathbf{E}. \checkmark$$

**Problem 2.23**

$$\begin{aligned} V(0) &= - \int_{\infty}^0 \mathbf{E} \cdot d\mathbf{l} = - \int_{\infty}^b \left( \frac{k}{\epsilon_0} \frac{(b-a)}{r^2} \right) dr - \int_b^a \left( \frac{k}{\epsilon_0} \frac{(r-a)}{r^2} \right) dr - \int_a^0 (0) dr = \frac{k}{\epsilon_0} \frac{(b-a)}{b} - \frac{k}{\epsilon_0} \left( \ln \left( \frac{a}{b} \right) + a \left( \frac{1}{a} - \frac{1}{b} \right) \right) \\ &= \frac{k}{\epsilon_0} \left\{ 1 - \frac{a}{b} - \ln \left( \frac{a}{b} \right) - 1 + \frac{a}{b} \right\} = \boxed{\frac{k}{\epsilon_0} \ln \left( \frac{b}{a} \right)}. \end{aligned}$$

**Problem 2.24**

Using Eq. 2.22 and the fields from Prob. 2.16:

$$V(b) - V(0) = - \int_0^b \mathbf{E} \cdot d\mathbf{l} = - \int_0^a \mathbf{E} \cdot d\mathbf{l} - \int_a^b \mathbf{E} \cdot d\mathbf{l} = -\frac{\rho}{2\epsilon_0} \int_0^a s \, ds - \frac{\rho a^2}{2\epsilon_0} \int_a^b \frac{1}{s} \, ds$$


$$= -\left(\frac{\rho}{2\epsilon_0}\right) \frac{s^2}{2} \Big|_0^a + \frac{\rho a^2}{2\epsilon_0} \ln s \Big|_a^b = \boxed{-\frac{\rho a^2}{4\epsilon_0} \left(1 + 2 \ln \left(\frac{b}{a}\right)\right)}.$$

**Problem 2.25**

$$(a) \quad V = \frac{1}{4\pi\epsilon_0} \frac{2q}{\sqrt{z^2 + \left(\frac{d}{2}\right)^2}}.$$

$$(b) \quad V = \frac{1}{4\pi\epsilon_0} \int_{-L}^L \frac{\lambda dx}{\sqrt{z^2 + x^2}} = \frac{\lambda}{4\pi\epsilon_0} \ln(x + \sqrt{z^2 + x^2}) \Big|_{-L}^L$$

$$= \frac{\lambda}{4\pi\epsilon_0} \ln \left[ \frac{L + \sqrt{z^2 + L^2}}{-L + \sqrt{z^2 + L^2}} \right] = \frac{\lambda}{2\pi\epsilon_0} \ln \left( \frac{L + \sqrt{z^2 + L^2}}{z} \right).$$



$$(c) \quad V = \frac{1}{4\pi\epsilon_0} \int_0^R \frac{\sigma 2\pi r dr}{\sqrt{r^2 + z^2}} = \frac{1}{4\pi\epsilon_0} 2\pi\sigma (\sqrt{r^2 + z^2}) \Big|_0^R = \boxed{\frac{\sigma}{2\epsilon_0} (\sqrt{R^2 + z^2} - z)}.$$

In each case, by symmetry  $\frac{\partial V}{\partial y} = \frac{\partial V}{\partial x} = 0$ .  $\therefore \mathbf{E} = -\frac{\partial V}{\partial z} \hat{\mathbf{z}}$ .

$$(a) \quad \mathbf{E} = -\frac{1}{4\pi\epsilon_0} 2q \left(-\frac{1}{2}\right) \frac{2z}{\left(z^2 + \left(\frac{d}{2}\right)^2\right)^{3/2}} \hat{\mathbf{z}} = \boxed{\frac{1}{4\pi\epsilon_0} \frac{2qz}{\left(z^2 + \left(\frac{d}{2}\right)^2\right)^{3/2}} \hat{\mathbf{z}}} \text{ (agrees with Ex. 2.1).}$$

$$(b) \quad \mathbf{E} = -\frac{\lambda}{4\pi\epsilon_0} \left\{ \frac{1}{(L + \sqrt{z^2 + L^2})} \frac{1}{2} \frac{1}{\sqrt{z^2 + L^2}} 2z - \frac{1}{(-L + \sqrt{z^2 + L^2})} \frac{1}{2} \frac{1}{\sqrt{z^2 + L^2}} 2z \right\} \hat{\mathbf{z}}$$

$$= -\frac{\lambda}{4\pi\epsilon_0} \frac{z}{\sqrt{z^2 + L^2}} \left\{ \frac{-L + \sqrt{z^2 + L^2} - L - \sqrt{z^2 + L^2}}{(z^2 + L^2) - L^2} \right\} \hat{\mathbf{z}} = \boxed{\frac{2L\lambda}{4\pi\epsilon_0} \frac{1}{z\sqrt{z^2 + L^2}} \hat{\mathbf{z}}} \text{ (agrees with Ex. 2.2).}$$


$$(c) \quad \mathbf{E} = -\frac{\sigma}{2\epsilon_0} \left\{ \frac{1}{2} \frac{1}{\sqrt{R^2 + z^2}} 2z - 1 \right\} \hat{\mathbf{z}} = \boxed{\frac{\sigma}{2\epsilon_0} \left[ 1 - \frac{z}{\sqrt{R^2 + z^2}} \right] \hat{\mathbf{z}}} \text{ (agrees with Prob. 2.6).}$$

If the right-hand charge in (a) is  $-q$ , then  $V = 0$ , which, naively, suggests  $\mathbf{E} = -\nabla V = \mathbf{0}$ , in contradiction with the answer to Prob. 2.2. The point is that we only know  $V$  on the  $z$  axis, and from this we cannot hope to compute  $E_x = -\frac{\partial V}{\partial x}$  or  $E_y = -\frac{\partial V}{\partial y}$ . That was OK in part (a), because we knew from symmetry that  $E_x = E_y = 0$ . But now  $\mathbf{E}$  points in the  $x$  direction, so knowing  $V$  on the  $z$  axis is insufficient to determine  $\mathbf{E}$ .

**Problem 2.26**

$$V(a) = \frac{1}{4\pi\epsilon_0} \int_0^{\sqrt{2}h} \left( \frac{\sigma 2\pi r}{r} \right) dz = \frac{2\pi\sigma}{4\pi\epsilon_0} \frac{1}{\sqrt{2}} (\sqrt{2}h) = \frac{\sigma h}{2\epsilon_0}$$

(where  $r = z/\sqrt{2}$ )




$$\begin{aligned}
V(\mathbf{b}) &= \frac{1}{4\pi\epsilon_0} \int_0^{\sqrt{2}h} \left( \frac{\sigma 2\pi r}{\tilde{r}} \right) dz \quad (\text{where } \tilde{r} = \sqrt{h^2 + z^2 - \sqrt{2}hz}) \\
&= \frac{2\pi\sigma}{4\pi\epsilon_0} \frac{1}{\sqrt{2}} \int_0^{\sqrt{2}h} \frac{z}{\sqrt{h^2 + z^2 - \sqrt{2}hz}} dz \\
&= \frac{\sigma}{2\sqrt{2}\epsilon_0} \left[ \sqrt{h^2 + z^2 - \sqrt{2}hz} + \frac{h}{\sqrt{2}} \ln(2\sqrt{h^2 + z^2 - \sqrt{2}hz} + 2z - \sqrt{2}h) \right] \Big|_0^{\sqrt{2}h} \\
&= \frac{\sigma}{2\sqrt{2}\epsilon_0} \left[ h + \frac{h}{\sqrt{2}} \ln(2h + 2\sqrt{2}h - \sqrt{2}h) - h - \frac{h}{\sqrt{2}} \ln(2h - \sqrt{2}h) \right] \\
&= \frac{\sigma}{2\sqrt{2}\epsilon_0} \frac{h}{\sqrt{2}} [\ln(2h + \sqrt{2}h) - \ln(2h - \sqrt{2}h)] = \frac{\sigma h}{4\epsilon_0} \ln \left( \frac{2 + \sqrt{2}}{2 - \sqrt{2}} \right) = \frac{\sigma h}{4\epsilon_0} \ln \left( \frac{(2 + \sqrt{2})^2}{2} \right) \\
&= \frac{\sigma h}{2\epsilon_0} \ln(1 + \sqrt{2}). \quad \therefore \boxed{V(\mathbf{a}) - V(\mathbf{b}) = \frac{\sigma h}{2\epsilon_0} [1 - \ln(1 + \sqrt{2})]}.
\end{aligned}$$

**Problem 2.27**

Cut the cylinder into slabs, as shown in the figure, and use result of Prob. 2.25c, with  $z \rightarrow x$  and  $\sigma \rightarrow \rho dx$ :

$$\begin{aligned}
V &= \frac{\rho}{2\epsilon_0} \int_{z-L/2}^{z+L/2} (\sqrt{R^2 + x^2} - x) dx \\
&= \frac{\rho}{2\epsilon_0} \frac{1}{2} [x\sqrt{R^2 + x^2} + R^2 \ln(x + \sqrt{R^2 + x^2}) - x^2] \Big|_{z-L/2}^{z+L/2} \\
&= \frac{\rho}{4\epsilon_0} \left\{ \left( z + \frac{L}{2} \right) \sqrt{R^2 + \left( z + \frac{L}{2} \right)^2} - \left( z - \frac{L}{2} \right) \sqrt{R^2 + \left( z - \frac{L}{2} \right)^2} + R^2 \ln \left[ \frac{z + \frac{L}{2} + \sqrt{R^2 + \left( z + \frac{L}{2} \right)^2}}{z - \frac{L}{2} + \sqrt{R^2 + \left( z - \frac{L}{2} \right)^2}} \right] - 2zL \right\}.
\end{aligned}$$

(Note:  $-(z + \frac{L}{2})^2 + (z - \frac{L}{2})^2 = -z^2 - zL - \frac{L^2}{4} + z^2 - zL + \frac{L^2}{4} = -2zL$ .)



$$\begin{aligned}
\mathbf{E} &= -\nabla V = -\hat{\mathbf{z}} \frac{\partial V}{\partial z} = -\frac{\hat{\mathbf{z}}\rho}{4\epsilon_0} \left\{ \sqrt{R^2 + \left( z + \frac{L}{2} \right)^2} + \frac{\left( z + \frac{L}{2} \right)^2}{\sqrt{R^2 + \left( z + \frac{L}{2} \right)^2}} - \sqrt{R^2 + \left( z - \frac{L}{2} \right)^2} - \frac{\left( z - \frac{L}{2} \right)^2}{\sqrt{R^2 + \left( z - \frac{L}{2} \right)^2}} \right. \\
&\quad \left. + R^2 \left[ \frac{1 + \frac{z + \frac{L}{2}}{\sqrt{R^2 + \left( z + \frac{L}{2} \right)^2}}}{z + \frac{L}{2} + \sqrt{R^2 + \left( z + \frac{L}{2} \right)^2}} - \frac{1 + \frac{z - \frac{L}{2}}{\sqrt{R^2 + \left( z - \frac{L}{2} \right)^2}}}{z - \frac{L}{2} + \sqrt{R^2 + \left( z - \frac{L}{2} \right)^2}} \right] - 2L \right\} \\
&\quad \underbrace{\frac{1}{\sqrt{R^2 + \left( z + \frac{L}{2} \right)^2}} - \frac{1}{\sqrt{R^2 + \left( z - \frac{L}{2} \right)^2}}}_{\text{}} \\
\mathbf{E} &= -\frac{\hat{\mathbf{z}}\rho}{4\epsilon_0} \left\{ 2\sqrt{R^2 + \left( z + \frac{L}{2} \right)^2} - 2\sqrt{R^2 + \left( z - \frac{L}{2} \right)^2} - 2L \right\}
\end{aligned}$$



$$= \frac{\rho}{2\epsilon_0} \left[ L - \sqrt{R^2 + \left(z + \frac{L}{2}\right)^2} + \sqrt{R^2 + \left(z - \frac{L}{2}\right)^2} \right] \hat{\mathbf{z}}.$$

**Problem 2.28**

Orient axes so  $P$  is on  $z$  axis.

$$V = \frac{1}{4\pi\epsilon_0} \int \frac{\rho}{\mathcal{L}} d\tau. \quad \left\{ \begin{array}{l} \text{Here } \rho \text{ is constant, } d\tau = r^2 \sin\theta dr d\theta d\phi, \\ \mathcal{L} = \sqrt{z^2 + r^2 - 2rz \cos\theta}. \end{array} \right.$$

$$V = \frac{\rho}{4\pi\epsilon_0} \int \frac{r^2 \sin\theta dr d\theta d\phi}{\sqrt{z^2 + r^2 - 2rz \cos\theta}}; \int_0^{2\pi} d\phi = 2\pi.$$

$$\int_0^\pi \frac{\sin\theta}{\sqrt{z^2 + r^2 - 2rz \cos\theta}} d\theta = \frac{1}{rz} \left( \sqrt{r^2 + z^2 - 2rz \cos\theta} \right) \Big|_0^\pi = \frac{1}{rz} \left( \sqrt{r^2 + z^2 + 2rz} - \sqrt{r^2 + z^2 - 2rz} \right)$$

$$= \frac{1}{rz} (r + z - |r - z|) = \begin{cases} 2/z, & \text{if } r < z, \\ 2/r, & \text{if } r > z. \end{cases}$$

$$\therefore V = \frac{\rho}{4\pi\epsilon_0} \cdot 2\pi \cdot 2 \left\{ \int_0^z \frac{1}{z} r^2 dr + \int_z^R \frac{1}{r} r^2 dr \right\} = \frac{\rho}{\epsilon_0} \left\{ \frac{1}{z} \frac{z^3}{3} + \frac{R^2 - z^2}{2} \right\} = \frac{\rho}{2\epsilon_0} \left( R^2 - \frac{z^2}{3} \right).$$

But  $\rho = \frac{q}{\frac{4}{3}\pi R^3}$ , so  $V(z) = \frac{1}{2\epsilon_0} \frac{3q}{4\pi R^3} \left( R^2 - \frac{z^2}{3} \right) = \frac{q}{8\pi\epsilon_0 R} \left( 3 - \frac{z^2}{R^2} \right)$ ;  $V(r) = \frac{q}{8\pi\epsilon_0 R} \left( 3 - \frac{r^2}{R^2} \right)$ . ✓

**Problem 2.29**


$$\begin{aligned} \nabla^2 V &= \frac{1}{4\pi\epsilon_0} \nabla^2 \int \left( \frac{\rho}{\mathcal{L}} \right) d\tau = \frac{1}{4\pi\epsilon_0} \int \rho(\mathbf{r}') \left( \nabla^2 \frac{1}{\mathcal{L}} \right) d\tau \quad (\text{since } \rho \text{ is a function of } \mathbf{r}', \text{ not } \mathbf{r}) \\ &= \frac{1}{4\pi\epsilon_0} \int \rho(\mathbf{r}') [-4\pi\delta^3(\mathbf{r} - \mathbf{r}')] d\tau = -\frac{1}{\epsilon_0} \rho(\mathbf{r}). \quad \checkmark \end{aligned}$$

**Problem 2.30.**

(a) Ex. 2.5:  $\mathbf{E}_{\text{above}} = \frac{\sigma}{2\epsilon_0} \hat{\mathbf{n}}$ ;  $\mathbf{E}_{\text{below}} = -\frac{\sigma}{2\epsilon_0} \hat{\mathbf{n}}$  ( $\hat{\mathbf{n}}$  always pointing up);  $\mathbf{E}_{\text{above}} - \mathbf{E}_{\text{below}} = \frac{\sigma}{\epsilon_0} \hat{\mathbf{n}}$ . ✓

Ex. 2.6: At each surface,  $E = 0$  one side and  $E = \frac{\sigma}{\epsilon_0}$  other side, so  $\Delta E = \frac{\sigma}{\epsilon_0}$ . ✓

Prob. 2.11:  $\mathbf{E}_{\text{out}} = \frac{\sigma R^2}{\epsilon_0 r^2} \hat{\mathbf{r}} = \frac{\sigma}{\epsilon_0} \hat{\mathbf{r}}$ ;  $\mathbf{E}_{\text{in}} = 0$ ; so  $\Delta \mathbf{E} = \frac{\sigma}{\epsilon_0} \hat{\mathbf{r}}$ . ✓

(b)  Outside:  $\oint \mathbf{E} \cdot d\mathbf{a} = E(2\pi s)l = \frac{1}{\epsilon_0} Q_{\text{enc}} = \frac{\sigma}{\epsilon_0} (2\pi R)l \Rightarrow \mathbf{E} = \frac{\sigma}{\epsilon_0} \frac{R}{s} \hat{\mathbf{s}} = \frac{\sigma}{\epsilon_0} \hat{\mathbf{s}}$  (at surface).  
Inside:  $Q_{\text{enc}} = 0$ , so  $\mathbf{E} = 0$ .  $\therefore \Delta \mathbf{E} = \frac{\sigma}{\epsilon_0} \hat{\mathbf{s}}$ . ✓


(c)  $V_{\text{out}} = \frac{R^2 \sigma}{\epsilon_0 r} = \frac{R\sigma}{\epsilon_0}$  (at surface);  $V_{\text{in}} = \frac{R\sigma}{\epsilon_0}$ ; so  $V_{\text{out}} = V_{\text{in}}$ . ✓

$$\frac{\partial V_{\text{out}}}{\partial r} = -\frac{R^2 \sigma}{\epsilon_0 r^2} = -\frac{\sigma}{\epsilon_0} \quad (\text{at surface}); \quad \frac{\partial V_{\text{in}}}{\partial r} = 0; \quad \text{so } \frac{\partial V_{\text{out}}}{\partial r} - \frac{\partial V_{\text{in}}}{\partial r} = -\frac{\sigma}{\epsilon_0}. \quad \checkmark$$

**Problem 2.31**

$$(a) V = \frac{1}{4\pi\epsilon_0} \sum \frac{q_i}{r_{ij}} = \frac{1}{4\pi\epsilon_0} \left\{ \frac{-q}{a} + \frac{q}{\sqrt{2}a} + \frac{-q}{a} \right\} = \frac{q}{4\pi\epsilon_0 a} \left( -2 + \frac{1}{\sqrt{2}} \right).$$

$$\therefore W_4 = qV = \boxed{\frac{q^2}{4\pi\epsilon_0 a} \left( -2 + \frac{1}{\sqrt{2}} \right)}.$$



$$(b) W_1 = 0, W_2 = \frac{1}{4\pi\epsilon_0} \left( \frac{-q^2}{a} \right); W_3 = \frac{1}{4\pi\epsilon_0} \left( \frac{q^2}{\sqrt{2}a} - \frac{q^2}{a} \right); W_4 = (\text{see (a)}).$$

$$W_{\text{tot}} = \frac{1}{4\pi\epsilon_0} \frac{q^2}{a} \left\{ -1 + \frac{1}{\sqrt{2}} - 1 - 2 + \frac{1}{\sqrt{2}} \right\} = \boxed{\frac{1}{4\pi\epsilon_0} \frac{2q^2}{a} \left( -2 + \frac{1}{\sqrt{2}} \right)}.$$

**Problem 2.32**

Conservation of energy (kinetic plus potential):

$$\frac{1}{2} m_A v_A^2 + \frac{1}{2} m_B v_B^2 + \frac{1}{4\pi\epsilon_0} \frac{q_A q_B}{r} = E.$$

At release  $v_A = v_B = 0$ ,  $r = a$ , so

$$E = \frac{1}{4\pi\epsilon_0} \frac{q_A q_B}{a}.$$

When they are very far apart ( $r \rightarrow \infty$ ) the potential energy is zero, so

$$\frac{1}{2} m_A v_A^2 + \frac{1}{2} m_B v_B^2 = \frac{1}{4\pi\epsilon_0} \frac{q_A q_B}{a}.$$

Meanwhile, conservation of momentum says  $m_A v_A = m_B v_B$ , or  $v_B = (m_A/m_B) v_A$ . So

$$\frac{1}{2} m_A v_A^2 + \frac{1}{2} m_B \left( \frac{m_A}{m_B} \right)^2 v_A^2 = \frac{1}{2} \left( \frac{m_A}{m_B} \right) (m_A + m_B) v_A^2 = \frac{1}{4\pi\epsilon_0} \frac{q_A q_B}{a}.$$

$$\boxed{v_A = \sqrt{\frac{1}{2\pi\epsilon_0} \frac{q_A q_B}{(m_A + m_B)a} \left( \frac{m_A}{m_B} \right)}; \quad v_B = \sqrt{\frac{1}{2\pi\epsilon_0} \frac{q_A q_B}{(m_A + m_B)a} \left( \frac{m_B}{m_A} \right)}}.$$

**Problem 2.33**

From Eq. 2.42, the energy of one charge is

$$W = \frac{1}{2} qV = \frac{1}{2} (2) \sum_{n=1}^{\infty} \frac{1}{4\pi\epsilon_0} \frac{(-1)^n q^2}{na} = \frac{q^2}{4\pi\epsilon_0 a} \sum_{n=1}^{\infty} \frac{(-1)^n}{n}.$$

(The factor of 2 out front counts the charges to the left as well as to the right of  $q$ .) The sum is  $-\ln 2$  (you can get it from the Taylor expansion of  $\ln(1+x)$ ):

$$\ln(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \dots$$

with  $x = 1$ . Evidently  $\boxed{\alpha = \ln 2}$ .

**Problem 2.34**

(a)  $W = \frac{1}{2} \int \rho V d\tau$ . From Prob. 2.21 (or Prob. 2.28):  $V = \frac{\rho}{2\epsilon_0} \left( R^2 - \frac{r^2}{3} \right) = \frac{1}{4\pi\epsilon_0} \frac{q}{2R} \left( 3 - \frac{r^2}{R^2} \right)$

$$\begin{aligned} W &= \frac{1}{2} \rho \frac{1}{4\pi\epsilon_0} \frac{q}{2R} \int_0^R \left( 3 - \frac{r^2}{R^2} \right) 4\pi r^2 dr = \frac{q\rho}{4\epsilon_0 R} \left[ 3 \frac{r^3}{3} - \frac{1}{R^2} \frac{r^5}{5} \right]_0^R = \frac{q\rho}{4\epsilon_0 R} \left( R^3 - \frac{R^3}{5} \right) \\ &= \frac{q\rho}{5\epsilon_0} R^2 = \frac{qR^2}{5\epsilon_0} \frac{q}{\frac{4}{3}\pi R^3} = \boxed{\frac{1}{4\pi\epsilon_0} \left( \frac{3}{5} \frac{q^2}{R} \right)}. \end{aligned}$$

(b)  $W = \frac{\epsilon_0}{2} \int E^2 d\tau$ . Outside ( $r > R$ )  $\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{\mathbf{r}}$ ; Inside ( $r < R$ )  $\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{R^3} r \hat{\mathbf{r}}$ .

$$\begin{aligned} \therefore W &= \frac{\epsilon_0}{2} \frac{1}{(4\pi\epsilon_0)^2} q^2 \left\{ \int_R^\infty \frac{1}{r^4} (r^2 4\pi dr) + \int_0^R \left( \frac{r}{R^3} \right)^2 (4\pi r^2 dr) \right\} \\ &= \frac{1}{4\pi\epsilon_0} \frac{q^2}{2} \left\{ \left( -\frac{1}{r} \right) \Big|_R^\infty + \frac{1}{R^6} \left( \frac{r^5}{5} \right) \Big|_0^R \right\} = \frac{1}{4\pi\epsilon_0} \frac{q^2}{2} \left( \frac{1}{R} + \frac{1}{5R} \right) = \frac{1}{4\pi\epsilon_0} \frac{3}{5} \frac{q^2}{R}. \checkmark \end{aligned}$$

(c)  $W = \frac{\epsilon_0}{2} \left\{ \oint_S \mathbf{V} \mathbf{E} \cdot d\mathbf{a} + \int_V E^2 d\tau \right\}$ , where  $V$  is large enough to enclose all the charge, but otherwise arbitrary. Let's use a sphere of radius  $a > R$ . Here  $V = \frac{1}{4\pi\epsilon_0} \frac{q}{r}$ .

$$\begin{aligned} W &= \frac{\epsilon_0}{2} \left\{ \int_{r=a} \left( \frac{1}{4\pi\epsilon_0} \frac{q}{r} \right) \left( \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \right) r^2 \sin\theta d\theta d\phi + \int_0^R E^2 d\tau + \int_R^a \left( \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \right)^2 (4\pi r^2 dr) \right\} \\ &= \frac{\epsilon_0}{2} \left\{ \frac{q^2}{(4\pi\epsilon_0)^2} \frac{1}{a} 4\pi + \frac{q^2}{(4\pi\epsilon_0)^2} \frac{4\pi}{5R} + \frac{1}{(4\pi\epsilon_0)^2} 4\pi q^2 \left( -\frac{1}{r} \right) \Big|_R^a \right\} \\ &= \frac{1}{4\pi\epsilon_0} \frac{q^2}{2} \left\{ \frac{1}{a} + \frac{1}{5R} - \frac{1}{a} + \frac{1}{R} \right\} = \frac{1}{4\pi\epsilon_0} \frac{3}{5} \frac{q^2}{R}. \checkmark \end{aligned}$$

As  $a \rightarrow \infty$ , the contribution from the surface integral  $\left( \frac{1}{4\pi\epsilon_0} \frac{q^2}{2a} \right)$  goes to zero, while the volume integral  $\left( \frac{1}{4\pi\epsilon_0} \frac{q^2}{2a} \left( \frac{6a}{5R} - 1 \right) \right)$  picks up the slack.

**Problem 2.35**


$$dW = d\bar{q} V = d\bar{q} \left( \frac{1}{4\pi\epsilon_0} \right) \frac{\bar{q}}{r}, \quad (\bar{q} = \text{charge on sphere of radius } r).$$

$$\bar{q} = \frac{4}{3} \pi r^3 \rho = q \frac{r^3}{R^3} \quad (q = \text{total charge on sphere}).$$

$$d\bar{q} = 4\pi r^2 dr \rho = \frac{4\pi r^2}{\frac{4}{3}\pi R^3} q dr = \frac{3q}{R^3} r^2 dr.$$

$$dW = \frac{1}{4\pi\epsilon_0} \left( \frac{qr^3}{R^3} \right) \frac{1}{r} \left( \frac{3q}{R^3} r^2 dr \right) = \frac{1}{4\pi\epsilon_0} \frac{3q^2}{R^6} r^4 dr$$

$$W = \frac{1}{4\pi\epsilon_0} \frac{3q^2}{R^6} \int_0^R r^4 dr = \frac{1}{4\pi\epsilon_0} \frac{3q^2}{R^6} \frac{R^5}{5} = \frac{1}{4\pi\epsilon_0} \left( \frac{3}{5} \frac{q^2}{R} \right). \checkmark$$




**Problem 2.36**

(a)  $W = \frac{\epsilon_0}{2} \int E^2 d\tau$ .  $\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} (a < r < b)$ , zero elsewhere.

$$W = \frac{\epsilon_0}{2} \left( \frac{q}{4\pi\epsilon_0} \right)^2 \int_a^b \left( \frac{1}{r^2} \right)^2 4\pi r^2 dr = \frac{q^2}{8\pi\epsilon_0} \int_a^b \frac{1}{r^2} = \boxed{\frac{q^2}{8\pi\epsilon_0} \left( \frac{1}{a} - \frac{1}{b} \right)}.$$

(b)  $W_1 = \frac{1}{8\pi\epsilon_0} \frac{q^2}{a}$ ,  $W_2 = \frac{1}{8\pi\epsilon_0} \frac{q^2}{b}$ ,  $\mathbf{E}_1 = \frac{1}{4\pi\epsilon_0} \frac{q}{r^2} \hat{\mathbf{r}} (r > a)$ ,  $\mathbf{E}_2 = \frac{1}{4\pi\epsilon_0} \frac{-q}{r^2} \hat{\mathbf{r}} (r > b)$ . So  $\mathbf{E}_1 \cdot \mathbf{E}_2 = \left( \frac{1}{4\pi\epsilon_0} \right)^2 \frac{-q^2}{r^4}$ , ( $r > b$ ), and hence  $\int \mathbf{E}_1 \cdot \mathbf{E}_2 d\tau = - \left( \frac{1}{4\pi\epsilon_0} \right)^2 q^2 \int_b^\infty \frac{1}{r^4} 4\pi r^2 dr = - \frac{q^2}{4\pi\epsilon_0 b}$ .  
 $W_{\text{tot}} = W_1 + W_2 + \epsilon_0 \int \mathbf{E}_1 \cdot \mathbf{E}_2 d\tau = \frac{1}{8\pi\epsilon_0} q^2 \left( \frac{1}{a} + \frac{1}{b} - \frac{2}{b} \right) = \frac{q^2}{8\pi\epsilon_0} \left( \frac{1}{a} - \frac{1}{b} \right) \checkmark$

**Problem 2.37**

$$\mathbf{E}_1 = \frac{1}{4\pi\epsilon_0} \frac{q_1}{r^2} \hat{\mathbf{r}}; \quad \mathbf{E}_2 = \frac{1}{4\pi\epsilon_0} \frac{q_2}{z^2} \hat{\mathbf{z}}; \quad W_i = \epsilon_0 \frac{q_1 q_2}{(4\pi\epsilon_0)^2} \int \frac{1}{r^2 z^2} \cos \beta r^2 \sin \theta dr d\theta d\phi,$$

where (from the figure)

$$z = \sqrt{r^2 + a^2 - 2ra \cos \theta}, \quad \cos \beta = \frac{(r - a \cos \theta)}{z}.$$

Therefore

$$W_i = \frac{q_1 q_2}{(4\pi)^2 \epsilon_0} 2\pi \int \frac{(r - a \cos \theta)}{z^3} \sin \theta dr d\theta.$$

It's simplest to do the  $r$  integral first, changing variables to  $z$  :

$$2z dz = (2r - 2a \cos \theta) dr \Rightarrow (r - a \cos \theta) dr = z dz.$$

As  $r : 0 \rightarrow \infty$ ,  $z : a \rightarrow \infty$ , so

$$W_i = \frac{q_1 q_2}{8\pi\epsilon_0} \int_0^\pi \left( \int_a^\infty \frac{1}{z^2} dz \right) \sin \theta d\theta.$$

The  $z$  integral is  $1/a$ , so

$$W_i = \frac{q_1 q_2}{8\pi\epsilon_0 a} \int_0^\pi \sin \theta d\theta = \boxed{\frac{q_1 q_2}{4\pi\epsilon_0 a}}.$$

Of course, this is precisely the interaction energy of two point charges.

**Problem 2.38**

(a)  $\sigma_R = \frac{q}{4\pi R^2}$ ;  $\sigma_a = \frac{-q}{4\pi a^2}$ ;  $\sigma_b = \frac{q}{4\pi b^2}$ .

$$(b) V(0) = -\int_{\infty}^0 \mathbf{E} \cdot d\mathbf{l} = -\int_{\infty}^b \left(\frac{1}{4\pi\epsilon_0} \frac{q}{r^2}\right) dr - \int_b^a (0) dr - \int_a^R \left(\frac{1}{4\pi\epsilon_0} \frac{q}{r^2}\right) dr - \int_R^0 (0) dr = \boxed{\frac{1}{4\pi\epsilon_0} \left(\frac{q}{b} + \frac{q}{R} - \frac{q}{a}\right)}.$$

$$(c) \boxed{\sigma_b \rightarrow 0} \text{ (the charge "drains off")}; V(0) = -\int_{\infty}^a (0) dr - \int_a^R \left(\frac{1}{4\pi\epsilon_0} \frac{q}{r^2}\right) dr - \int_R^0 (0) dr = \boxed{\frac{1}{4\pi\epsilon_0} \left(\frac{q}{R} - \frac{q}{a}\right)}.$$

**Problem 2.39**

$$(a) \boxed{\sigma_a = -\frac{q_a}{4\pi a^2}}; \boxed{\sigma_b = -\frac{q_b}{4\pi b^2}}; \boxed{\sigma_R = \frac{q_a + q_b}{4\pi R^2}}.$$

$$(b) \boxed{\mathbf{E}_{\text{out}} = \frac{1}{4\pi\epsilon_0} \frac{q_a + q_b}{r^2} \hat{\mathbf{r}}}, \text{ where } \mathbf{r} = \text{vector from center of large sphere}.$$

$$(c) \boxed{\mathbf{E}_a = \frac{1}{4\pi\epsilon_0} \frac{q_a}{r_a^2} \hat{\mathbf{r}}_a, \quad \mathbf{E}_b = \frac{1}{4\pi\epsilon_0} \frac{q_b}{r_b^2} \hat{\mathbf{r}}_b}, \text{ where } \mathbf{r}_a \text{ (} \mathbf{r}_b \text{) is the vector from center of cavity } a \text{ (} b \text{)}.$$

$$(d) \boxed{\text{Zero}}.$$

$$(e) \sigma_R \text{ changes (but not } \sigma_a \text{ or } \sigma_b \text{); } \mathbf{E}_{\text{outside}} \text{ changes (but not } \mathbf{E}_a \text{ or } \mathbf{E}_b \text{); force on } q_a \text{ and } q_b \text{ still zero.}$$

**Problem 2.40**

(a)  $\boxed{\text{No.}}$  For example, if it is very close to the wall, it will induce charge of the opposite sign on the wall, and it will be attracted.

(b)  $\boxed{\text{No.}}$  Typically it will be attractive, but see footnote 12 for an extraordinary counterexample.

**Problem 2.41**

Between the plates,  $E = 0$ ; outside the plates  $E = \sigma/\epsilon_0 = Q/\epsilon_0 A$ . So

$$P = \frac{\epsilon_0}{2} E^2 = \frac{\epsilon_0}{2} \frac{Q^2}{\epsilon_0^2 A^2} = \boxed{\frac{Q^2}{2\epsilon_0 A^2}}.$$


**Problem 2.42**

Inside,  $\mathbf{E} = \mathbf{0}$ ; outside,  $\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2} \hat{\mathbf{r}}$ ; so

$$\mathbf{E}_{\text{ave}} = \frac{1}{2} \frac{1}{4\pi\epsilon_0} \frac{Q}{R^2} \hat{\mathbf{r}}; \quad f_z = \sigma(E_{\text{ave}})_z; \quad \sigma = \frac{Q}{4\pi R^2}.$$

$$F_z = \int f_z da = \int \left(\frac{Q}{4\pi R^2}\right) \frac{1}{2} \left(\frac{1}{4\pi\epsilon_0} \frac{Q}{R^2}\right) \cos\theta R^2 \sin\theta d\theta d\phi$$

$$= \frac{1}{2\epsilon_0} \left(\frac{Q}{4\pi R}\right)^2 2\pi \int_0^{\pi/2} \sin\theta \cos\theta d\theta = \frac{1}{\pi\epsilon_0} \left(\frac{Q}{4R}\right)^2 \left(\frac{1}{2} \sin^2\theta\right) \Big|_0^{\pi/2} = \frac{1}{2\pi\epsilon_0} \left(\frac{Q}{4R}\right)^2 = \boxed{\frac{Q^2}{32\pi R^2 \epsilon_0}}.$$

**Problem 2.43**

Say the charge on the inner cylinder is  $Q$ , for a length  $L$ . The field is given by Gauss's law:

$\int \mathbf{E} \cdot d\mathbf{a} = E \cdot 2\pi s \cdot L = \frac{1}{\epsilon_0} Q_{\text{enc}} = \frac{1}{\epsilon_0} Q \Rightarrow \mathbf{E} = \frac{Q}{2\pi\epsilon_0 L} \frac{1}{s} \hat{\mathbf{s}}$ . Potential difference between the cylinders is

$$V(b) - V(a) = -\int_a^b \mathbf{E} \cdot d\mathbf{l} = -\frac{Q}{2\pi\epsilon_0 L} \int_a^b \frac{1}{s} ds = -\frac{Q}{2\pi\epsilon_0 L} \ln\left(\frac{b}{a}\right).$$

As set up here,  $a$  is at the higher potential, so  $V = V(a) - V(b) = \frac{Q}{2\pi\epsilon_0 L} \ln\left(\frac{b}{a}\right)$ .

$$C = \frac{Q}{V} = \frac{2\pi\epsilon_0 L}{\ln\left(\frac{b}{a}\right)}, \text{ so capacitance per unit length is } \boxed{\frac{2\pi\epsilon_0}{\ln\left(\frac{b}{a}\right)}}.$$

**Problem 2.44**


$$(a) W = (\text{force}) \times (\text{distance}) = (\text{pressure}) \times (\text{area}) \times (\text{distance}) = \boxed{\frac{\epsilon_0}{2} E^2 A \epsilon}.$$

(b)  $W = (\text{energy per unit volume}) \times (\text{decrease in volume}) = \left(\epsilon_0 \frac{E^2}{2}\right) (A\epsilon)$ . Same as (a), confirming that the energy lost is equal to the work done.

**Problem 2.45**

From Prob. 2.4, the field at height  $z$  above the center of a square loop (side  $a$ ) is

$$\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{4\lambda a z}{\left(z^2 + \frac{a^2}{4}\right) \sqrt{z^2 + \frac{a^2}{2}}} \hat{\mathbf{z}}.$$



Here  $\lambda \rightarrow \sigma \frac{da}{2}$  (see figure), and we integrate over  $a$  from 0 to  $\bar{a}$ :

$$\begin{aligned} E &= \frac{1}{4\pi\epsilon_0} 2\sigma z \int_0^{\bar{a}} \frac{a da}{\left(z^2 + \frac{a^2}{4}\right) \sqrt{z^2 + \frac{a^2}{2}}} . \text{ Let } u = \frac{a^2}{4}, \text{ so } a da = 2 du. \\ &= \frac{1}{4\pi\epsilon_0} 4\sigma z \int_0^{\bar{a}^2/4} \frac{du}{(u + z^2) \sqrt{2u + z^2}} = \frac{\sigma z}{\pi\epsilon_0} \left[ \frac{2}{z} \tan^{-1} \left( \frac{\sqrt{2u + z^2}}{z} \right) \right]_0^{\bar{a}^2/4} \\ &= \frac{2\sigma}{\pi\epsilon_0} \left\{ \tan^{-1} \left( \frac{\sqrt{\frac{\bar{a}^2}{2} + z^2}}{z} \right) - \tan^{-1}(1) \right\}; \end{aligned}$$

$$\boxed{\mathbf{E} = \frac{2\sigma}{\pi\epsilon_0} \left[ \tan^{-1} \sqrt{1 + \frac{a^2}{2z^2}} - \frac{\pi}{4} \right] \hat{\mathbf{z}}} = \frac{\sigma}{\pi\epsilon_0} \tan^{-1} \left( \frac{a^2}{4z \sqrt{z^2 + (a^2/2)}} \right) \hat{\mathbf{z}}.$$

$$a \rightarrow \infty \text{ (infinite plane): } E = \frac{2\sigma}{\pi\epsilon_0} [\tan^{-1}(\infty) - \frac{\pi}{4}] = \frac{2\sigma}{\pi\epsilon_0} \left( \frac{\pi}{2} - \frac{\pi}{4} \right) = \frac{\sigma}{2\epsilon_0}. \checkmark$$

$$z \gg a \text{ (point charge): Let } f(x) = \tan^{-1} \sqrt{1+x} - \frac{\pi}{4}, \text{ and expand as a Taylor series:}$$

$$f(x) = f(0) + x f'(0) + \frac{1}{2} x^2 f''(0) + \dots$$

$$\text{Here } f(0) = \tan^{-1}(1) - \frac{\pi}{4} = \frac{\pi}{4} - \frac{\pi}{4} = 0; f'(x) = \frac{1}{1+(1+x)} \frac{1}{2} \frac{1}{\sqrt{1+x}} = \frac{1}{2(2+x)\sqrt{1+x}}, \text{ so } f'(0) = \frac{1}{4}, \text{ so}$$

$$f(x) = \frac{1}{4} x + ( ) x^2 + ( ) x^3 + \dots$$

$$\text{Thus (since } \frac{a^2}{2z^2} = x \ll 1), E \approx \frac{2\sigma}{\pi\epsilon_0} \left( \frac{1}{4} \frac{a^2}{2z^2} \right) = \frac{1}{4\pi\epsilon_0} \frac{\sigma a^2}{z^2} = \frac{1}{4\pi\epsilon_0} \frac{q}{z^2}. \checkmark$$

**Problem 2.46**

$$\begin{aligned}
\rho &= \epsilon_0 \nabla \cdot \mathbf{E} = \epsilon_0 \left\{ \frac{1}{r^2} \frac{\partial}{\partial r} \left( r^2 \frac{3k}{r} \right) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \theta} \left( \sin \theta \frac{2k \sin \theta \cos \theta \sin \phi}{r} \right) + \frac{1}{r \sin \theta} \frac{\partial}{\partial \phi} \left( \frac{k \sin \theta \cos \phi}{r} \right) \right\} \\
&= \epsilon_0 \left[ \frac{1}{r^2} 3k + \frac{1}{r \sin \theta} \frac{2k \sin \phi (2 \sin \theta \cos^2 \theta - \sin^3 \theta)}{r} + \frac{1}{r \sin \theta} \frac{(-k \sin \theta \sin \phi)}{r} \right] \\
&= \frac{k \epsilon_0}{r^2} [3 + 2 \sin \phi (2 \cos^2 \theta - \sin^2 \theta) - \sin \phi] = \frac{k \epsilon_0}{r^2} [3 + \sin \phi (4 \cos^2 \theta - 2 + 2 \cos^2 \theta - 1)] \\
&= \frac{3k \epsilon_0}{r^2} [1 + \sin \phi (2 \cos^2 \theta - 1)] = \boxed{\frac{3k \epsilon_0}{r^2} (1 + \sin \phi \cos 2\theta)}.
\end{aligned}$$

**Problem 2.47**

From Prob. 2.12, the field inside a uniformly charged sphere is:  $\mathbf{E} = \frac{1}{4\pi\epsilon_0} \frac{Q}{R^3} \mathbf{r}$ . So the force per unit volume is  $\mathbf{f} = \rho \mathbf{E} = \left( \frac{Q}{\frac{4}{3}\pi R^3} \right) \left( \frac{Q}{4\pi\epsilon_0 R^3} \right) \mathbf{r} = \frac{3}{\epsilon_0} \left( \frac{Q}{4\pi R^3} \right)^2 \mathbf{r}$ , and the force in the  $z$  direction on  $d\tau$  is:


$$dF_z = f_z d\tau = \frac{3}{\epsilon_0} \left( \frac{Q}{4\pi R^3} \right)^2 r \cos \theta (r^2 \sin \theta dr d\theta d\phi).$$

The total force on the “northern” hemisphere is:

$$\begin{aligned}
F_z &= \int f_z d\tau = \frac{3}{\epsilon_0} \left( \frac{Q}{4\pi R^3} \right)^2 \int_0^R r^3 dr \int_0^{\pi/2} \cos \theta \sin \theta d\theta \int_0^{2\pi} d\phi \\
&= \frac{3}{\epsilon_0} \left( \frac{Q}{4\pi R^3} \right)^2 \left( \frac{R^4}{4} \right) \left( \frac{\sin^2 \theta}{2} \Big|_0^{\pi/2} \right) (2\pi) = \boxed{\frac{3Q^2}{64\pi\epsilon_0 R^2}}.
\end{aligned}$$

**Problem 2.48**

$$\begin{aligned}
V_{\text{center}} &= \frac{1}{4\pi\epsilon_0} \int \frac{\sigma}{z} da = \frac{1}{4\pi\epsilon_0} \frac{\sigma}{R} \int da = \frac{1}{4\pi\epsilon_0} \frac{\sigma}{R} (2\pi R^2) = \frac{\sigma R}{2\epsilon_0} \\
V_{\text{pole}} &= \frac{1}{4\pi\epsilon_0} \int \frac{\sigma}{z} da, \text{ with } \begin{cases} da = 2\pi R^2 \sin \theta d\theta, \\ z^2 = R^2 + R^2 - 2R^2 \cos \theta = 2R^2(1 - \cos \theta). \end{cases} \\
&= \frac{1}{4\pi\epsilon_0} \frac{\sigma(2\pi R^2)}{R\sqrt{2}} \int_0^{\pi/2} \frac{\sin \theta d\theta}{\sqrt{1 - \cos \theta}} = \frac{\sigma R}{2\sqrt{2}\epsilon_0} (2\sqrt{1 - \cos \theta}) \Big|_0^{\pi/2} \\
&= \frac{\sigma R}{\sqrt{2}\epsilon_0} (1 - 0) = \frac{\sigma R}{\sqrt{2}\epsilon_0}. \quad \therefore V_{\text{pole}} - V_{\text{center}} = \boxed{\frac{\sigma R}{2\epsilon_0} (\sqrt{2} - 1)}.
\end{aligned}$$

**Problem 2.49**

First let's determine the electric field inside and outside the sphere, using Gauss's law:

$$\epsilon_0 \oint \mathbf{E} \cdot d\mathbf{a} = \epsilon_0 4\pi r^2 E = Q_{\text{enc}} = \int \rho d\tau = \int (k\bar{r}) \bar{r}^2 \sin \theta d\bar{r} d\theta d\phi = 4\pi k \int_0^r \bar{r}^3 d\bar{r} = \begin{cases} \pi k r^4 & (r < R), \\ \pi k R^4 & (r > R). \end{cases}$$

So  $\mathbf{E} = \frac{k}{4\epsilon_0} r^2 \hat{\mathbf{r}}$  ( $r < R$ );  $\mathbf{E} = \frac{kR^4}{4\epsilon_0 r^2} \hat{\mathbf{r}}$  ( $r > R$ ).

Method I:

$$\begin{aligned}
 W &= \frac{\epsilon_0}{2} \int E^2 d\tau \quad (\text{Eq. 2.45}) = \frac{\epsilon_0}{2} \int_0^R \left( \frac{kr^2}{4\epsilon_0} \right)^2 4\pi r^2 dr + \frac{\epsilon_0}{2} \int_R^\infty \left( \frac{kR^4}{4\epsilon_0 r^2} \right)^2 4\pi r^2 dr \\
 &= 4\pi \frac{\epsilon_0}{2} \left( \frac{k}{4\epsilon_0} \right)^2 \left\{ \int_0^R r^6 dr + R^8 \int_R^\infty \frac{1}{r^2} dr \right\} = \frac{\pi k^2}{8\epsilon_0} \left\{ \frac{R^7}{7} + R^8 \left( -\frac{1}{r} \right) \Big|_R^\infty \right\} = \frac{\pi k^2}{8\epsilon_0} \left( \frac{R^7}{7} + R^7 \right) \\
 &= \boxed{\frac{\pi k^2 R^7}{7\epsilon_0}}.
 \end{aligned}$$

Method II:

$$\begin{aligned}
 W &= \frac{1}{2} \int \rho V d\tau \quad (\text{Eq. 2.43}). \\
 \text{For } r < R, \quad V(r) &= - \int_\infty^r \mathbf{E} \cdot d\mathbf{l} = - \int_\infty^R \left( \frac{kR^4}{4\epsilon_0 r^2} \right) dr - \int_R^r \left( \frac{kr^2}{4\epsilon_0} \right) dr = -\frac{k}{4\epsilon_0} \left\{ R^4 \left( -\frac{1}{r} \right) \Big|_\infty^R + \frac{r^3}{3} \Big|_R^r \right\} \\
 &= -\frac{k}{4\epsilon_0} \left( -R^3 + \frac{r^3}{3} - \frac{R^3}{3} \right) = \frac{k}{3\epsilon_0} \left( R^3 - \frac{r^3}{4} \right). \\
 \therefore W &= \frac{1}{2} \int_0^R (kr) \left[ \frac{k}{3\epsilon_0} \left( R^3 - \frac{r^3}{4} \right) \right] 4\pi r^2 dr = \frac{2\pi k^2}{3\epsilon_0} \int_0^R \left( R^3 r^3 - \frac{1}{4} r^6 \right) dr \\
 &= \frac{2\pi k^2}{3\epsilon_0} \left\{ R^3 \frac{R^4}{4} - \frac{1}{4} \frac{R^7}{7} \right\} = \frac{\pi k^2 R^7}{2 \cdot 3\epsilon_0} \left( \frac{6}{7} \right) = \frac{\pi k^2 R^7}{7\epsilon_0}. \quad \checkmark
 \end{aligned}$$

### Problem 2.50

$$\mathbf{E} = -\nabla V = -A \frac{\partial}{\partial r} \left( \frac{e^{-\lambda r}}{r} \right) \hat{\mathbf{r}} = -A \left\{ \frac{r(-\lambda)e^{-\lambda r} - e^{-\lambda r}}{r^2} \right\} \hat{\mathbf{r}} = \boxed{Ae^{-\lambda r} (1 + \lambda r) \frac{\hat{\mathbf{r}}}{r^2}}.$$

$\rho = \epsilon_0 \nabla \cdot \mathbf{E} = \epsilon_0 A \left\{ e^{-\lambda r} (1 + \lambda r) \nabla \cdot \left( \frac{\hat{\mathbf{r}}}{r^2} \right) + \frac{\hat{\mathbf{r}}}{r^2} \cdot \nabla (e^{-\lambda r} (1 + \lambda r)) \right\}$ . But  $\nabla \cdot \left( \frac{\hat{\mathbf{r}}}{r^2} \right) = 4\pi \delta^3(\mathbf{r})$  (Eq. 1.99), and  $e^{-\lambda r} (1 + \lambda r) \delta^3(\mathbf{r}) = \delta^3(\mathbf{r})$  (Eq. 1.88). Meanwhile,

$$\nabla (e^{-\lambda r} (1 + \lambda r)) = \hat{\mathbf{r}} \frac{\partial}{\partial r} (e^{-\lambda r} (1 + \lambda r)) = \hat{\mathbf{r}} \{-\lambda e^{-\lambda r} (1 + \lambda r) + e^{-\lambda r} \lambda\} = \hat{\mathbf{r}} (-\lambda^2 r e^{-\lambda r}).$$

$$\text{So } \frac{\hat{\mathbf{r}}}{r^2} \cdot \nabla (e^{-\lambda r} (1 + \lambda r)) = -\frac{\lambda^2}{r} e^{-\lambda r}, \text{ and } \boxed{\rho = \epsilon_0 A \left[ 4\pi \delta^3(\mathbf{r}) - \frac{\lambda^2}{r} e^{-\lambda r} \right]}.$$

$$Q = \int \rho d\tau = \epsilon_0 A \left\{ 4\pi \int \delta^3(\mathbf{r}) d\tau - \lambda^2 \int \frac{e^{-\lambda r}}{r} 4\pi r^2 dr \right\} = \epsilon_0 A \left( 4\pi - \lambda^2 4\pi \int_0^\infty r e^{-\lambda r} dr \right).$$

$$\text{But } \int_0^\infty r e^{-\lambda r} dr = \frac{1}{\lambda^2}, \text{ so } Q = 4\pi \epsilon_0 A \left( 1 - \frac{\lambda^2}{\lambda^2} \right) = \boxed{\text{zero}}.$$

### Problem 2.51

$$V = \frac{1}{4\pi \epsilon_0} \int \frac{\sigma}{r} da = \frac{\sigma}{4\pi \epsilon_0} \int_0^R \int_0^{2\pi} \frac{1}{\sqrt{R^2 + s^2 - 2Rs \cos \phi}} s ds d\phi.$$

Let  $u \equiv s/R$ . Then

$$V = \frac{2\sigma R}{4\pi \epsilon_0} \int_0^1 \left( \int_0^\pi \frac{u}{\sqrt{1 + u^2 - 2u \cos \phi}} d\phi \right) du.$$



The (double) integral is a pure number; Mathematica says it is 2. So

$$V = \frac{\sigma R}{\pi \epsilon_0}.$$


### Problem 2.52

- (a) Potential of  $+\lambda$  is  $V_+ = -\frac{\lambda}{2\pi\epsilon_0} \ln\left(\frac{s_+}{a}\right)$ , where  $s_+$  is distance from  $\lambda_+$  (Prob. 2.22).  
Potential of  $-\lambda$  is  $V_- = +\frac{\lambda}{2\pi\epsilon_0} \ln\left(\frac{s_-}{a}\right)$ , where  $s_-$  is distance from  $\lambda_-$ .

$$\therefore \text{Total } V = \frac{\lambda}{2\pi\epsilon_0} \ln\left(\frac{s_-}{s_+}\right).$$

Now  $s_+ = \sqrt{(y-a)^2 + z^2}$ , and  $s_- = \sqrt{(y+a)^2 + z^2}$ , so

$$V(x, y, z) = \frac{\lambda}{2\pi\epsilon_0} \ln\left(\frac{\sqrt{(y+a)^2 + z^2}}{\sqrt{(y-a)^2 + z^2}}\right) = \frac{\lambda}{4\pi\epsilon_0} \ln\left[\frac{(y+a)^2 + z^2}{(y-a)^2 + z^2}\right].$$



- (b) Equipotentials are given by  $\frac{(y+a)^2 + z^2}{(y-a)^2 + z^2} = e^{(4\pi\epsilon_0 V_0/\lambda)} = k = \text{constant}$ . That is:  
 $y^2 + 2ay + a^2 + z^2 = k(y^2 - 2ay + a^2 + z^2) \Rightarrow y^2(k-1) + z^2(k-1) + a^2(k-1) - 2ay(k+1) = 0$ , or  
 $y^2 + z^2 + a^2 - 2ay\left(\frac{k+1}{k-1}\right) = 0$ . The equation for a *circle*, with center at  $(y_0, 0)$  and radius  $R$ , is  
 $(y - y_0)^2 + z^2 = R^2$ , or  $y^2 + z^2 + (y_0^2 - R^2) - 2yy_0 = 0$ .


Evidently the equipotentials *are* circles, with  $y_0 = a\left(\frac{k+1}{k-1}\right)$  and

$$a^2 = y_0^2 - R^2 \Rightarrow R^2 = y_0^2 - a^2 = a^2\left(\frac{k+1}{k-1}\right)^2 - a^2 = a^2 \frac{(k^2 + 2k + 1 - k^2 - 2k - 1)}{(k-1)^2} = a^2 \frac{4k}{(k-1)^2}, \text{ or}$$

$$R = \frac{2a\sqrt{k}}{|k-1|}; \text{ or, in terms of } V_0:$$

$$y_0 = a \frac{e^{4\pi\epsilon_0 V_0/\lambda} + 1}{e^{4\pi\epsilon_0 V_0/\lambda} - 1} = a \frac{e^{2\pi\epsilon_0 V_0/\lambda} + e^{-2\pi\epsilon_0 V_0/\lambda}}{e^{2\pi\epsilon_0 V_0/\lambda} - e^{-2\pi\epsilon_0 V_0/\lambda}} = a \coth\left(\frac{2\pi\epsilon_0 V_0}{\lambda}\right).$$

$$R = 2a \frac{e^{2\pi\epsilon_0 V_0/\lambda}}{e^{4\pi\epsilon_0 V_0/\lambda} - 1} = a \frac{2}{(e^{2\pi\epsilon_0 V_0/\lambda} - e^{-2\pi\epsilon_0 V_0/\lambda})} = \frac{a}{\sinh\left(\frac{2\pi\epsilon_0 V_0}{\lambda}\right)} = a \operatorname{csch}\left(\frac{2\pi\epsilon_0 V_0}{\lambda}\right).$$



**Problem 2.53**

(a)  $\nabla^2 V = -\frac{\rho}{\epsilon_0}$  (Eq. 2.24), so  $\boxed{\frac{d^2 V}{dx^2} = -\frac{1}{\epsilon_0} \rho.}$

(b)  $qV = \frac{1}{2}mv^2 \rightarrow \boxed{v = \sqrt{\frac{2qV}{m}}.}$

(c)  $dq = A\rho dx$ ;  $\frac{dq}{dt} = a\rho \frac{dx}{dt} = \boxed{A\rho v = I}$  (constant). (Note:  $\rho$ , hence also  $I$ , is negative.)

(d)  $\frac{d^2 V}{dx^2} = -\frac{1}{\epsilon_0} \rho = -\frac{1}{\epsilon_0} \frac{I}{Av} = -\frac{I}{\epsilon_0 A} \sqrt{\frac{m}{2qV}} \Rightarrow \boxed{\frac{d^2 V}{dx^2} = \beta V^{-1/2}}$ , where  $\beta = -\frac{I}{\epsilon_0 A} \sqrt{\frac{m}{2q}}$ .  
(Note:  $I$  is negative, so  $\beta$  is positive;  $q$  is positive.)

(e) Multiply by  $V' = \frac{dV}{dx}$ :

$$V' \frac{dV'}{dx} = \beta V^{-1/2} \frac{dV}{dx} \Rightarrow \int V' dV' = \beta \int V^{-1/2} dV \Rightarrow \frac{1}{2} V'^2 = 2\beta V^{1/2} + \text{constant}.$$

But  $V(0) = V'(0) = 0$  (cathode is at potential zero, and field at cathode is zero), so the constant is zero, and

$$V'^2 = 4\beta V^{1/2} \Rightarrow \frac{dV}{dx} = 2\sqrt{\beta} V^{1/4} \Rightarrow V^{-1/4} dV = 2\sqrt{\beta} dx;$$

$$\int V^{-1/4} dV = 2\sqrt{\beta} \int dx \Rightarrow \frac{4}{3} V^{3/4} = 2\sqrt{\beta} x + \text{constant}.$$

But  $V(0) = 0$ , so this constant is also zero.

$$V^{3/4} = \frac{3}{2} \sqrt{\beta} x, \text{ so } V(x) = \left(\frac{3}{2} \sqrt{\beta}\right)^{4/3} x^{4/3}, \text{ or } V(x) = \left(\frac{9}{4} \beta\right)^{2/3} x^{4/3} = \left(\frac{81 I^2 m}{32 \epsilon_0^2 A^2 q}\right)^{1/3} x^{4/3}.$$

In terms of  $V_0$  (instead of  $I$ ):  $\boxed{V(x) = V_0 \left(\frac{x}{d}\right)^{4/3}}$  (see graph).


Without space-charge,  $V$  would increase linearly:  $V(x) = V_0 \left(\frac{x}{d}\right)$ .

$$\rho = -\epsilon_0 \frac{d^2 V}{dx^2} = -\epsilon_0 V_0 \frac{1}{d^{4/3}} \frac{4}{3} \cdot \frac{1}{3} x^{-2/3} = \boxed{-\frac{4\epsilon_0 V_0}{9(d^2 x)^{2/3}}}.$$

$$v = \sqrt{\frac{2q}{m}} \sqrt{V} = \boxed{\sqrt{2qV_0/m} \left(\frac{x}{d}\right)^{2/3}}.$$

(f)  $V(d) = V_0 = \left(\frac{81 I^2 m}{32 \epsilon_0^2 A^2 q}\right)^{1/3} d^{4/3} \Rightarrow V_0^3 = \frac{81 m d^4}{32 \epsilon_0^2 A^2 q} I^2$ ;  $I^2 = \frac{32 \epsilon_0^2 A^2 q}{81 m d^4} V_0^3$ ;

$$I = \frac{4\sqrt{2} \epsilon_0 A \sqrt{q}}{9\sqrt{m} d^2} V_0^{3/2} = K V_0^{3/2}, \text{ where } \boxed{K = \frac{4\epsilon_0 A}{9d^2} \sqrt{\frac{2q}{m}}}.$$

**Problem 2.54**

(a)  $\boxed{\mathbf{E} = \frac{1}{4\pi\epsilon_0} \int \frac{\rho \hat{\mathbf{z}}}{z^2} \left(1 + \frac{z}{\lambda}\right) e^{-z/\lambda} d\tau.}$

(b) Yes. The field of a point charge at the origin is radial and symmetric, so  $\nabla \times \mathbf{E} = \mathbf{0}$ , and hence this is also true (by superposition) for any *collection* of charges.


$$\begin{aligned} \text{(c)} \quad V &= - \int_{\infty}^r \mathbf{E} \cdot d\mathbf{l} = - \frac{1}{4\pi\epsilon_0} q \int_{\infty}^r \frac{1}{r^2} \left(1 + \frac{r}{\lambda}\right) e^{-r/\lambda} dr \\ &= \frac{1}{4\pi\epsilon_0} q \int_r^{\infty} \frac{1}{r^2} \left(1 + \frac{r}{\lambda}\right) e^{-r/\lambda} dr = \frac{q}{4\pi\epsilon_0} \left\{ \int_r^{\infty} \frac{1}{r^2} e^{-r/\lambda} dr + \frac{1}{\lambda} \int_r^{\infty} \frac{1}{r} e^{-r/\lambda} dr \right\}. \end{aligned}$$

Now  $\int \frac{1}{r^2} e^{-r/\lambda} dr = -\frac{e^{-r/\lambda}}{r} - \frac{1}{\lambda} \int \frac{e^{-r/\lambda}}{r} dr \leftarrow$  exactly right to kill the last term. Therefore

$$V(r) = \frac{q}{4\pi\epsilon_0} \left\{ -\frac{e^{-r/\lambda}}{r} \Big|_r^{\infty} \right\} = \boxed{\frac{q}{4\pi\epsilon_0} \frac{e^{-r/\lambda}}{r}}.$$

$$\begin{aligned} \text{(d)} \quad \oint_S \mathbf{E} \cdot d\mathbf{a} &= \frac{1}{4\pi\epsilon_0} q \frac{1}{R^2} \left(1 + \frac{R}{\lambda}\right) e^{-R/\lambda} 4\pi R^2 = \frac{q}{\epsilon_0} \left(1 + \frac{R}{\lambda}\right) e^{-R/\lambda}. \\ \int_V V d\tau &= \frac{q}{4\pi\epsilon_0} \int_0^R \frac{e^{-r/\lambda}}{r} r^2 4\pi dr = \frac{q}{\epsilon_0} \int_0^R r e^{-r/\lambda} dr = \frac{q}{\epsilon_0} \left[ \frac{e^{-r/\lambda}}{(1/\lambda)^2} \left(-\frac{r}{\lambda} - 1\right) \right]_0^R \\ &= \lambda^2 \frac{q}{\epsilon_0} \left\{ -e^{-R/\lambda} \left(1 + \frac{R}{\lambda}\right) + 1 \right\}. \\ \therefore \oint_S \mathbf{E} \cdot d\mathbf{a} + \frac{1}{\lambda^2} \int_V V d\tau &= \frac{q}{\epsilon_0} \left\{ \left(1 + \frac{R}{\lambda}\right) e^{-R/\lambda} - \left(1 + \frac{R}{\lambda}\right) e^{-R/\lambda} + 1 \right\} = \frac{q}{\epsilon_0}. \quad \text{qed} \end{aligned}$$

(e) Does the result in (d) hold for a *nonspherical* surface? Suppose we make a “dent” in the sphere—pushing a patch (area  $R^2 \sin \theta d\theta d\phi$ ) from radius  $R$  out to radius  $S$  (area  $S^2 \sin \theta d\theta d\phi$ ).




$$\begin{aligned} \Delta \oint \mathbf{E} \cdot d\mathbf{a} &= \frac{q}{4\pi\epsilon_0} \left\{ \frac{1}{S^2} \left(1 + \frac{S}{\lambda}\right) e^{-S/\lambda} (S^2 \sin \theta d\theta d\phi) - \frac{1}{R^2} \left(1 + \frac{R}{\lambda}\right) e^{-R/\lambda} (R^2 \sin \theta d\theta d\phi) \right\} \\ &= \frac{q}{4\pi\epsilon_0} \left[ \left(1 + \frac{S}{\lambda}\right) e^{-S/\lambda} - \left(1 + \frac{R}{\lambda}\right) e^{-R/\lambda} \right] \sin \theta d\theta d\phi. \end{aligned}$$

$$\begin{aligned} \Delta \frac{1}{\lambda^2} \int V d\tau &= \frac{1}{\lambda^2} \frac{q}{4\pi\epsilon_0} \int \frac{e^{-r/\lambda}}{r} r^2 \sin \theta dr d\theta d\phi = \frac{1}{\lambda^2} \frac{q}{4\pi\epsilon_0} \sin \theta d\theta d\phi \int_R^S r e^{-r/\lambda} dr \\ &= -\frac{q}{4\pi\epsilon_0} \sin \theta d\theta d\phi \left( e^{-r/\lambda} \left(1 + \frac{r}{\lambda}\right) \right) \Big|_R^S \\ &= -\frac{q}{4\pi\epsilon_0} \left[ \left(1 + \frac{S}{\lambda}\right) e^{-S/\lambda} - \left(1 + \frac{R}{\lambda}\right) e^{-R/\lambda} \right] \sin \theta d\theta d\phi. \end{aligned}$$

So the change in  $\frac{1}{\lambda^2} \int V d\tau$  exactly compensates for the change in  $\oint \mathbf{E} \cdot d\mathbf{a}$ , and we get  $\frac{1}{\epsilon_0} q$  for the total using the dented sphere, just as we did with the perfect sphere. Any closed surface can be built up by successive distortions of the sphere, so the result holds for all shapes. By superposition, if there are many charges inside, the total is  $\frac{1}{\epsilon_0} Q_{\text{enc}}$ . Charges *outside* do not contribute (in the argument above we found that  $\oint \mathbf{E} \cdot d\mathbf{a} = 0$  for this volume  $\oint \mathbf{E} \cdot d\mathbf{a} + \frac{1}{\lambda^2} \int V d\tau = 0$ —and, again, the sum is not changed by distortions of the surface, as long as  $q$  remains outside). So the new “Gauss’s Law” holds for *any* charge configuration.

(f) In differential form, “Gauss’s law” reads:  $\nabla \cdot \mathbf{E} + \frac{1}{\lambda^2} V = \frac{1}{\epsilon_0} \rho$ , or, putting it all in terms of  $\mathbf{E}$ :

$\nabla \cdot \mathbf{E} - \frac{1}{\lambda^2} \int \mathbf{E} \cdot d\mathbf{l} = \frac{1}{\epsilon_0} \rho$ . Since  $\mathbf{E} = -\nabla V$ , this also yields “Poisson’s equation”:  $-\nabla^2 V + \frac{1}{\lambda^2} V = \frac{1}{\epsilon_0} \rho$ .



(g) Refer to “Gauss’s law” in differential form (f). Since  $\mathbf{E}$  is zero, inside a conductor (otherwise charge would move, and in such a direction as to cancel the field),  $V$  is constant (inside), and hence  $\rho$  is uniform, throughout the volume. Any “extra” charge must reside on the surface. (The fraction at the surface depends on  $\lambda$ , and on the shape of the conductor.)

### Problem 2.55

$$\rho = \epsilon_0 \nabla \cdot \mathbf{E} = \epsilon_0 \frac{\partial}{\partial x}(ax) = \boxed{\epsilon_0 a} \text{ (constant everywhere).}$$

The same charge density would be compatible (as far as Gauss’s law is concerned) with  $\mathbf{E} = ay\hat{\mathbf{y}}$ , for instance, or  $\mathbf{E} = (\frac{a}{3})\mathbf{r}$ , etc. The point is that Gauss’s law (and  $\nabla \times \mathbf{E} = \mathbf{0}$ ) by themselves *do not determine the field*—like any differential equations, they must be supplemented by appropriate *boundary conditions*. Ordinarily, these are so “obvious” that we impose them almost subconsciously (“ $E$  must go to zero far from the source charges”)—or we appeal to symmetry to resolve the ambiguity (“the field must be the same—in magnitude—on both sides of an infinite plane of surface charge”). But in this case there are *no* natural boundary conditions, and no persuasive symmetry conditions, to fix the answer. The question “What is the electric field produced by a uniform charge density filling all of space?” is simply *ill-posed*: it does not give us sufficient information to determine the answer. (Incidentally, it won’t help to appeal to Coulomb’s law ( $\mathbf{E} = \frac{1}{4\pi\epsilon_0} \int \rho \frac{\hat{\mathbf{r}}}{r^2} d\tau$ )—the integral is hopelessly indefinite, in this case.)

### Problem 2.56

Compare Newton’s law of universal gravitation to Coulomb’s law:

$$\mathbf{F} = -G \frac{m_1 m_2}{r^2} \hat{\mathbf{r}}; \quad \mathbf{F} = \frac{1}{4\pi\epsilon_0} \frac{q_1 q_2}{r^2} \hat{\mathbf{r}}.$$

Evidently  $\frac{1}{4\pi\epsilon_0} \rightarrow G$  and  $q \rightarrow m$ . The gravitational energy of a sphere (translating Prob. 2.34) is therefore

$$\boxed{W_{\text{grav}} = \frac{3}{5} G \frac{M^2}{R}}.$$

Now,  $G = 6.67 \times 10^{-11} \text{ N m}^2/\text{kg}^2$ , and for the sun  $M = 1.99 \times 10^{30} \text{ kg}$ ,  $R = 6.96 \times 10^8 \text{ m}$ , so the sun's gravitational energy is  $W = 2.28 \times 10^{41} \text{ J}$ . At the current rate this energy would be dissipated in a time

$$t = \frac{W}{P} = \frac{2.28 \times 10^{41}}{3.86 \times 10^{26}} = 5.90 \times 10^{14} \text{ s} = \boxed{1.87 \times 10^7 \text{ years.}}$$

### Problem 2.57

First eliminate  $z$ , using the formula for the ellipsoid:

$$\sigma(x, y) = \frac{Q}{4\pi ab} \frac{1}{\sqrt{c^2(x^2/a^4) + c^2(y^2/b^4) + 1 - (x^2/a^2) - (y^2/b^2)}}.$$

Now (for parts (a) and (b)) set  $c \rightarrow 0$ , “squashing” the ellipsoid down to an ellipse in the  $xy$  plane:

$$\sigma(x, y) = \frac{Q}{2\pi ab} \frac{1}{\sqrt{1 - (x/a)^2 - (y/b)^2}}.$$

(I multiplied by 2 to count both surfaces.)

(a) For the circular disk, set  $a = b = R$  and let  $r \equiv \sqrt{x^2 + y^2}$ .  $\sigma(r) = \frac{Q}{2\pi R} \frac{1}{\sqrt{R^2 - r^2}}.$

(b) For the ribbon, let  $Q/2b \equiv \Lambda$ , and then take the limit  $b \rightarrow \infty$ :  $\sigma(x) = \frac{\Lambda}{2\pi} \frac{1}{\sqrt{a^2 - x^2}}.$

(c) Let  $b = c$ ,  $r \equiv \sqrt{y^2 + z^2}$ , making an ellipsoid of revolution:



$$\frac{x^2}{a^2} + \frac{r^2}{c^2} = 1, \quad \text{with } \sigma = \frac{Q}{4\pi ac^2} \frac{1}{\sqrt{x^2/a^4 + r^2/c^4}}.$$

The charge on a ring of width  $dx$  is

$$dq = \sigma 2\pi r ds, \quad \text{where } ds = \sqrt{dx^2 + dr^2} = dx \sqrt{1 + (dr/dx)^2}.$$

Now  $\frac{2x dx}{a^2} + \frac{2r dr}{c^2} = 0 \Rightarrow \frac{dr}{dx} = -\frac{c^2 x}{a^2 r}$ , so  $ds = dx \sqrt{1 + \frac{c^4 x^2}{a^4 r^2}} = dx \frac{c^2}{r} \sqrt{x^2/a^4 + r^2/c^4}$ . Thus

$$\lambda(x) = \frac{dq}{dx} = 2\pi r \frac{Q}{4\pi ac^2} \frac{1}{\sqrt{x^2/a^4 + r^2/c^4}} \frac{c^2}{r} \sqrt{x^2/a^4 + r^2/c^4} = \boxed{\frac{Q}{2a}}. \quad (\text{Constant!})$$

**Problem 2.58**

(a) One such point is on the  $x$  axis (see diagram) at  $x = r$ . Here the field is

$$E_x = \frac{q}{4\pi\epsilon_0} \left[ \frac{1}{(a+r)^2} - 2\frac{\cos\theta}{b^2} \right] = 0, \quad \text{or} \quad \frac{2\cos\theta}{b^2} = \frac{1}{(a+r)^2}.$$

Now,

$$\cos\theta = \frac{(a/2) - r}{b}; \quad b^2 = \left(\frac{a}{2} - r\right)^2 + \left(\frac{\sqrt{3}}{2}a\right)^2 = (a^2 - ar + r^2).$$

Therefore

$$\frac{2[(a/2) - r]}{(a^2 - ar + r^2)^{3/2}} = \frac{1}{(a+r)^2}. \quad \text{To simplify, let } \frac{r}{a} \equiv u :$$

$$\frac{(1-2u)}{(1-u+u^2)^{3/2}} = \frac{1}{(1+u)^2}, \quad \text{or} \quad (1-2u)^2(1+u)^4 = (1-u+u^2)^3.$$

Multiplying out each side:

$$1 - 6u^2 - 4u^3 + 9u^4 + 12u^5 + 4u^6 = 1 - 3u + 6u^2 - 7u^3 + 6u^4 - 3u^5 + u^6,$$


or

$$3u - 12u^2 + 3u^3 + 3u^4 + 15u^5 + 3u^6 = 0.$$

$u = 0$  is a solution (of course—the center of the triangle); factoring out  $3u$  we are left with a quintic equation:

$$1 - 4u + u^2 + u^3 + 5u^4 + u^5 = 0.$$

According to Mathematica, this has two complex roots, and one negative root. The two remaining solutions are  $u = 0.284718$  and  $u = 0.626691$ . The latter is outside the triangle, and clearly spurious. So  $\boxed{r = 0.284718a}$ . (The other two places where  $\mathbf{E} = \mathbf{0}$  are at the symmetrically located points, of course.)



(b) For the square:

$$E_x = \frac{q}{4\pi\epsilon_0} \left( 2 \frac{\cos \theta_+}{b_+^2} - 2 \frac{\cos \theta_-}{b_-^2} \right) = 0 \quad \Rightarrow \quad \frac{\cos \theta_+}{b_+^2} = \frac{\cos \theta_-}{b_-^2},$$

where

$$\cos \theta_{\pm} = \frac{(a/\sqrt{2}) \pm r}{b_{\pm}}; \quad b_{\pm}^2 = \left( \frac{a}{\sqrt{2}} \right)^2 + \left( \frac{a}{\sqrt{2}} \pm r \right)^2 = a^2 \pm \sqrt{2}ar + r^2.$$

Thus

$$\frac{(a/\sqrt{2}) + r}{(a^2 + \sqrt{2}ar + r^2)^{3/2}} = \frac{(a/\sqrt{2}) - r}{(a^2 - \sqrt{2}ar + r^2)^{3/2}}.$$

To simplify, let  $w \equiv \sqrt{2}r/a$ ; then

$$\frac{1+w}{(2+2w+w^2)^{3/2}} = \frac{1-w}{(2-2w+w^2)^{3/2}}, \quad \text{or} \quad (1+w)^2(2-2w+w^2)^3 = (1-w)^2(2+2w+w^2)^3.$$

Multiplying out the left side:


$$8 - 8w - 4w^2 + 16w^3 - 10w^4 - 2w^5 + 7w^6 - 4w^7 + w^8 = (\text{same thing with } w \rightarrow -w).$$

The even powers cancel, leaving

$$8w - 16w^3 + 2w^5 + 4w^7 = 0, \quad \text{or} \quad 4 - 8v + v^2 + 2v^3 = 0,$$

where  $v \equiv w^2$ . According to Mathematica, this cubic equation has one negative root, one root that is spurious (the point lies outside the square), and  $v = 0.598279$ , which yields

$$r = \sqrt{\frac{v}{2}} a = \boxed{0.546936 a}.$$



For the pentagon:

$$E_x = \frac{q}{4\pi\epsilon_0} \left( \frac{1}{(a+r)^2} + 2\frac{\cos\theta}{b^2} - 2\frac{\cos\phi}{c^2} \right) = 0,$$

where

$$\cos\theta = \frac{a \cos(2\pi/5) + r}{b}, \quad \cos\phi = \frac{a \cos(\pi/5) - r}{c};$$

$$b^2 = [a \cos(2\pi/5) + r]^2 + [a \sin(2\pi/5)]^2 = a^2 + r^2 + 2ar \cos(2\pi/5),$$

$$c^2 = [a \cos(\pi/5) - r]^2 + [a \sin(\pi/5)]^2 = a^2 + r^2 - 2ar \cos(\pi/5).$$

$$\frac{1}{(a+r)^2} + 2\frac{r + a \cos(2\pi/5)}{[a^2 + r^2 + 2ar \cos(2\pi/5)]^{3/2}} + 2\frac{r - a \cos(\pi/5)}{[a^2 + r^2 - 2ar \cos(\pi/5)]^{3/2}} = 0.$$

Mathematica gives the solution  $\boxed{r = 0.688917 a}.$

For an  $n$ -sided regular polygon there are evidently  $n$  such points, lying on the radial spokes that bisect the sides; their distance from the center appears to grow monotonically with  $n$ :  $r(3) = 0.285$ ,  $r(4) = 0.547$ ,  $r(5) = 0.689$ ,  $\dots$ . As  $n \rightarrow \infty$  they fill out a circle that (in the limit) coincides with the ring of charge itself.

**Problem 2.59** The theorem is *false*. For example, suppose the conductor is a neutral sphere and the external field is due to a nearby positive point charge  $q$ . A negative charge will be induced on the near side of the sphere (and a positive charge on the far side), so the force will be *attractive* (toward  $q$ ). If we now reverse the sign of  $q$ , the induced charges will also reverse, but the force will still be attractive.

If the external field is *uniform*, then the net force on the induced charges is zero, and the total force on the conductor is  $Q\mathbf{E}_e$ , which *does* switch signs if  $\mathbf{E}_e$  is reversed. So the “theorem” is valid in this very special case.



**Problem 2.60** The initial configuration consists of a point charge  $q$  at the center,  $-q$  induced on the inner surface, and  $+q$  on the outer surface. What is the energy of this configuration? Imagine assembling it piece-by-piece. First bring in  $q$  and place it at the origin—this takes no work. Now bring in  $-q$  and spread it over the surface at  $a$ —using the method in Prob. 2.35, this takes work  $-q^2/(8\pi\epsilon_0 a)$ . Finally, bring in  $+q$  and spread it over the surface at  $b$ —this costs  $q^2/(8\pi\epsilon_0 b)$ . Thus the energy of the initial configuration is


$$W_i = -\frac{q^2}{8\pi\epsilon_0} \left( \frac{1}{a} - \frac{1}{b} \right).$$

The final configuration is a neutral shell and a distant point charge—the energy is zero. Thus the work necessary to go from the initial to the final state is

$$W = W_f - W_i = \frac{q^2}{8\pi\epsilon_0} \left( \frac{1}{a} - \frac{1}{b} \right).$$

---

**Problem 2.61**



Suppose the  $n$  point charges are evenly spaced around the circle, with the  $j$ th particle at angle  $j(2\pi/n)$ . According to Eq. 2.42, the energy of the configuration is

$$W_n = n \frac{1}{2} qV,$$

where  $V$  is the potential due to the  $(n - 1)$  other charges, at charge #  $n$  (on the  $x$  axis).

$$V = \frac{1}{4\pi\epsilon_0} q \sum_{j=1}^{n-1} \frac{1}{r_j}, \quad r_j = 2R \sin\left(\frac{j\pi}{n}\right)$$

(see the figure). So

$$W_n = \frac{q^2}{4\pi\epsilon_0 R} \frac{n}{4} \sum_{j=1}^{n-1} \frac{1}{\sin(j\pi/n)} = \frac{q^2}{4\pi\epsilon_0 R} \Omega_n.$$

Mathematica says

$$\Omega_{10} = \frac{10}{4} \sum_{j=1}^9 \frac{1}{\sin(j\pi/10)} = 38.6245$$

$$\Omega_{11} = \frac{11}{4} \sum_{j=1}^{10} \frac{1}{\sin(j\pi/11)} = \boxed{48.5757}$$

$$\Omega_{12} = \frac{12}{4} \sum_{j=1}^{11} \frac{1}{\sin(j\pi/12)} = \boxed{59.8074}$$

If  $(n-1)$  charges are on the circle (energy  $\Omega_{n-1}q^2/4\pi\epsilon_0 R$ ), and the  $n$ th is at the center, the total energy is

$$W_n = [\Omega_{n-1} + (n-1)] \frac{q^2}{4\pi\epsilon_0 R}.$$

For

$$n = 11 : \quad \Omega_{10} + 10 = 38.6245 + 10 = \boxed{48.6245} > \Omega_{11}$$

$$n = 12 : \quad \Omega_{11} + 11 = 48.5757 + 11 = \boxed{59.5757} < \Omega_{12}$$

Thus a lower energy is achieved for 11 charges if they are all at the rim, but for 12 it is better to put one at the center.

---

## Chapter 3

# Potential

### Problem 3.1

The argument is exactly the same as in Sect. 3.1.4, except that since  $z < R$ ,  $\sqrt{z^2 + R^2 - 2zR} = (R - z)$ , instead of  $(z - R)$ . Hence  $V_{\text{ave}} = \frac{q}{4\pi\epsilon_0} \frac{1}{2zR} [(z + R) - (R - z)] = \frac{1}{4\pi\epsilon_0} \frac{q}{R}$ . If there is more than one charge inside the sphere, the average potential due to interior charges is  $\frac{1}{4\pi\epsilon_0} \frac{Q_{\text{enc}}}{R}$ , and the average due to exterior charges is  $V_{\text{center}}$ , so  $V_{\text{ave}} = V_{\text{center}} + \frac{Q_{\text{enc}}}{4\pi\epsilon_0 R}$ . ✓

### Problem 3.2

A stable equilibrium is a point of local minimum in the potential energy. Here the potential energy is  $qV$ . But we know that Laplace's equation allows no local minima for  $V$ . What *looks* like a minimum, in the figure, must in fact be a saddle point, and the box "leaks" through the center of each face.

### Problem 3.3

Laplace's equation in *spherical* coordinates, for  $V$  dependent only on  $r$ , reads:

$$\nabla^2 V = \frac{1}{r^2} \frac{d}{dr} \left( r^2 \frac{dV}{dr} \right) = 0 \Rightarrow r^2 \frac{dV}{dr} = c \text{ (constant)} \Rightarrow \frac{dV}{dr} = \frac{c}{r^2} \Rightarrow \boxed{V = -\frac{c}{r} + k.}$$

*Example:* potential of a uniformly charged sphere.

In *cylindrical* coordinates:  $\nabla^2 V = \frac{1}{s} \frac{d}{ds} \left( s \frac{dV}{ds} \right) = 0 \Rightarrow s \frac{dV}{ds} = c \Rightarrow \frac{dV}{ds} = \frac{c}{s} \Rightarrow \boxed{V = c \ln s + k.}$

*Example:* potential of a long wire.

### Problem 3.4

Refer to Fig. 3.3, letting  $\alpha$  be the angle between  $\mathbf{r}$  and the  $z$  axis. Obviously,  $\mathbf{E}_{\text{ave}}$  points in the  $-\hat{\mathbf{z}}$  direction, so

$$\mathbf{E}_{\text{ave}} = \frac{1}{4\pi R^2} \oint \mathbf{E} da = -\hat{\mathbf{z}} \frac{1}{4\pi R^2} \frac{q}{4\pi\epsilon_0} \int \frac{1}{r^2} \cos \alpha da.$$

By the law of cosines,

$$\begin{aligned} R^2 &= z^2 + r^2 - 2zr \cos \alpha \Rightarrow \cos \alpha = \frac{z^2 + r^2 - R^2}{2zr}, \\ r^2 &= R^2 + z^2 - 2Rz \cos \theta \Rightarrow \frac{\cos \alpha}{r^2} = \frac{z^2 + r^2 - R^2}{2zr^3} = \frac{z - R \cos \theta}{(R^2 + z^2 - 2Rz \cos \theta)^{3/2}}. \end{aligned}$$

$$\begin{aligned}\mathbf{E}_{\text{ave}} &= -\hat{\mathbf{z}} \frac{q}{16\pi^2 R^2 \epsilon_0} \int \frac{z - R \cos \theta}{(R^2 + z^2 - 2Rz \cos \theta)^{3/2}} R^2 \sin \theta d\theta d\phi \\ &= -\frac{q\hat{\mathbf{z}}}{8\pi\epsilon_0} \int_0^\pi \frac{z - R \cos \theta}{(R^2 + z^2 - 2Rz \cos \theta)^{3/2}} \sin \theta d\theta = -\frac{q\hat{\mathbf{z}}}{8\pi\epsilon_0} \int_{-1}^1 \frac{z - Ru}{(R^2 + z^2 - 2Rzu)^{3/2}} du\end{aligned}$$

(where  $u \equiv \cos \theta$ ). The integral is

$$\begin{aligned}I &= \frac{1}{R\sqrt{R^2 + z^2 - 2Rzu}} \Big|_{-1}^1 - \frac{1}{2Rz^2} \left( \sqrt{R^2 + z^2 - 2Rzu} + \frac{R^2 + z^2}{\sqrt{R^2 + z^2 - 2Rzu}} \right) \Big|_{-1}^1 \\ &= \frac{1}{R} \left( \frac{1}{|z - R|} - \frac{1}{z + R} \right) - \frac{1}{2Rz^2} \left[ |z - R| - (z + R) + (R^2 + z^2) \left( \frac{1}{|z - R|} - \frac{1}{z + R} \right) \right].\end{aligned}$$

(a) If  $z > R$ ,

$$\begin{aligned}I &= \frac{1}{R} \left( \frac{1}{z - R} - \frac{1}{z + R} \right) - \frac{1}{2Rz^2} \left[ (z - R) - (z + R) + (R^2 + z^2) \left( \frac{1}{z - R} - \frac{1}{z + R} \right) \right] \\ &= \frac{1}{R} \left( \frac{2R}{z^2 - R^2} \right) - \frac{1}{2Rz^2} \left[ -2R + (R^2 + z^2) \frac{2R}{z^2 - R^2} \right] = \frac{2}{z^2}.\end{aligned}$$

So

$$\mathbf{E}_{\text{ave}} = -\frac{1}{4\pi\epsilon_0} \frac{q}{z^2} \hat{\mathbf{z}},$$

the same as the field at the center. By superposition the same holds for any *collection* of charges outside the sphere.

(b) If  $z < R$ ,

$$\begin{aligned}I &= \frac{1}{R} \left( \frac{1}{R - z} - \frac{1}{z + R} \right) - \frac{1}{2Rz^2} \left[ (R - z) - (z + R) + (R^2 + z^2) \left( \frac{1}{R - z} - \frac{1}{z + R} \right) \right] \\ &= \frac{1}{R} \left( \frac{2z}{R^2 - z^2} \right) - \frac{1}{2Rz^2} \left[ -2z + (R^2 + z^2) \frac{2z}{R^2 - z^2} \right] = 0.\end{aligned}$$

So

$$\mathbf{E}_{\text{ave}} = \mathbf{0}.$$

By superposition the same holds for any *collection* of charges inside the sphere.

### Problem 3.5

Same as proof of second uniqueness theorem, up to the equation  $\oint_S V_3 \mathbf{E}_3 \cdot d\mathbf{a} = -\int_V (E_3)^2 d\tau$ . But on each surface, either  $V_3 = 0$  (if  $V$  is specified on the surface), or else  $E_{3\perp} = 0$  (if  $\frac{\partial V}{\partial n} = -E_\perp$  is specified). So  $\oint_V (E_3)^2 = 0$ , and hence  $\mathbf{E}_2 = \mathbf{E}_1$ . qed

### Problem 3.6

Putting  $U = T = V_3$  into Green's identity:

$$\int_V [V_3 \nabla^2 V_3 + \nabla V_3 \cdot \nabla V_3] d\tau = \oint_S V_3 \nabla V_3 \cdot d\mathbf{a}. \quad \text{But } \nabla^2 V_3 = \nabla^2 V_1 - \nabla^2 V_2 = -\frac{\rho}{\epsilon_0} + \frac{\rho}{\epsilon_0} = 0, \text{ and } \nabla V_3 = -\mathbf{E}_3.$$

So  $\int_V E_3^2 d\tau = -\oint_S V_3 \mathbf{E}_3 \cdot d\mathbf{a}$ , and the rest is the same as before.

### Problem 3.7

Place image charges  $+2q$  at  $z = -d$  and  $-q$  at  $z = -3d$ . Total force on  $+q$  is

$$\mathbf{F} = \frac{q}{4\pi\epsilon_0} \left[ \frac{-2q}{(2d)^2} + \frac{2q}{(4d)^2} + \frac{-q}{(6d)^2} \right] \hat{\mathbf{z}} = \frac{q^2}{4\pi\epsilon_0 d^2} \left( -\frac{1}{2} + \frac{1}{8} - \frac{1}{36} \right) \hat{\mathbf{z}} = \boxed{-\frac{1}{4\pi\epsilon_0} \left( \frac{29q^2}{72d^2} \right) \hat{\mathbf{z}}}.$$

**Problem 3.8**

(a) From Fig. 3.13:  $z = \sqrt{r^2 + a^2 - 2ra \cos \theta}$ ;  $z' = \sqrt{r^2 + b^2 - 2rb \cos \theta}$ . Therefore:

$$\begin{aligned} \frac{q'}{z'} &= -\frac{R}{a} \frac{q}{\sqrt{r^2 + b^2 - 2rb \cos \theta}} \quad (\text{Eq. 3.15}), \text{ while } b = \frac{R^2}{a} \quad (\text{Eq. 3.16}). \\ &= -\frac{q}{\left(\frac{a}{R}\right) \sqrt{r^2 + \frac{R^4}{a^2} - 2r\frac{R^2}{a} \cos \theta}} = -\frac{q}{\sqrt{\left(\frac{ar}{R}\right)^2 + R^2 - 2ra \cos \theta}}. \end{aligned}$$

Therefore:

$$V(r, \theta) = \frac{1}{4\pi\epsilon_0} \left( \frac{q}{z} + \frac{q'}{z'} \right) = \frac{q}{4\pi\epsilon_0} \left\{ \frac{1}{\sqrt{r^2 + a^2 - 2ra \cos \theta}} - \frac{1}{\sqrt{R^2 + (ra/R)^2 - 2ra \cos \theta}} \right\}.$$

Clearly, when  $r = R$ ,  $V \rightarrow 0$ .

(b)  $\sigma = -\epsilon_0 \frac{\partial V}{\partial n}$  (Eq. 2.49). In this case,  $\frac{\partial V}{\partial n} = \frac{\partial V}{\partial r}$  at the point  $r = R$ . Therefore,

$$\begin{aligned} \sigma(\theta) &= -\epsilon_0 \left( \frac{q}{4\pi\epsilon_0} \right) \left\{ -\frac{1}{2} (r^2 + a^2 - 2ra \cos \theta)^{-3/2} (2r - 2a \cos \theta) \right. \\ &\quad \left. + \frac{1}{2} (R^2 + (ra/R)^2 - 2ra \cos \theta)^{-3/2} \left( \frac{a^2}{R^2} 2r - 2a \cos \theta \right) \right\} \Big|_{r=R} \\ &= -\frac{q}{4\pi} \left\{ -(R^2 + a^2 - 2Ra \cos \theta)^{-3/2} (R - a \cos \theta) + (R^2 + a^2 - 2Ra \cos \theta)^{-3/2} \left( \frac{a^2}{R} - a \cos \theta \right) \right\} \\ &= \frac{q}{4\pi} (R^2 + a^2 - 2Ra \cos \theta)^{-3/2} \left[ R - a \cos \theta - \frac{a^2}{R} + a \cos \theta \right] \\ &= \frac{q}{4\pi R} (R^2 - a^2) (R^2 + a^2 - 2Ra \cos \theta)^{-3/2}. \\ q_{\text{induced}} &= \int \sigma da = \frac{q}{4\pi R} (R^2 - a^2) \int (R^2 + a^2 - 2Ra \cos \theta)^{-3/2} R^2 \sin \theta d\theta d\phi \\ &= \frac{q}{4\pi R} (R^2 - a^2) 2\pi R^2 \left[ -\frac{1}{Ra} (R^2 + a^2 - 2Ra \cos \theta)^{-1/2} \right] \Big|_0^\pi \\ &= \frac{q}{2a} (a^2 - R^2) \left[ \frac{1}{\sqrt{R^2 + a^2 + 2Ra}} - \frac{1}{\sqrt{R^2 + a^2 - 2Ra}} \right]. \\ \text{But } a > R \text{ (else } q \text{ would be inside), so } \sqrt{R^2 + a^2 - 2Ra} &= a - R. \\ &= \frac{q}{2a} (a^2 - R^2) \left[ \frac{1}{(a + R)} - \frac{1}{(a - R)} \right] = \frac{q}{2a} [(a - R) - (a + R)] = \frac{q}{2a} (-2R) \\ &= -\frac{qR}{a} = q'. \end{aligned}$$

(c) The force on  $q$ , due to the sphere, is the same as the force of the image charge  $q'$ , to wit:

$$F = \frac{1}{4\pi\epsilon_0} \frac{qq'}{(a-b)^2} = \frac{1}{4\pi\epsilon_0} \left( -\frac{R}{a} q^2 \right) \frac{1}{(a - R^2/a)^2} = -\frac{1}{4\pi\epsilon_0} \frac{q^2 Ra}{(a^2 - R^2)^2}.$$

To bring  $q$  in from infinity to  $a$ , then, we do work

$$W = \frac{q^2 R}{4\pi\epsilon_0} \int_{\infty}^a \frac{\bar{a}}{(\bar{a}^2 - R^2)^2} d\bar{a} = \frac{q^2 R}{4\pi\epsilon_0} \left[ -\frac{1}{2} \frac{1}{(\bar{a}^2 - R^2)} \right]_{\infty}^a = \boxed{-\frac{1}{4\pi\epsilon_0} \frac{q^2 R}{2(a^2 - R^2)}}.$$

### Problem 3.9

Place a second image charge,  $q''$ , at the *center* of the sphere; this will not alter the fact that the sphere is an *equipotential*, but merely *increase* that potential from zero to  $V_0 = \frac{1}{4\pi\epsilon_0} \frac{q''}{R}$ ;

$$\boxed{q'' = 4\pi\epsilon_0 V_0 R \text{ at center of sphere.}}$$


For a *neutral* sphere,  $q' + q'' = 0$ .

$$\begin{aligned} F &= \frac{1}{4\pi\epsilon_0} q \left( \frac{q''}{a^2} + \frac{q'}{(a-b)^2} \right) = \frac{qq'}{4\pi\epsilon_0} \left( -\frac{1}{a^2} + \frac{1}{(a-b)^2} \right) \\ &= \frac{qq'}{4\pi\epsilon_0} \frac{b(2a-b)}{a^2(a-b)^2} = \frac{q(-Rq/a)(R^2/a)(2a-R^2/a)}{4\pi\epsilon_0 a^2(a-R^2/a)^2} \\ &= -\boxed{\frac{q^2}{4\pi\epsilon_0} \left( \frac{R}{a} \right)^3 \frac{(2a^2 - R^2)}{(a^2 - R^2)^2}}. \end{aligned}$$


(Drop the minus sign, because the problem asks for the force of *attraction*.)

### Problem 3.10

(a) Image problem:  $\lambda$  above,  $-\lambda$  below. Potential was found in Prob. 2.52:



$$\begin{aligned} V(y, z) &= \frac{2\lambda}{4\pi\epsilon_0} \ln(s_-/s_+) = \frac{\lambda}{4\pi\epsilon_0} \ln(s_-^2/s_+^2) \\ &= \boxed{\frac{\lambda}{4\pi\epsilon_0} \ln \left\{ \frac{y^2 + (z+d)^2}{y^2 + (z-d)^2} \right\}} \end{aligned}$$



(b)  $\sigma = -\epsilon_0 \frac{\partial V}{\partial n}$ . Here  $\frac{\partial V}{\partial n} = \frac{\partial V}{\partial z}$ , evaluated at  $z = 0$ .

$$\begin{aligned} \sigma(y) &= -\epsilon_0 \frac{\lambda}{4\pi\epsilon_0} \left\{ \frac{1}{y^2 + (z+d)^2} 2(z+d) - \frac{1}{y^2 + (z-d)^2} 2(z-d) \right\} \Big|_{z=0} \\ &= -\frac{2\lambda}{4\pi} \left\{ \frac{d}{y^2 + d^2} - \frac{-d}{y^2 + d^2} \right\} = \boxed{-\frac{\lambda d}{\pi(y^2 + d^2)}}. \end{aligned}$$


*Check:* Total charge induced on a strip of width  $l$  parallel to the  $y$  axis:

$$\begin{aligned} q_{\text{ind}} &= -\frac{l\lambda d}{\pi} \int_{-\infty}^{\infty} \frac{1}{y^2 + d^2} dy = -\frac{l\lambda d}{\pi} \left[ \frac{1}{d} \tan^{-1} \left( \frac{y}{d} \right) \right]_{-\infty}^{\infty} = -\frac{l\lambda d}{\pi} \left[ \frac{\pi}{2} - \left( -\frac{\pi}{2} \right) \right] \\ &= -\lambda l. \quad \text{Therefore } \lambda_{\text{ind}} = -\lambda, \text{ as it should be.} \end{aligned}$$

**Problem 3.11**

The image configuration is as shown.

$$V(x, y) = \frac{q}{4\pi\epsilon_0} \left\{ \frac{1}{\sqrt{(x-a)^2 + (y-b)^2 + z^2}} + \frac{1}{\sqrt{(x+a)^2 + (y+b)^2 + z^2}} - \frac{1}{\sqrt{(x+a)^2 + (y-b)^2 + z^2}} - \frac{1}{\sqrt{(x-a)^2 + (y+b)^2 + z^2}} \right\}.$$



$$\mathbf{F} = \frac{q^2}{4\pi\epsilon_0} \left\{ -\frac{1}{(2a)^2} \hat{\mathbf{x}} - \frac{1}{(2b)^2} \hat{\mathbf{y}} + \frac{1}{(2\sqrt{a^2 + b^2})^2} [\cos \theta \hat{\mathbf{x}} + \sin \theta \hat{\mathbf{y}}] \right\},$$


where  $\cos \theta = a/\sqrt{a^2 + b^2}$ ,  $\sin \theta = b/\sqrt{a^2 + b^2}$ .

$$\mathbf{F} = \frac{q^2}{16\pi\epsilon_0} \left\{ \left[ \frac{a}{(a^2 + b^2)^{3/2}} - \frac{1}{a^2} \right] \hat{\mathbf{x}} + \left[ \frac{b}{(a^2 + b^2)^{3/2}} - \frac{1}{b^2} \right] \hat{\mathbf{y}} \right\}.$$

$$W = \frac{1}{2} \frac{1}{4\pi\epsilon_0} \left[ \frac{-q^2}{(2a)} + \frac{-q^2}{(2b)} + \frac{q^2}{(2\sqrt{a^2 + b^2})} \right] = \frac{q^2}{16\pi\epsilon_0} \left[ \frac{1}{\sqrt{a^2 + b^2}} - \frac{1}{a} - \frac{1}{b} \right].$$


For this to work,  $\theta$  must be an integer divisor of  $180^\circ$ . Thus  $180^\circ$ ,  $90^\circ$ ,  $60^\circ$ ,  $45^\circ$ , etc., are OK, but no others. It works for  $45^\circ$ , say, with the charges as shown.

(Note the strategy: to make the  $x$  axis an equipotential ( $V = 0$ ), you place the image charge (1) in the reflection point. To make the  $45^\circ$  line an equipotential, you place charge (2) at the image point. But that screws up the  $x$  axis, so you must now insert image (3) to balance (2). Moreover, to make the  $45^\circ$  line  $V = 0$  you also need (4), to balance (1). But now, to restore the  $x$  axis to  $V = 0$  you need (5) to balance (4), and so on.



why it works for  $\theta = 45^\circ$

The reason this doesn't work for *arbitrary* angles is that you are eventually forced to place an image charge *within the original region of interest*, and that's not allowed—all images must go *outside* the region, or you're no longer dealing with the (same) problem at all.)



why it *doesn't* work for  $\theta = 135^\circ$

**Problem 3.12**

From Prob. 2.52 (with  $y_0 \rightarrow d$ ):  $V = \frac{\lambda}{4\pi\epsilon_0} \ln \left[ \frac{(x+a)^2 + y^2}{(x-a)^2 + y^2} \right]$ , where  $a^2 = y_0^2 - R^2 \Rightarrow a = \sqrt{d^2 - R^2}$ ,

and

$$\left\{ \begin{array}{l} a \coth(2\pi\epsilon_0 V_0/\lambda) = d \\ a \operatorname{csch}(2\pi\epsilon_0 V_0/\lambda) = R \end{array} \right\} \Rightarrow (\text{dividing}) \quad \frac{d}{R} = \cosh \left( \frac{2\pi\epsilon_0 V_0}{\lambda} \right), \text{ or } \lambda = \frac{2\pi\epsilon_0 V_0}{\cosh^{-1}(d/R)}.$$

**Problem 3.13**

$$V(x, y) = \sum_{n=1}^{\infty} C_n e^{-n\pi x/a} \sin(n\pi y/a) \quad (\text{Eq. 3.30}), \quad \text{where } C_n = \frac{2}{a} \int_0^a V_0(y) \sin(n\pi y/a) dy \quad (\text{Eq. 3.34}).$$

In this case  $V_0(y) = \begin{cases} +V_0, & \text{for } 0 < y < a/2 \\ -V_0, & \text{for } a/2 < y < a \end{cases}$ . Therefore,

$$\begin{aligned} C_n &= \frac{2}{a} V_0 \left\{ \int_0^{a/2} \sin(n\pi y/a) dy - \int_{a/2}^a \sin(n\pi y/a) dy \right\} = \frac{2V_0}{a} \left\{ -\frac{\cos(n\pi y/a)}{(n\pi/a)} \Big|_0^{a/2} + \frac{\cos(n\pi y/a)}{(n\pi/a)} \Big|_{a/2}^a \right\} \\ &= \frac{2V_0}{n\pi} \left\{ -\cos\left(\frac{n\pi}{2}\right) + \cos(0) + \cos(n\pi) - \cos\left(\frac{n\pi}{2}\right) \right\} = \frac{2V_0}{n\pi} \left\{ 1 + (-1)^n - 2\cos\left(\frac{n\pi}{2}\right) \right\}. \end{aligned}$$

The term in curly brackets is:

$$\left\{ \begin{array}{ll} n=1 & : 1 - 1 - 2\cos(\pi/2) = 0, \\ n=2 & : 1 + 1 - 2\cos(\pi) = 4, \\ n=3 & : 1 - 1 - 2\cos(3\pi/2) = 0, \\ n=4 & : 1 + 1 - 2\cos(2\pi) = 0, \end{array} \right\} \text{ etc. (Zero if } n \text{ is odd or divisible by 4, otherwise 4.)}$$

Therefore

$$C_n = \begin{cases} 8V_0/n\pi, & n = 2, 6, 10, 14, \text{ etc. (in general, } 4j+2, \text{ for } j = 0, 1, 2, \dots), \\ 0, & \text{otherwise.} \end{cases}$$

So

$$V(x, y) = \frac{8V_0}{\pi} \sum_{n=2,6,10,\dots} \frac{e^{-n\pi x/a} \sin(n\pi y/a)}{n} = \frac{8V_0}{\pi} \sum_{j=0}^{\infty} \frac{e^{-(4j+2)\pi x/a} \sin[(4j+2)\pi y/a]}{(4j+2)}.$$

### Problem 3.14

$$V(x, y) = \frac{4V_0}{\pi} \sum_{n=1,3,5,\dots} \frac{1}{n} e^{-n\pi x/a} \sin(n\pi y/a) \quad (\text{Eq. 3.36}); \quad \sigma = -\epsilon_0 \frac{\partial V}{\partial n} \quad (\text{Eq. 2.49}).$$

So

$$\begin{aligned} \sigma(y) &= -\epsilon_0 \frac{\partial}{\partial x} \left\{ \frac{4V_0}{\pi} \sum \frac{1}{n} e^{-n\pi x/a} \sin(n\pi y/a) \right\} \Big|_{x=0} = -\epsilon_0 \frac{4V_0}{\pi} \sum \frac{1}{n} \left(-\frac{n\pi}{a}\right) e^{-n\pi x/a} \sin(n\pi y/a) \Big|_{x=0} \\ &= \frac{4\epsilon_0 V_0}{a} \sum_{n=1,3,5,\dots} \sin(n\pi y/a). \end{aligned}$$

Or, using the closed form 3.37:

$$\begin{aligned} V(x, y) &= \frac{2V_0}{\pi} \tan^{-1} \left( \frac{\sin(\pi y/a)}{\sinh(\pi x/a)} \right) \Rightarrow \sigma = -\epsilon_0 \frac{2V_0}{\pi} \frac{1}{1 + \frac{\sin^2(\pi y/a)}{\sinh^2(\pi x/a)}} \left( \frac{-\sin(\pi y/a)}{\sinh^2(\pi x/a)} \right) \frac{\pi}{a} \cosh(\pi x/a) \Big|_{x=0} \\ &= \frac{2\epsilon_0 V_0}{a} \frac{\sin(\pi y/a) \cosh(\pi x/a)}{\sin^2(\pi y/a) + \sinh^2(\pi x/a)} \Big|_{x=0} = \frac{2\epsilon_0 V_0}{a} \frac{1}{\sin(\pi y/a)}. \end{aligned}$$

[*Comment:* Technically, the series solution for  $\sigma$  is defective, since term-by-term differentiation has produced a (naively) non-convergent sum. More sophisticated definitions of convergence permit one to work with series of this form, but it is better to sum the series *first* and *then* differentiate (the second method.)]



**Summation of series Eq. 3.36**

$$V(x, y) = \frac{4V_0}{\pi} I, \text{ where } I \equiv \sum_{n=1,3,5,\dots} \frac{1}{n} e^{-n\pi x/a} \sin(n\pi y/a).$$

Now  $\sin w = \mathcal{I}m(e^{iw})$ , so

$$I = \mathcal{I}m \sum_n \frac{1}{n} e^{-n\pi x/a} e^{in\pi y/a} = \mathcal{I}m \sum_n \frac{1}{n} \mathcal{Z}^n,$$

where  $\mathcal{Z} \equiv e^{-\pi(x-iy)/a}$ . Now

$$\begin{aligned} \sum_{1,3,5,\dots} \frac{1}{n} \mathcal{Z}^n &= \sum_{j=0}^{\infty} \frac{1}{(2j+1)} \mathcal{Z}^{(2j+1)} = \int_0^{\mathcal{Z}} \left\{ \sum_{j=0}^{\infty} u^{2j} \right\} du \\ &= \int_0^{\mathcal{Z}} \frac{1}{1-u^2} du = \frac{1}{2} \ln \left( \frac{1+\mathcal{Z}}{1-\mathcal{Z}} \right) = \frac{1}{2} \ln (Re^{i\theta}) = \frac{1}{2} (\ln R + i\theta), \end{aligned}$$

where  $Re^{i\theta} = \frac{1+\mathcal{Z}}{1-\mathcal{Z}}$ . Therefore

$$\begin{aligned} I &= \mathcal{I}m \left\{ \frac{1}{2} (\ln R + i\theta) \right\} = \frac{1}{2} \theta. \quad \text{But } \frac{1+\mathcal{Z}}{1-\mathcal{Z}} = \frac{1+e^{-\pi(x-iy)/a}}{1-e^{-\pi(x-iy)/a}} = \frac{(1+e^{-\pi(x-iy)/a})(1-e^{-\pi(x+iy)/a})}{(1-e^{-\pi(x-iy)/a})(1-e^{-\pi(x+iy)/a})} \\ &= \frac{1+e^{-\pi x/a} (e^{i\pi y/a} - e^{-i\pi y/a}) - e^{-2\pi x/a}}{|1-e^{-\pi(x-iy)/a}|^2} = \frac{1+2ie^{-\pi x/a} \sin(\pi y/a) - e^{-2\pi x/a}}{|1-e^{-\pi(x-iy)/a}|^2}, \end{aligned}$$

so

$$\tan \theta = \frac{2e^{-\pi x/a} \sin(\pi y/a)}{1-e^{-2\pi x/a}} = \frac{2 \sin(\pi y/a)}{e^{\pi x/a} - e^{-\pi x/a}} = \frac{\sin(\pi y/a)}{\sinh(\pi x/a)}.$$


Therefore

$$I = \frac{1}{2} \tan^{-1} \left( \frac{\sin(\pi y/a)}{\sinh(\pi x/a)} \right), \text{ and } \boxed{V(x, y) = \frac{2V_0}{\pi} \tan^{-1} \left( \frac{\sin(\pi y/a)}{\sinh(\pi x/a)} \right)}.$$

**Problem 3.15**

(a)  $\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} = 0$ , with boundary conditions

$$\left\{ \begin{array}{l} \text{(i)} \quad V(x, 0) = 0, \\ \text{(ii)} \quad V(x, a) = 0, \\ \text{(iii)} \quad V(0, y) = 0, \\ \text{(iv)} \quad V(b, y) = V_0(y). \end{array} \right\}$$



As in Ex. 3.4, separation of variables yields

$$V(x, y) = (Ae^{kx} + Be^{-kx}) (C \sin ky + D \cos ky).$$

Here (i)  $\Rightarrow D = 0$ , (iii)  $\Rightarrow B = -A$ , (ii)  $\Rightarrow ka$  is an integer multiple of  $\pi$ :

$$V(x, y) = AC \left( e^{n\pi x/a} - e^{-n\pi x/a} \right) \sin(n\pi y/a) = (2AC) \sinh(n\pi x/a) \sin(n\pi y/a).$$

But  $(2AC)$  is a constant, and the most general linear combination of separable solutions consistent with (i), (ii), (iii) is

$$V(x, y) = \sum_{n=1}^{\infty} C_n \sinh(n\pi x/a) \sin(n\pi y/a).$$

It remains to determine the coefficients  $C_n$  so as to fit boundary condition (iv):

$$\sum C_n \sinh(n\pi b/a) \sin(n\pi y/a) = V_0(y). \text{ Fourier's trick } \Rightarrow C_n \sinh(n\pi b/a) = \frac{2}{a} \int_0^a V_0(y) \sin(n\pi y/a) dy.$$

Therefore

$$C_n = \frac{2}{a \sinh(n\pi b/a)} \int_0^a V_0(y) \sin(n\pi y/a) dy.$$

$$(b) \ C_n = \frac{2}{a \sinh(n\pi b/a)} V_0 \int_0^a \sin(n\pi y/a) dy = \frac{2V_0}{a \sinh(n\pi b/a)} \times \begin{cases} 0, & \text{if } n \text{ is even,} \\ \frac{2a}{n\pi}, & \text{if } n \text{ is odd.} \end{cases}$$

$$V(x, y) = \frac{4V_0}{\pi} \sum_{n=1,3,5,\dots} \frac{\sinh(n\pi x/a) \sin(n\pi y/a)}{n \sinh(n\pi b/a)}.$$

### Problem 3.16

Same format as Ex. 3.5, only the boundary conditions are:

$$\begin{cases} \text{(i)} & V = 0 & \text{when } x = 0, \\ \text{(ii)} & V = 0 & \text{when } x = a, \\ \text{(iii)} & V = 0 & \text{when } y = 0, \\ \text{(iv)} & V = 0 & \text{when } y = a, \\ \text{(v)} & V = 0 & \text{when } z = 0, \\ \text{(vi)} & V = V_0 & \text{when } z = a. \end{cases}$$

This time we want sinusoidal functions in  $x$  and  $y$ , exponential in  $z$ :

$$X(x) = A \sin(kx) + B \cos(kx), \quad Y(y) = C \sin.ly) + D \cos.ly), \quad Z(z) = E e^{\sqrt{k^2+l^2}z} + G e^{-\sqrt{k^2+l^2}z}.$$

(i)  $\Rightarrow B = 0$ ; (ii)  $\Rightarrow k = n\pi/a$ ; (iii)  $\Rightarrow D = 0$ ; (iv)  $\Rightarrow l = m\pi/a$ ; (v)  $\Rightarrow E + G = 0$ . Therefore

$$Z(z) = 2E \sinh(\pi \sqrt{n^2 + m^2} z/a).$$

Putting this all together, and combining the constants, we have:

$$V(x, y, z) = \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} C_{n,m} \sin(n\pi x/a) \sin(m\pi y/a) \sinh(\pi \sqrt{n^2 + m^2} z/a).$$

It remains to evaluate the constants  $C_{n,m}$ , by imposing boundary condition (vi):

$$V_0 = \sum \sum \left[ C_{n,m} \sinh(\pi \sqrt{n^2 + m^2}) \right] \sin(n\pi x/a) \sin(m\pi y/a).$$

According to Eqs. 3.50 and 3.51:

$$C_{n,m} \sinh(\pi\sqrt{n^2+m^2}) = \left(\frac{2}{a}\right)^2 V_0 \int_0^a \int_0^a \sin(n\pi x/a) \sin(m\pi y/a) dx dy = \begin{cases} 0, & \text{if } n \text{ or } m \text{ is even,} \\ \frac{16V_0}{\pi^2 nm}, & \text{if both are odd.} \end{cases}$$

Therefore

$$V(x, y, z) = \frac{16V_0}{\pi^2} \sum_{n=1,3,5,\dots} \sum_{m=1,3,5,\dots} \frac{1}{nm} \sin(n\pi x/a) \sin(m\pi y/a) \frac{\sinh(\pi\sqrt{n^2+m^2}z/a)}{\sinh(\pi\sqrt{n^2+m^2})}.$$

Consider the superposition of *six* such cubes, one with  $V_0$  on each of the six faces. The result is a cube with  $V_0$  on its entire surface, so the potential at the center is  $V_0$ . Evidently the potential at the center of the original cube (with  $V_0$  on just one face) is one sixth of this:  $V_0/6$ . To check it, put in  $x = y = z = a/2$ :

$$V(a/2, a/2, a/2) = \frac{16V_0}{\pi^2} \sum_{n=1,3,5,\dots} \sum_{m=1,3,5,\dots} \frac{1}{nm} \sin(n\pi/2) \sin(m\pi/2) \frac{\sinh(\pi\sqrt{n^2+m^2}/2)}{\sinh(\pi\sqrt{n^2+m^2})}.$$

Let  $n \equiv 2i+1$ ,  $m \equiv 2j+1$ , and note that  $\sinh(2u) = 2 \sinh(u) \cosh(u)$ . The double sum is then

$$S = \frac{1}{2} \sum_{i=0}^{\infty} \sum_{j=0}^{\infty} \frac{(-1)^{i+j}}{(2i+1)(2j+1)} \operatorname{sech} \left[ \pi \sqrt{(2i+1)^2 + (2j+1)^2} / 2 \right].$$

Setting the upper limits at  $i = 3$ ,  $j = 3$  (or above) Mathematica returns  $S = 0.102808$ , which (to 6 digits) is equal to  $\pi^2/96$ , confirming (at least, numerically) that  $V(a/2, a/2, a/2) = V_0/6$ .

### Problem 3.17

$$\begin{aligned} P_3(x) &= \frac{1}{8 \cdot 6} \frac{d^3}{dx^3} (x^2 - 1)^3 = \frac{1}{48} \frac{d^2}{dx^2} 3(x^2 - 1)^2 2x = \frac{1}{8} \frac{d^2}{dx^2} x(x^2 - 1)^2 \\ &= \frac{1}{8} \frac{d}{dx} [(x^2 - 1)^2 + 2x(x^2 - 1)2x] = \frac{1}{8} \frac{d}{dx} [(x^2 - 1)(x^2 - 1 + 4x^2)] \\ &= \frac{1}{8} \frac{d}{dx} [(x^2 - 1)(5x^2 - 1)] = \frac{1}{8} [2x(5x^2 - 1) + (x^2 - 1)10x] \\ &= \frac{1}{4} (5x^3 - x + 5x^3 - 5x) = \frac{1}{4} (10x^3 - 6x) = \boxed{\frac{5}{2}x^3 - \frac{3}{2}x}. \end{aligned}$$

We need to show that  $P_3(\cos \theta)$  satisfies

$$\frac{1}{\sin \theta} \frac{d}{d\theta} \left( \sin \theta \frac{dP}{d\theta} \right) = -l(l+1)P, \text{ with } l = 3,$$

where  $P_3(\cos \theta) = \frac{1}{2} \cos \theta (5 \cos^2 \theta - 3)$ .

$$\begin{aligned} \frac{dP_3}{d\theta} &= \frac{1}{2} [-\sin \theta (5 \cos^2 \theta - 3) + \cos \theta (10 \cos \theta (-\sin \theta))] = -\frac{1}{2} \sin \theta (5 \cos^2 \theta - 3 + 10 \cos^2 \theta) \\ &= -\frac{3}{2} \sin \theta (5 \cos^2 \theta - 1). \end{aligned}$$

$$\begin{aligned}\frac{\partial}{\partial \theta} \left( \sin \theta \frac{dP_3}{d\theta} \right) &= -\frac{3}{2} \frac{d}{d\theta} [\sin^2 \theta (5 \cos^2 \theta - 1)] = -\frac{3}{2} [2 \sin \theta \cos \theta (5 \cos^2 \theta - 1) + \sin^2 \theta (-10 \cos \theta \sin \theta)] \\ &= -3 \sin \theta \cos \theta [5 \cos^2 \theta - 1 - 5 \sin^2 \theta].\end{aligned}$$

$$\begin{aligned}\frac{1}{\sin \theta} \frac{d}{d\theta} \left( \sin \theta \frac{dP}{d\theta} \right) &= -3 \cos \theta [5 \cos^2 - 1 - 5 (1 - \cos^2 \theta)] = -3 \cos \theta (10 \cos^2 \theta - 6) \\ &= -3 \cdot 4 \cdot \frac{1}{2} \cos \theta (5 \cos^2 \theta - 3) = -l(l+1)P_3. \quad \text{qed}\end{aligned}$$

$$\int_{-1}^1 P_1(x) P_3(x) dx = \int_{-1}^1 (x) \frac{1}{2} (5x^3 - 3x) dx = \frac{1}{2} (x^5 - x^3) \Big|_{-1}^1 = \frac{1}{2} (1 - 1 + 1 - 1) = 0. \quad \checkmark$$

**Problem 3.18**

(a) *Inside:*  $V(r, \theta) = \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta)$  (Eq. 3.66) where

$$A_l = \frac{(2l+1)}{2R^l} \int_0^{\pi} V_0(\theta) P_l(\cos \theta) \sin \theta d\theta \quad (\text{Eq. 3.69}).$$

In this case  $V_0(\theta) = V_0$  comes outside the integral, so

$$A_l = \frac{(2l+1)V_0}{2R^l} \int_0^{\pi} P_l(\cos \theta) \sin \theta d\theta.$$

But  $P_0(\cos \theta) = 1$ , so the integral can be written

$$\int_0^{\pi} P_0(\cos \theta) P_l(\cos \theta) \sin \theta d\theta = \begin{cases} 0, & \text{if } l \neq 0 \\ 2, & \text{if } l = 0 \end{cases} \quad (\text{Eq. 3.68}).$$

Therefore

$$A_l = \begin{cases} 0, & \text{if } l \neq 0 \\ V_0, & \text{if } l = 0 \end{cases}.$$

Plugging this into the general form:

$$V(r, \theta) = A_0 r^0 P_0(\cos \theta) = \boxed{V_0}.$$

The potential is *constant throughout the sphere*.

*Outside:*  $V(r, \theta) = \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta)$  (Eq. 3.72), where

$$\begin{aligned}B_l &= \frac{(2l+1)}{2} R^{l+1} \int_0^{\pi} V_0(\theta) P_l(\cos \theta) \sin \theta d\theta \quad (\text{Eq. 3.73}). \\ &= \frac{(2l+1)}{2} R^{l+1} V_0 \int_0^{\pi} P_l(\cos \theta) \sin \theta d\theta = \begin{cases} 0, & \text{if } l \neq 0 \\ RV_0, & \text{if } l = 0 \end{cases}.\end{aligned}$$

Therefore  $V(r, \theta) = V_0 \frac{R}{r}$  (i.e. equals  $V_0$  at  $r = R$ , then falls off like  $\frac{1}{r}$ ).

(b)

$$V(r, \theta) = \begin{cases} \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta), & \text{for } r \leq R \text{ (Eq. 3.78)} \\ \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta), & \text{for } r \geq R \text{ (Eq. 3.79)} \end{cases},$$

where

$$B_l = R^{2l+1} A_l \quad (\text{Eq. 3.81})$$

and

$$\begin{aligned} A_l &= \frac{1}{2\epsilon_0 R^{l-1}} \int_0^\pi \sigma_0(\theta) P_l(\cos \theta) \sin \theta d\theta \quad (\text{Eq. 3.84}) \\ &= \frac{1}{2\epsilon_0 R^{l-1}} \sigma_0 \int_0^\pi P_l(\cos \theta) \sin \theta d\theta = \begin{cases} 0, & \text{if } l \neq 0 \\ R\sigma_0/\epsilon_0, & \text{if } l = 0 \end{cases}. \end{aligned}$$

Therefore

$$V(r, \theta) = \begin{cases} \frac{R\sigma_0}{\epsilon_0}, & \text{for } r \leq R \\ \frac{R^2\sigma_0}{\epsilon_0} \frac{1}{r}, & \text{for } r \geq R \end{cases}.$$

Note: in terms of the total charge  $Q = 4\pi R^2 \sigma_0$ ,

$$V(r, \theta) = \begin{cases} \frac{1}{4\pi\epsilon_0} \frac{Q}{R}, & \text{for } r \leq R \\ \frac{1}{4\pi\epsilon_0} \frac{Q}{r}, & \text{for } r \geq R \end{cases}.$$

### Problem 3.19

$$V_0(\theta) = k \cos(3\theta) = k [4 \cos^3 \theta - 3 \cos \theta] = k [\alpha P_3(\cos \theta) + \beta P_1(\cos \theta)].$$

(I know that any 3<sup>rd</sup> order polynomial can be expressed as a linear combination of the first four Legendre polynomials; in this case, since the polynomial is *odd*, I only need  $P_1$  and  $P_3$ .)

$$4 \cos^3 \theta - 3 \cos \theta = \alpha \left[ \frac{1}{2} (5 \cos^3 \theta - 3 \cos \theta) \right] + \beta \cos \theta = \frac{5\alpha}{2} \cos^3 \theta + \left( \beta - \frac{3}{2}\alpha \right) \cos \theta,$$

so

$$4 = \frac{5\alpha}{2} \Rightarrow \alpha = \frac{8}{5}; \quad -3 = \beta - \frac{3}{2}\alpha = \beta - \frac{3}{2} \cdot \frac{8}{5} = \beta - \frac{12}{5} \Rightarrow \beta = \frac{12}{5} - 3 = -\frac{3}{5}.$$

Therefore

$$V_0(\theta) = \frac{k}{5} [8P_3(\cos \theta) - 3P_1(\cos \theta)].$$

Now

$$V(r, \theta) = \left\{ \begin{array}{l} \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta), \text{ for } r \leq R \text{ (Eq. 3.66)} \\ \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta), \text{ for } r \geq R \text{ (Eq. 3.72)} \end{array} \right\},$$

where

$$\begin{aligned} A_l &= \frac{(2l+1)}{2R^l} \int_0^\pi V_0(\theta) P_l(\cos \theta) \sin \theta d\theta \quad (\text{Eq. 3.69}) \\ &= \frac{(2l+1)}{2R^l} \frac{k}{5} \left\{ 8 \int_0^\pi P_3(\cos \theta) P_l(\cos \theta) \sin \theta d\theta - 3 \int_0^\pi P_1(\cos \theta) P_l(\cos \theta) \sin \theta d\theta \right\} \\ &= \frac{k}{5} \frac{(2l+1)}{2R^l} \left\{ 8 \frac{2}{(2l+1)} \delta_{l3} - 3 \frac{2}{(2l+1)} \delta_{l1} \right\} = \frac{k}{5} \frac{1}{R^l} [8 \delta_{l3} - 3 \delta_{l1}] \\ &= \begin{cases} 8k/5R^3, & \text{if } l = 3 \\ -3k/5R, & \text{if } l = 1 \end{cases} \text{ (zero otherwise).} \end{aligned}$$

Therefore

$$V(r, \theta) = -\frac{3k}{5R} r P_1(\cos \theta) + \frac{8k}{5R^3} r^3 P_3(\cos \theta) = \boxed{\frac{k}{5} \left[ 8 \left( \frac{r}{R} \right)^3 P_3(\cos \theta) - 3 \left( \frac{r}{R} \right) P_1(\cos \theta) \right]},$$

or

$$\frac{k}{5} \left\{ 8 \left( \frac{r}{R} \right)^3 \frac{1}{2} [5 \cos^3 \theta - 3 \cos \theta] - 3 \left( \frac{r}{R} \right) \cos \theta \right\} \Rightarrow \boxed{V(r, \theta) = \frac{k}{5} \frac{r}{R} \cos \theta \left\{ 4 \left( \frac{r}{R} \right)^2 [5 \cos^2 \theta - 3] - 3 \right\}}$$

(for  $r \leq R$ ). Meanwhile,  $B_l = A_l R^{2l+1}$  (Eq. 3.81—this follows from the continuity of  $V$  at  $R$ ). Therefore

$$B_l = \begin{cases} 8kR^4/5, & \text{if } l = 3 \\ -3kR^2/5, & \text{if } l = 1 \end{cases} \text{ (zero otherwise).}$$

So

$$V(r, \theta) = \frac{-3kR^2}{5} \frac{1}{r^2} P_1(\cos \theta) + \frac{8kR^4}{5} \frac{1}{r^4} P_3(\cos \theta) = \boxed{\frac{k}{5} \left[ 8 \left( \frac{R}{r} \right)^4 P_3(\cos \theta) - 3 \left( \frac{R}{r} \right)^2 P_1(\cos \theta) \right]},$$

or

$$\boxed{V(r, \theta) = \frac{k}{5} \left( \frac{R}{r} \right)^2 \cos \theta \left\{ 4 \left( \frac{R}{r} \right)^2 [5 \cos^2 \theta - 3] - 3 \right\}}$$

(for  $r \geq R$ ). Finally, using Eq. 3.83:

$$\begin{aligned}
 \sigma(\theta) &= \epsilon_0 \sum_{l=0}^{\infty} (2l+1) A_l R^{l-1} P_l(\cos \theta) = \epsilon_0 [3A_1 P_1 + 7A_3 R^2 P_3] \\
 &= \epsilon_0 \left[ 3 \left( -\frac{3k}{5R} \right) P_1 + 7 \left( \frac{8k}{5R^3} \right) R^2 P_3 \right] = \boxed{\frac{\epsilon_0 k}{5R} [-9P_1(\cos \theta) + 56P_3(\cos \theta)]} \\
 &= \frac{\epsilon_0 k}{5R} \left[ -9 \cos \theta + \frac{56}{2} (5 \cos^3 \theta - 3 \cos \theta) \right] = \frac{\epsilon_0 k}{5R} \cos \theta [-9 + 28 \cdot 5 \cos^2 \theta - 28 \cdot 3] \\
 &= \boxed{\frac{\epsilon_0 k}{5R} \cos \theta [140 \cos^2 \theta - 93]}.
 \end{aligned}$$

### Problem 3.20

Use Eq. 3.83:  $\sigma(\theta) = \epsilon_0 \sum_{l=0}^{\infty} (2l+1) A_l R^{l-1} P_l(\cos \theta)$ . But Eq. 3.69 says:  $A_l = \frac{2l+1}{2R^l} \int_0^\pi V_0(\theta) P_l(\cos \theta) \sin \theta d\theta$ .

Putting them together:

$$\sigma(\theta) = \frac{\epsilon_0}{2R} \sum_{l=0}^{\infty} (2l+1)^2 C_l P_l(\cos \theta), \quad \text{with } C_l = \int_0^\pi V_0(\theta) P_l(\cos \theta) \sin \theta d\theta. \quad \text{qed}$$

### Problem 3.21

Set  $V = 0$  on the equatorial plane, far from the sphere. Then the potential is the same as Ex. 3.8 *plus* the potential of a uniformly charged spherical shell:

$$\boxed{V(r, \theta) = -E_0 \left( r - \frac{R^3}{r^2} \right) \cos \theta + \frac{1}{4\pi\epsilon_0} \frac{Q}{r}.$$

### Problem 3.22

$$(a) \quad V(r, \theta) = \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta) \quad (r > R), \quad \text{so } V(r, 0) = \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(1) = \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} = \frac{\sigma}{2\epsilon_0} [\sqrt{r^2 + R^2} - r].$$

Since  $r > R$  in this region,  $\sqrt{r^2 + R^2} = r\sqrt{1 + (R/r)^2} = r \left[ 1 + \frac{1}{2}(R/r)^2 - \frac{1}{8}(R/r)^4 + \dots \right]$ , so

$$\sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} = \frac{\sigma}{2\epsilon_0} r \left[ 1 + \frac{1}{2} \frac{R^2}{r^2} - \frac{1}{8} \frac{R^4}{r^4} + \dots - 1 \right] = \frac{\sigma}{2\epsilon_0} \left( \frac{R^2}{2r} - \frac{R^4}{8r^3} + \dots \right).$$

Comparing like powers of  $r$ , I see that  $B_0 = \frac{\sigma R^2}{4\epsilon_0}$ ,  $B_1 = 0$ ,  $B_2 = -\frac{\sigma R^4}{16\epsilon_0}$ ,  $\dots$ . Therefore

$$\begin{aligned}
 V(r, \theta) &= \frac{\sigma R^2}{4\epsilon_0} \left[ \frac{1}{r} - \frac{R^2}{4r^3} P_2(\cos \theta) + \dots \right], \\
 &= \frac{\sigma R^2}{4\epsilon_0 r} \left[ 1 - \frac{1}{8} \left( \frac{R}{r} \right)^2 (3 \cos^2 \theta - 1) + \dots \right],
 \end{aligned} \quad (\text{for } r > R).$$

(b)  $V(r, \theta) = \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta)$  ( $r < R$ ). In the northern hemisphere,  $0 \leq \theta \leq \pi/2$ ,

$$V(r, 0) = \sum_{l=0}^{\infty} A_l r^l = \frac{\sigma}{2\epsilon_0} \left[ \sqrt{r^2 + R^2} - r \right].$$

Since  $r < R$  in this region,  $\sqrt{r^2 + R^2} = R\sqrt{1 + (r/R)^2} = R \left[ 1 + \frac{1}{2}(r/R)^2 - \frac{1}{8}(r/R)^4 + \dots \right]$ . Therefore

$$\sum_{l=0}^{\infty} A_l r^l = \frac{\sigma}{2\epsilon_0} \left[ R + \frac{1}{2} \frac{r^2}{R} - \frac{1}{8} \frac{r^4}{R^3} + \dots - r \right].$$

Comparing like powers:  $A_0 = \frac{\sigma}{2\epsilon_0} R$ ,  $A_1 = -\frac{\sigma}{2\epsilon_0}$ ,  $A_2 = \frac{\sigma}{4\epsilon_0 R}$ ,  $\dots$ , so

$$\begin{aligned} V(r, \theta) &= \frac{\sigma}{2\epsilon_0} \left[ R - r P_1(\cos \theta) + \frac{1}{2R} r^2 P_2(\cos \theta) + \dots \right], \\ &= \frac{\sigma R}{2\epsilon_0} \left[ 1 - \left( \frac{r}{R} \right) \cos \theta + \frac{1}{4} \left( \frac{r}{R} \right)^2 (3 \cos^2 \theta - 1) + \dots \right], \end{aligned} \quad (\text{for } r < R, \text{ northern hemisphere}).$$

In the southern hemisphere we'll have to go for  $\theta = \pi$ , using  $P_l(-1) = (-1)^l$ .

$$V(r, \pi) = \sum_{l=0}^{\infty} (-1)^l \bar{A}_l r^l = \frac{\sigma}{2\epsilon_0} \left[ \sqrt{r^2 + R^2} - r \right].$$

(I put an overbar on  $\bar{A}_l$  to distinguish it from the northern  $A_l$ ). The only difference is the sign of  $\bar{A}_1$ :  $\bar{A}_1 = +(\sigma/2\epsilon_0)$ ,  $\bar{A}_0 = A_0$ ,  $\bar{A}_2 = A_2$ . So:

$$\begin{aligned} V(r, \theta) &= \frac{\sigma}{2\epsilon_0} \left[ R + r P_1(\cos \theta) + \frac{1}{2R} r^2 P_2(\cos \theta) + \dots \right], \\ &= \frac{\sigma R}{2\epsilon_0} \left[ 1 + \left( \frac{r}{R} \right) \cos \theta + \frac{1}{4} \left( \frac{r}{R} \right)^2 (3 \cos^2 \theta - 1) + \dots \right], \end{aligned} \quad (\text{for } r < R, \text{ southern hemisphere}).$$

---

### Problem 3.23

$$V(r, \theta) = \begin{cases} \sum_{l=0}^{\infty} A_l r^l P_l(\cos \theta), & (r \leq R) \text{ (Eq. 3.78),} \\ \sum_{l=0}^{\infty} \frac{B_l}{r^{l+1}} P_l(\cos \theta), & (r \geq R) \text{ (Eq. 3.79),} \end{cases}$$



where  $B_l = A_l R^{2l+1}$  (Eq. 3.81) and

$$\begin{aligned} A_l &= \frac{1}{2\epsilon_0 R^{l-1}} \int_0^\pi \sigma_0(\theta) P_l(\cos \theta) \sin \theta d\theta \quad (\text{Eq. 3.84}) \\ &= \frac{1}{2\epsilon_0 R^{l-1}} \sigma_0 \left\{ \int_0^{\pi/2} P_l(\cos \theta) \sin \theta d\theta - \int_{\pi/2}^\pi P_l(\cos \theta) \sin \theta d\theta \right\} \quad (\text{let } x = \cos \theta) \\ &= \frac{\sigma_0}{2\epsilon_0 R^{l-1}} \left\{ \int_0^1 P_l(x) dx - \int_{-1}^0 P_l(x) dx \right\}. \end{aligned}$$

Now  $P_l(-x) = (-1)^l P_l(x)$ , since  $P_l(x)$  is even, for even  $l$ , and odd, for odd  $l$ . Therefore

$$\int_{-1}^0 P_l(x) dx = \int_1^0 P_l(-x) d(-x) = (-1)^l \int_0^1 P_l(x) dx,$$

and hence

$$A_l = \frac{\sigma_0}{2\epsilon_0 R^{l-1}} [1 - (-1)^l] \int_0^1 P_l(x) dx = \begin{cases} 0, & \text{if } l \text{ is even} \\ \frac{\sigma_0}{\epsilon_0 R^{l-1}} \int_0^1 P_l(x) dx, & \text{if } l \text{ is odd} \end{cases}.$$

So  $A_0 = A_2 = A_4 = A_6 = 0$ , and all we need are  $A_1$ ,  $A_3$ , and  $A_5$ .

$$\begin{aligned} \int_0^1 P_1(x) dx &= \int_0^1 x dx = \frac{x^2}{2} \Big|_0^1 = \frac{1}{2}. \\ \int_0^1 P_3(x) dx &= \frac{1}{2} \int_0^1 (5x^3 - 3x) dx = \frac{1}{2} \left( 5\frac{x^4}{4} - 3\frac{x^2}{2} \right) \Big|_0^1 = \frac{1}{2} \left( \frac{5}{4} - \frac{3}{2} \right) = -\frac{1}{8}. \\ \int_0^1 P_5(x) dx &= \frac{1}{8} \int_0^1 (63x^5 - 70x^3 + 15x) dx = \frac{1}{8} \left( 63\frac{x^6}{6} - 70\frac{x^4}{4} + 15\frac{x^2}{2} \right) \Big|_0^1 \\ &= \frac{1}{8} \left( \frac{21}{2} - \frac{35}{2} + \frac{15}{2} \right) = \frac{1}{16} (36 - 35) = \frac{1}{16}. \end{aligned}$$

Therefore

$$A_1 = \frac{\sigma_0}{\epsilon_0} \left( \frac{1}{2} \right); \quad A_3 = \frac{\sigma_0}{\epsilon_0 R^2} \left( -\frac{1}{8} \right); \quad A_5 = \frac{\sigma_0}{\epsilon_0 R^4} \left( \frac{1}{16} \right); \quad \text{etc.}$$

and

$$B_1 = \frac{\sigma_0}{\epsilon_0} R^3 \left( \frac{1}{2} \right); \quad B_3 = \frac{\sigma_0}{\epsilon_0} R^5 \left( -\frac{1}{8} \right); \quad B_5 = \frac{\sigma_0}{\epsilon_0} R^7 \left( \frac{1}{16} \right); \quad \text{etc.}$$

Thus

$$V(r, \theta) = \begin{cases} \frac{\sigma_0 r}{2\epsilon_0} \left[ P_1(\cos \theta) - \frac{1}{4} \left( \frac{r}{R} \right)^2 P_3(\cos \theta) + \frac{1}{8} \left( \frac{r}{R} \right)^4 P_5(\cos \theta) + \dots \right], & (r \leq R), \\ \frac{\sigma_0 R^3}{2\epsilon_0 r^2} \left[ P_1(\cos \theta) - \frac{1}{4} \left( \frac{R}{r} \right)^2 P_3(\cos \theta) + \frac{1}{8} \left( \frac{R}{r} \right)^4 P_5(\cos \theta) + \dots \right], & (r \geq R). \end{cases}$$

### Problem 3.24

$$\frac{1}{s} \frac{\partial}{\partial s} \left( s \frac{\partial V}{\partial s} \right) + \frac{1}{s^2} \frac{\partial^2 V}{\partial \phi^2} = 0.$$

Look for solutions of the form  $V(s, \phi) = S(s)\Phi(\phi)$ :

$$\frac{1}{s} \Phi \frac{d}{ds} \left( s \frac{dS}{ds} \right) + \frac{1}{s^2} S \frac{d^2 \Phi}{d\phi^2} = 0.$$

Multiply by  $s^2$  and divide by  $V = S\Phi$ :

$$\frac{s}{S} \frac{d}{ds} \left( s \frac{dS}{ds} \right) + \frac{1}{\Phi} \frac{d^2 \Phi}{d\phi^2} = 0.$$

Since the first term involves  $s$  only, and the second  $\phi$  only, each is a constant:

$$\frac{s}{S} \frac{d}{ds} \left( s \frac{dS}{ds} \right) = C_1, \quad \frac{1}{\Phi} \frac{d^2 \Phi}{d\phi^2} = C_2, \quad \text{with } C_1 + C_2 = 0.$$

Now  $C_2$  must be negative (else we get exponentials for  $\Phi$ , which do not return to their original value—as geometrically they *must*—when  $\phi$  is increased by  $2\pi$ ).

$$C_2 = -k^2. \quad \text{Then } \frac{d^2 \Phi}{d\phi^2} = -k^2 \Phi \Rightarrow \Phi = A \cos k\phi + B \sin k\phi.$$

Moreover, since  $\Phi(\phi + 2\pi) = \Phi(\phi)$ ,  $k$  must be an integer:  $k = 0, 1, 2, 3, \dots$  (negative integers are just repeats, but  $k = 0$  must be included, since  $\Phi = A$  (a constant) is OK).

$s \frac{d}{ds} \left( s \frac{dS}{ds} \right) = k^2 S$  can be solved by  $S = s^n$ , provided  $n$  is chosen right:

$$s \frac{d}{ds} (s n s^{n-1}) = n s \frac{d}{ds} (s^n) = n^2 s s^{n-1} = n^2 s^n = k^2 S \Rightarrow n = \pm k.$$

Evidently the general solution is  $S(s) = C s^k + D s^{-k}$ , unless  $k = 0$ , in which case we have only *one* solution to a *second-order* equation—namely,  $S = \text{constant}$ . So we must treat  $k = 0$  separately. *One* solution is a constant—but what's the other? Go back to the differential equation for  $S$ , and put in  $k = 0$ :

$$s \frac{d}{ds} \left( s \frac{dS}{ds} \right) = 0 \Rightarrow s \frac{dS}{ds} = \text{constant} = C \Rightarrow \frac{dS}{ds} = \frac{C}{s} \Rightarrow dS = C \frac{ds}{s} \Rightarrow S = C \ln s + D \text{ (another constant)}.$$

So the second solution in this case is  $\ln s$ . [How about  $\Phi$ ? That *too* reduces to a single solution,  $\Phi = A$ , in the case  $k = 0$ . What's the second solution here? Well, putting  $k = 0$  into the  $\Phi$  equation:

$$\frac{d^2 \Phi}{d\phi^2} = 0 \Rightarrow \frac{d\Phi}{d\phi} = \text{constant} = B \Rightarrow \Phi = B\phi + A.$$

But a term of the form  $B\phi$  is unacceptable, since it does not return to its initial value when  $\phi$  is augmented by  $2\pi$ .] *Conclusion:* The general solution with cylindrical symmetry is


$$V(s, \phi) = a_0 + b_0 \ln s + \sum_{k=1}^{\infty} [s^k (a_k \cos k\phi + b_k \sin k\phi) + s^{-k} (c_k \cos k\phi + d_k \sin k\phi)].$$

Yes: the potential of a line charge goes like  $\ln s$ , which *is* included.

### Problem 3.25

Picking  $V = 0$  on the  $yz$  plane, with  $\mathbf{E}_0$  in the  $x$  direction, we have (Eq. 3.74):

$$\left\{ \begin{array}{ll} \text{(i)} & V = 0, \\ \text{(ii)} & V \rightarrow -E_0 x = -E_0 s \cos \phi, \text{ for } s \gg R. \end{array} \right\} \quad \text{when } s = R,$$



Evidently  $a_0 = b_0 = b_k = d_k = 0$ , and  $a_k = c_k = 0$  except for  $k = 1$ :

$$V(s, \phi) = \left( a_1 s + \frac{c_1}{s} \right) \cos \phi.$$

(i)  $\Rightarrow c_1 = -a_1 R^2$ ; (ii)  $\rightarrow a_1 = -E_0$ . Therefore

$$V(s, \phi) = \left( -E_0 s + \frac{E_0 R^2}{s} \right) \cos \phi, \quad \text{or} \quad V(s, \phi) = E_0 s \left[ \left( \frac{R}{s} \right)^2 - 1 \right] \cos \phi.$$

$$\sigma = -\epsilon_0 \left. \frac{\partial V}{\partial s} \right|_{s=R} = -\epsilon_0 E_0 \left( -\frac{R^2}{s^2} - 1 \right) \cos \phi \Big|_{s=R} = 2\epsilon_0 E_0 \cos \phi.$$

### Problem 3.26

*Inside:*  $V(s, \phi) = a_0 + \sum_{k=1}^{\infty} s^k (a_k \cos k\phi + b_k \sin k\phi)$ . (In this region  $\ln s$  and  $s^{-k}$  are no good—they blow up at  $s = 0$ .)

*Outside:*  $V(s, \phi) = \bar{a}_0 + \sum_{k=1}^{\infty} \frac{1}{s^k} (c_k \cos k\phi + d_k \sin k\phi)$ . (Here  $\ln s$  and  $s^k$  are no good at  $s \rightarrow \infty$ ).

$$\sigma = -\epsilon_0 \left( \frac{\partial V_{\text{out}}}{\partial s} - \frac{\partial V_{\text{in}}}{\partial s} \right) \Big|_{s=R} \quad (\text{Eq. 2.36}).$$

Thus

$$a \sin 5\phi = -\epsilon_0 \sum_{k=1}^{\infty} \left\{ -\frac{k}{R^{k+1}} (c_k \cos k\phi + d_k \sin k\phi) - k R^{k-1} (a_k \cos k\phi + b_k \sin k\phi) \right\}.$$

Evidently  $a_k = c_k = 0$ ;  $b_k = d_k = 0$  except  $k = 5$ ;  $a = 5\epsilon_0 \left( \frac{1}{R^6} d_5 + R^4 b_5 \right)$ . Also,  $V$  is continuous at  $s = R$ :

$a_0 + R^5 b_5 \sin 5\phi = \bar{a}_0 + \frac{1}{R^5} d_5 \sin 5\phi$ . So  $a_0 = \bar{a}_0$  (might as well choose both zero);  $R^5 b_5 = R^{-5} d_5$ , or  $d_5 = R^{10} b_5$ .

Combining these results:  $a = 5\epsilon_0 (R^4 b_5 + R^4 b_5) = 10\epsilon_0 R^4 b_5$ ;  $b_5 = \frac{a}{10\epsilon_0 R^4}$ ;  $d_5 = \frac{aR^6}{10\epsilon_0}$ . Therefore

$$V(s, \phi) = \frac{a \sin 5\phi}{10\epsilon_0} \left\{ \begin{array}{l} s^5/R^4, \text{ for } s < R, \\ R^6/s^5, \text{ for } s > R. \end{array} \right\}$$

**Problem 3.27** Since  $\mathbf{r}$  is on the  $z$  axis, the angle  $\alpha$  is just the polar angle  $\theta$  (I'll drop the primes, for simplicity).

*Monopole term:*

$$\int \rho d\tau = kR \int \left[ \frac{1}{r^2} (R - 2r) \sin \theta \right] r^2 \sin \theta dr d\theta d\phi.$$

But the  $r$  integral is

$$\int_0^R (R - 2r) dr = (Rr - r^2)|_0^R = R^2 - R^2 = 0.$$

So the monopole term is zero.

*Dipole term:*

$$\int r \cos \theta \rho d\tau = kR \int (r \cos \theta) \left[ \frac{1}{r^2} (R - 2r) \sin \theta \right] r^2 \sin \theta dr d\theta d\phi.$$

But the  $\theta$  integral is

$$\int_0^\pi \sin^2 \theta \cos \theta d\theta = \left. \frac{\sin^3 \theta}{3} \right|_0^\pi = \frac{1}{3}(0 - 0) = 0.$$

So the dipole contribution is likewise zero.

*Quadrupole term:*

$$\int r^2 \left( \frac{3}{2} \cos^2 \theta - \frac{1}{2} \right) \rho d\tau = \frac{1}{2} kR \int r^2 (3 \cos^2 \theta - 1) \left[ \frac{1}{r^2} (R - 2r) \sin \theta \right] r^2 \sin \theta dr d\theta d\phi.$$

*$r$  integral:*

$$\int_0^R r^2 (R - 2r) dr = \left( \frac{r^3}{3} R - \frac{r^4}{2} \right) \Big|_0^R = \frac{R^4}{3} - \frac{R^4}{2} = -\frac{R^4}{6}.$$

*$\theta$  integral:*

$$\begin{aligned} \int_0^\pi \underbrace{(3 \cos^2 \theta - 1)}_{3(1 - \sin^2 \theta) - 1 = 2 - 3 \sin^2 \theta} \sin^2 \theta d\theta &= 2 \int_0^\pi \sin^2 \theta d\theta - 3 \int_0^\pi \sin^4 \theta d\theta \\ &= 2 \left( \frac{\pi}{2} \right) - 3 \left( \frac{3\pi}{8} \right) = \pi \left( 1 - \frac{9}{8} \right) = -\frac{\pi}{8}. \end{aligned}$$

*$\phi$  integral:*

$$\int_0^{2\pi} d\phi = 2\pi.$$


The whole integral is:

$$\frac{1}{2} kR \left( -\frac{R^4}{6} \right) \left( -\frac{\pi}{8} \right) (2\pi) = \frac{k\pi^2 R^5}{48}.$$

For point  $P$  on the  $z$  axis ( $r \rightarrow z$  in Eq. 3.95) the approximate potential is

$$V(z) \cong \frac{1}{4\pi\epsilon_0} \frac{k\pi^2 R^5}{48z^3}. \quad (\text{Quadrupole.})$$

### Problem 3.28



For a line charge,  $\rho(\mathbf{r}') d\tau' \rightarrow \lambda(\mathbf{r}') dl'$ , which in this case becomes  $\lambda R d\phi'$ .

$$\begin{aligned} \mathbf{r} &= r \sin \theta \cos \phi \hat{\mathbf{x}} + r \sin \theta \sin \phi \hat{\mathbf{y}} + r \cos \theta \hat{\mathbf{z}}, \\ \mathbf{r}' &= R \cos \phi' + R \sin \phi' \hat{\mathbf{y}}, \quad \text{so} \\ \mathbf{r} \cdot \mathbf{r}' &= r R \sin \theta \cos \phi \cos \phi' + r R \sin \theta \sin \phi \sin \phi' = r R \cos \alpha, \\ \cos \alpha &= \sin \theta (\cos \phi \cos \phi' + \sin \phi \sin \phi'). \end{aligned}$$

$n = 0$  :

$$\int \rho(\mathbf{r}') d\tau' \rightarrow \lambda R \int_0^{2\pi} d\phi' = 2\pi R \lambda; \quad V_0 = \frac{1}{4\pi\epsilon_0} \frac{2\pi R \lambda}{r} = \boxed{\frac{\lambda}{2\epsilon_0} \frac{R}{r}}.$$

$n = 1$  :

$$\int r' \cos \alpha \rho(\mathbf{r}') d\tau' \rightarrow \int R \cos \alpha \lambda R d\phi' = \lambda R^2 \sin \theta \int_0^{2\pi} (\cos \phi \cos \phi' + \sin \phi \sin \phi') d\phi' = 0; \quad V_1 = \boxed{0}.$$

$n = 2$  :

$$\begin{aligned} \int (r')^2 P_2(\cos \alpha) \rho(\mathbf{r}') d\tau' &\rightarrow \int R^2 \left( \frac{3}{2} \cos^2 \alpha - \frac{1}{2} \right) \lambda R d\phi' = \frac{\lambda R^3}{2} \int \left[ 3 \sin^2 \theta (\cos \phi \cos \phi' + \sin \phi \sin \phi')^2 - 1 \right] d\phi' \\ &= \frac{\lambda R^3}{2} \left[ 3 \sin^2 \theta \left( \cos^2 \phi \int_0^{2\pi} \cos^2 \phi' d\phi' + \sin^2 \phi \int_0^{2\pi} \sin^2 \phi' d\phi' + 2 \sin \phi \cos \phi \int_0^{2\pi} \sin \phi' \cos \phi' d\phi' \right) - \int_0^{2\pi} d\phi' \right] \\ &= \frac{\lambda R^3}{2} [3 \sin^2 \theta (\pi \cos^2 \phi + \pi \sin^2 \phi + 0) - 2\pi] = \frac{\pi \lambda R^3}{2} (3 \sin^2 \theta - 2) = -\pi \lambda R^3 \left( \frac{3}{2} \cos^2 \theta - \frac{1}{2} \right). \end{aligned}$$

So

$$V_2 = \boxed{-\frac{\lambda}{8\epsilon_0} \frac{R^3}{r^3} (3 \cos^2 \theta - 1) = -\frac{\lambda}{4\epsilon_0} \frac{R^3}{r^3} P_2(\cos \theta)}.$$

**Problem 3.29**

$\mathbf{p} = (3qa - qa)\hat{\mathbf{z}} + (-2qa - 2q(-a))\hat{\mathbf{y}} = 2qa\hat{\mathbf{z}}$ . Therefore

$$V \cong \frac{1}{4\pi\epsilon_0} \frac{\mathbf{p} \cdot \hat{\mathbf{r}}}{r^2},$$

and  $\mathbf{p} \cdot \hat{\mathbf{r}} = 2qa\hat{\mathbf{z}} \cdot \hat{\mathbf{r}} = 2qa \cos \theta$ , so

$$V \cong \frac{1}{4\pi\epsilon_0} \frac{2qa \cos \theta}{r^2}. \quad (\text{Dipole.})$$

**Problem 3.30**

(a) By symmetry,  $\mathbf{p}$  is clearly in the  $z$  direction:  $\mathbf{p} = p\hat{\mathbf{z}}$ ;  $p = \int z\rho d\tau \Rightarrow \int z\sigma da$ .

$$\begin{aligned} p &= \int (R \cos \theta)(k \cos \theta) R^2 \sin \theta d\theta d\phi = 2\pi R^3 k \int_0^\pi \cos^2 \theta \sin \theta d\theta = 2\pi R^3 k \left( -\frac{\cos^3 \theta}{3} \right) \Big|_0^\pi \\ &= \frac{2}{3} \pi R^3 k [1 - (-1)] = \frac{4\pi R^3 k}{3}; \quad \boxed{\mathbf{p} = \frac{4\pi R^3 k}{3} \hat{\mathbf{z}}.} \end{aligned}$$

(b)

$$V \cong \frac{1}{4\pi\epsilon_0} \frac{4\pi R^3 k \cos \theta}{3 r^2} = \boxed{\frac{k R^3 \cos \theta}{3\epsilon_0 r^2}}. \quad (\text{Dipole.})$$

This is *also* the *exact* potential. *Conclusion:* all multiple moments of this distribution (except the dipole) are exactly zero.

**Problem 3.31**

Using Eq. 3.94 with  $r' = d/2$  and  $\alpha = \theta$  (Fig. 3.26):

$$\frac{1}{z_+} = \frac{1}{r} \sum_{n=0}^{\infty} \left( \frac{d}{2r} \right)^n P_n(\cos \theta);$$

for  $z_-$ , we let  $\theta \rightarrow 180^\circ + \theta$ , so  $\cos \theta \rightarrow -\cos \theta$ :

$$\frac{1}{z_-} = \frac{1}{r} \sum_{n=0}^{\infty} \left( \frac{d}{2r} \right)^n P_n(-\cos \theta).$$

But  $P_n(-x) = (-1)^n P_n(x)$ , so

$$V = \frac{1}{4\pi\epsilon_0} q \left( \frac{1}{z_+} - \frac{1}{z_-} \right) = \frac{1}{4\pi\epsilon_0} q \frac{1}{r} \sum_{n=0}^{\infty} \left( \frac{d}{2r} \right)^n [P_n(\cos \theta) - P_n(-\cos \theta)] = \frac{2q}{4\pi\epsilon_0 r} \sum_{n=1,3,5,\dots} \left( \frac{d}{2r} \right)^n P_n(\cos \theta).$$

Therefore

$$\begin{aligned} V_{\text{dip}} &= \frac{2q}{4\pi\epsilon_0} \frac{1}{r} \frac{d}{2r} P_1(\cos \theta) = \frac{qd \cos \theta}{4\pi\epsilon_0 r^2}, \quad \text{while} \quad \boxed{V_{\text{quad}} = 0.} \\ V_{\text{oct}} &= \frac{2q}{4\pi\epsilon_0 r} \left( \frac{d}{2r} \right)^3 P_3(\cos \theta) = \frac{2q}{4\pi\epsilon_0} \frac{d^3}{8r^4} \frac{1}{2} (5 \cos^3 \theta - 3 \cos \theta) = \boxed{\frac{qd^3}{4\pi\epsilon_0} \frac{1}{8r^4} (5 \cos^3 \theta - 3 \cos \theta)}. \end{aligned}$$

**Problem 3.32**

$$\begin{aligned}
\text{(a) (i) } Q &= \boxed{2q}, \quad \text{(ii) } \mathbf{p} = \boxed{3qa \hat{\mathbf{z}}}, \quad \text{(iii) } V \cong \frac{1}{4\pi\epsilon_0} \left[ \frac{Q}{r} + \frac{\mathbf{p} \cdot \hat{\mathbf{r}}}{r^2} \right] = \boxed{\frac{1}{4\pi\epsilon_0} \left[ \frac{2q}{r} + \frac{3qa \cos \theta}{r^2} \right]}. \\
\text{(b) (i) } Q &= \boxed{2q}, \quad \text{(ii) } \mathbf{p} = \boxed{qa \hat{\mathbf{z}}}, \quad \text{(iii) } V \cong \boxed{\frac{1}{4\pi\epsilon_0} \left[ \frac{2q}{r} + \frac{qa \cos \theta}{r^2} \right]}. \\
\text{(c) (i) } Q &= \boxed{2q}, \quad \text{(ii) } \mathbf{p} = \boxed{3qa \hat{\mathbf{y}}}, \quad \text{(iii) } V \cong \boxed{\frac{1}{4\pi\epsilon_0} \left[ \frac{2q}{r} + \frac{3qa \sin \theta \sin \phi}{r^2} \right]} \quad (\text{from Eq. 1.64, } \hat{\mathbf{y}} \cdot \hat{\mathbf{r}} = \sin \theta \sin \phi).
\end{aligned}$$

**Problem 3.33**

$$\begin{aligned}
\text{(a) This point is at } r &= a, \theta = \frac{\pi}{2}, \phi = 0, \text{ so } \mathbf{E} = \frac{p}{4\pi\epsilon_0 a^3} \hat{\boldsymbol{\theta}} = \frac{p}{4\pi\epsilon_0 a^3} (-\hat{\mathbf{z}}); \mathbf{F} = q\mathbf{E} = \boxed{-\frac{pq}{4\pi\epsilon_0 a^3} \hat{\mathbf{z}}}. \\
\text{(b) Here } r &= a, \theta = 0, \text{ so } \mathbf{E} = \frac{p}{4\pi\epsilon_0 a^3} (2\hat{\mathbf{r}}) = \frac{2p}{4\pi\epsilon_0 a^3} \hat{\mathbf{z}}. \quad \boxed{\mathbf{F} = \frac{2pq}{4\pi\epsilon_0 a^3} \hat{\mathbf{z}}}. \\
\text{(c) } W &= q[V(0,0,a) - V(a,0,0)] = \frac{qp}{4\pi\epsilon_0 a^2} \left[ \cos(0) - \cos\left(\frac{\pi}{2}\right) \right] = \boxed{\frac{pq}{4\pi\epsilon_0 a^2}}.
\end{aligned}$$

**Problem 3.34**

$$Q = -q, \text{ so } V_{\text{mono}} = \frac{1}{4\pi\epsilon_0} \frac{-q}{r}; \quad \mathbf{p} = qa \hat{\mathbf{z}}, \quad \text{so } V_{\text{dip}} = \frac{1}{4\pi\epsilon_0} \frac{qa \cos \theta}{r^2}. \text{ Therefore}$$

$$\boxed{V(r, \theta) \cong \frac{q}{4\pi\epsilon_0} \left( -\frac{1}{r} + \frac{a \cos \theta}{r^2} \right)} \quad \boxed{\mathbf{E}(r, \theta) \cong \frac{q}{4\pi\epsilon_0} \left[ -\frac{1}{r^2} \hat{\mathbf{r}} + \frac{a}{r^3} (2 \cos \theta \hat{\mathbf{r}} + \sin \theta \hat{\boldsymbol{\theta}}) \right]}.$$

**Problem 3.35** The total charge is zero, so the dominant term is the dipole. We need the dipole moment of this configuration. It obviously points in the  $z$  direction, and for the southern hemisphere ( $\theta : \frac{\pi}{2} \rightarrow \pi$ )  $\rho$  switches sign but so does  $z$ , so

$$\begin{aligned}
p &= \int z \rho d\tau = 2\rho_0 \int_{\theta=0}^{\pi/2} r \cos \theta r^2 \sin \theta dr d\theta d\phi = 2\rho_0 (2\pi) \int_0^R r^3 dr \int_0^{\pi/2} \cos \theta \sin \theta d\theta \\
&= 4\pi\rho_0 \frac{R^4}{4} \frac{\sin^2 \theta}{2} \Big|_0^{\pi/2} = \frac{\pi\rho_0 R^4}{2}.
\end{aligned}$$

Therefore (Eq. 3.103)

$$\mathbf{E} \approx \boxed{\frac{\pi\rho_0 R^4}{8\pi\epsilon_0 r^3} (2 \cos \theta \hat{\mathbf{r}} + \sin \theta \hat{\boldsymbol{\theta}})}.$$

**Problem 3.36**

$\mathbf{p} = (\mathbf{p} \cdot \hat{\mathbf{r}}) \hat{\mathbf{r}} + (\mathbf{p} \cdot \hat{\boldsymbol{\theta}}) \hat{\boldsymbol{\theta}} = p \cos \theta \hat{\mathbf{r}} - p \sin \theta \hat{\boldsymbol{\theta}}$  (Fig. 3.36). So  $3(\mathbf{p} \cdot \hat{\mathbf{r}}) \hat{\mathbf{r}} - \mathbf{p} = 3p \cos \theta \hat{\mathbf{r}} - p \cos \theta \hat{\mathbf{r}} + p \sin \theta \hat{\boldsymbol{\theta}} = 2p \cos \theta \hat{\mathbf{r}} + p \sin \theta \hat{\boldsymbol{\theta}}$ . So Eq. 3.104  $\equiv$  Eq. 3.103.  $\checkmark$

**Problem 3.37**

$$V_{\text{ave}}(R) = \frac{1}{4\pi R^2} \int V(\mathbf{r}) da, \text{ where the integral is over the surface of a sphere of radius } R. \text{ Now } da =$$

$R^2 \sin \theta d\theta d\phi$ , so  $V_{\text{ave}}(R) = \frac{1}{4\pi} \int V(R, \theta, \phi) \sin \theta d\theta d\phi$ .

$$\begin{aligned} \frac{dV_{\text{ave}}}{dR} &= \frac{1}{4\pi} \int \frac{\partial V}{\partial R} \sin \theta d\theta d\phi = \frac{1}{4\pi} \int (\nabla V \cdot \hat{\mathbf{r}}) \sin \theta d\theta d\phi = \frac{1}{4\pi R^2} \int (\nabla V) \cdot (R^2 \sin \theta d\theta d\phi \hat{\mathbf{r}}) \\ &= \frac{1}{4\pi R^2} \int (\nabla V) \cdot d\mathbf{a} = \frac{1}{4\pi R^2} \int (\nabla^2 V) d\tau = 0. \end{aligned}$$

(The final integral, from the divergence theorem, is over the *volume* of the sphere, where by assumption the Laplacian of  $V$  is zero.) So  $V_{\text{ave}}$  is independent of  $R$ —the same for all spheres, regardless of their radius—and hence (taking the limit as  $R \rightarrow 0$ ),  $V_{\text{ave}}(R) = V(0)$ .  $\text{qed}$


**Problem 3.38** At a point  $(x, y)$  on the plane the field of  $q$  is

$$\mathbf{E}_q = \frac{1}{4\pi\epsilon_0} \frac{q}{z^3} \hat{\mathbf{z}}, \quad \text{and} \quad \mathbf{r} = x \hat{\mathbf{x}} + y \hat{\mathbf{y}} - d \hat{\mathbf{z}},$$

so its  $z$  component is  $-\frac{q}{4\pi\epsilon_0} \frac{d}{(x^2 + y^2 + d^2)^{3/2}}$ . Meanwhile, the field of  $\sigma$  (just below the surface) is  $-\frac{\sigma}{2\epsilon_0}$ , (Eq. 2.17). (Of course, this is for a *uniform* surface charge, but as long as we are infinitesimally far away  $\sigma$  is *e-effectively* uniform.) The total field inside the conductor is zero, so

$$-\frac{q}{4\pi\epsilon_0} \frac{d}{(x^2 + y^2 + d^2)^{3/2}} - \frac{\sigma}{2\epsilon_0} = 0 \quad \Rightarrow \quad \sigma(x, y) = -\frac{qd}{2\pi(x^2 + y^2 + d^2)^{3/2}}. \quad \checkmark$$

**Problem 3.39**



The image configuration is shown in the figure; the positive image charge forces cancel in pairs. The net force of the negative image charges is:

$$\begin{aligned} F &= \frac{1}{4\pi\epsilon_0} q^2 \left\{ \frac{1}{[2(a-x)]^2} + \frac{1}{[2a+2(a-x)]^2} + \frac{1}{[4a+2(a-x)]^2} + \dots \right. \\ &\quad \left. - \frac{1}{(2x)^2} - \frac{1}{(2a+2x)^2} - \frac{1}{(4a+2x)^2} - \dots \right\} \\ &= \frac{1}{4\pi\epsilon_0} \frac{q^2}{4} \left\{ \left[ \frac{1}{(a-x)^2} + \frac{1}{(2a-x)^2} + \frac{1}{(3a-x)^2} + \dots \right] - \left[ \frac{1}{x^2} + \frac{1}{(a+x)^2} + \frac{1}{(2a+x)^2} + \dots \right] \right\}. \end{aligned}$$


When  $a \rightarrow \infty$  (i.e.  $a \gg x$ ) only the  $\frac{1}{x^2}$  term survives:  $F = -\frac{1}{4\pi\epsilon_0} \frac{q^2}{(2x)^2} \checkmark$  (same as for only *one* plane—Eq. 3.12). When  $x = a/2$ ,

$$F = \frac{1}{4\pi\epsilon_0} \frac{q^2}{4} \left\{ \left[ \frac{1}{(a/2)^2} + \frac{1}{(3a/2)^2} + \frac{1}{(5a/2)^2} + \dots \right] - \left[ \frac{1}{(a/2)^2} + \frac{1}{(3a/2)^2} + \frac{1}{(5a/2)^2} + \dots \right] \right\} = 0. \quad \checkmark$$

**Problem 3.40**

Following Prob. 2.52, we place image line charges  $-\lambda$  at  $y = b$  and  $+\lambda$  at  $y = -b$  (here  $y$  is the horizontal axis,  $z$  vertical).





In the solution to Prob. 2.52 substitute:

$$a \rightarrow \frac{a-b}{2}, \quad y_0 \rightarrow \frac{a+b}{2} \quad \text{so} \quad \left(\frac{a-b}{2}\right)^2 = \left(\frac{a+b}{2}\right)^2 - R^2 \Rightarrow b = \frac{R^2}{a}.$$

$$\begin{aligned} V &= \frac{\lambda}{4\pi\epsilon_0} \left[ \ln\left(\frac{s_3^2}{s_4^2}\right) + \ln\left(\frac{s_1^2}{s_2^2}\right) \right] = \frac{\lambda}{4\pi\epsilon_0} \ln\left(\frac{s_1^2 s_3^2}{s_2^2 s_4^2}\right) \\ &= \frac{\lambda}{4\pi\epsilon_0} \ln \left\{ \frac{[(y+a)^2 + z^2][(y-b)^2 + z^2]}{[(y-a)^2 + z^2][(y+b)^2 + z^2]} \right\}, \quad \text{or, using } y = s \cos \phi, \quad z = s \sin \phi, \\ &= \boxed{\frac{\lambda}{4\pi\epsilon_0} \ln \left\{ \frac{(s^2 + a^2 + 2as \cos \phi)[(as/R)^2 + R^2 - 2as \cos \phi]}{(a^2 + a^2 - 2as \cos \phi)[(as/R)^2 + R^2 + 2as \cos \phi]} \right\}}. \end{aligned}$$

**Problem 3.41** Same as Problem 3.9, only this time we want  $q' + q'' = q$ , so  $q'' = q - q'$ :

$$F = \frac{q}{4\pi\epsilon_0} \left( \frac{q''}{a^2} + \frac{q'}{(a-b)^2} \right) = \frac{q^2}{4\pi\epsilon_0 a^2} + \frac{qq'}{4\pi\epsilon_0} \left( -\frac{1}{a^2} + \frac{1}{(a-b)^2} \right).$$

The second term is identical to Problem 3.9, and I'll just quote the answer from there:

$$F = \frac{q^2}{4\pi\epsilon_0 a^3} \left[ a - R^3 \frac{(2a^2 - R^2)}{(a^2 - R^2)^2} \right].$$


(a)  $F = 0 \Rightarrow a(a^2 - R^2)^2 = R^3(2a^2 - R^2)$ , or (letting  $x \equiv a/R$ ),  $x(x^2 - 1)^2 - 2x^2 + 1 = 0$ . We want a real root greater than 1; Mathematica delivers  $x = (1 + \sqrt{5})/2 = 1.61803$ , so  $a = 1.61803R = \boxed{5.66311 \text{ \AA}}$ .

(b) Let  $a_0 = x_0 R$  be the minimum value of  $a$ . The work necessary is


$$\begin{aligned} W &= - \int_{\infty}^{a_0} F da = \frac{q^2}{4\pi\epsilon_0} \int_{a_0}^{\infty} \frac{1}{a^3} \left[ a - R^3 \frac{(2a^2 - R^2)}{(a^2 - R^2)^2} \right] da = \frac{q^2}{4\pi\epsilon_0 R} \int_{x_0}^{\infty} \left[ \frac{1}{x^2} - \frac{(2x^2 - 1)}{x^3(x^2 - 1)^2} \right] dx \\ &= \frac{q^2}{4\pi\epsilon_0 R} \left[ \frac{1 + 2x_0 - 2x_0^3}{2x_0^2(1 - x_0^2)} \right]. \end{aligned}$$

Putting in  $x_0 = (1 + \sqrt{5})/2$ , Mathematica says the term in square brackets is  $1/2$  (this is not an accident; see footnote 6 on page 127), so  $W = \frac{q^2}{8\pi\epsilon_0 R}$ . Numerically,


$$W = \frac{(1.60 \times 10^{-19})^2}{8\pi(8.85 \times 10^{-12})(5.66 \times 10^{-10})} \text{ J} = 2.03 \times 10^{-19} \text{ J} = \boxed{1.27 \text{ eV}}.$$

**Problem 3.42**

The first configuration on the right is precisely Example 3.4, but unfortunately the second configuration is not the same as Problem 3.15:




We could reconstruct Problem 3.15 with the modified boundaries, but let's see if we can't twist it around by an astute change of variables. Suppose we let  $x \rightarrow y$ ,  $y \rightarrow u$ ,  $a \rightarrow c$ ,  $b \rightarrow a$ , and  $V_0 \rightarrow V_1$ :



This is closer; making the changes in the solution to Problem 3.15 we have (for this configuration)

$$V(u, y) = \frac{4V_1}{\pi} \sum_{n=1,3,5,\dots} \frac{\sinh(n\pi y/c) \sin(n\pi u/c)}{n \sinh(n\pi a/c)}.$$

Now let  $c \rightarrow 2b$  and  $u \rightarrow x + b$ , and the configuration is just what we want:



The potential for this configuration is

$$V(x, y) = \frac{4V_1}{\pi} \sum_{n=1,3,5,\dots} \frac{\sinh(n\pi y/2b) \sin(n\pi(x+b)/2b)}{n \sinh(n\pi a/2b)}.$$